

Local zeta regularization and the scalar Casimir effect I. A general approach based on integral kernels

Davide Fermi^a, Livio Pizzocchero^{b(1)}

^a Dipartimento di Matematica, Università di Milano
Via C. Saldini 50, I-20133 Milano, Italy
e-mail: davide.fermi@unimi.it

^b Dipartimento di Matematica, Università di Milano
Via C. Saldini 50, I-20133 Milano, Italy
and Istituto Nazionale di Fisica Nucleare, Sezione di Milano, Italy
e-mail: livio.pizzocchero@unimi.it

Abstract

This is the first one of a series of papers about zeta regularization of the divergences appearing in the vacuum expectation value (VEV) of several local and global observables in quantum field theory. More precisely we consider a quantized, neutral scalar field on a domain in any spatial dimension, with arbitrary boundary conditions and, possibly, in presence of an external classical potential. We analyze, in particular, the VEV of the stress-energy tensor, the corresponding boundary forces and the total energy, thus taking into account both local and global aspects of the Casimir effect. In comparison with the wide existing literature on these subjects, we try to develop a more systematic approach, allowing to treat specific configurations by mere application of a general machinery. The present Part I is mainly devoted to setting up this general framework; at the end of the paper, this is exemplified in a very simple case. In Parts II, III and IV we will consider more engaging applications, indicated in the Introduction of the present work.

Keywords: Local Casimir effect, renormalization, zeta regularization.

AMS Subject classifications: 81T55, 83C47.

PACS: 03.70.+k, 11.10.Gh, 41.20.Cv .

¹Corresponding author

Contents

1	Introduction	3
2	Zeta regularization for a scalar field	6
2.1	General setting.	6
2.2	Zeta regularization.	8
2.3	A remark.	11
2.4	Staticity features of the VEV of $\hat{T}_{\mu\nu}$	11
2.5	Conformal and non-conformal parts of the stress-energy tensor. . . .	12
2.6	Total energy and pressure on the boundary.	12
3	Expressions of the zeta regularized stress-energy VEV in terms of integral kernels	13
3.2	Basics on integral kernels.	13
3.3	The operator $\mathcal{A}$	14
3.4	The Green function.	15
3.5	A digression on complex powers.	16
3.6	The Dirichlet kernel.	16
3.7	Some remarks concerning the Dirichlet kernel and its derivatives. . .	18
3.8	The regularized propagator and stress-energy VEV: connections with the Dirichlet kernel.	18
3.9	The heat kernel, the cylinder kernel and some variations.	22
3.10	The case of a non-negative $\mathcal{A}$. The heat and cylinder kernels. . . .	24
3.11	Connections between the cylinder kernel and a $(d+1)$ -dimensional Green function.	25
3.12	Behaviour of the heat and cylinder kernels for small and large $\mathbf{t}$	26
3.13	The Dirichlet kernel as the Mellin transform of the heat or cylinder kernel.	28
3.14	Analytic continuation of Mellin transforms via integration by parts. .	30
3.15	Analytic continuation of Mellin transforms via complex integration. .	32
3.16	Other kernels, and their relations with $\mathbf{D}_s$	35
3.17	The case of product domains. Factorization of the heat kernel. . . .	35
3.18	The case of a slab: reduction to a lower-dimensional problem. . . .	37
4	Total energy and forces on the boundary	38
4.1	The total energy.	38
4.1.1	Reduced energy for a slab configuration.	41
4.2	Pressure on the boundary.	42
4.3	Explicit expression for the (regularized) pressure.	43

4.4	An equivalent characterization of boundary forces.	43
4.5	Integrated forces on the boundary.	45
4.6	A comment on some previous “anomalies”.	46
5	Some variations of the previous schemes	46
5.1	The basic Hilbert space when $\mathbf{0}$ is an isolated point of $\sigma(\mathcal{A})$; the case of Neumann and periodic boundary conditions.	47
5.2	The case where $\mathbf{0}$ is in the continuous spectrum of $\mathcal{A}$	48
5.3	Some variations involving the spatial domain.	52
6	The case of a massless field on the segment	53
6.1	Introducing the problem for arbitrary boundary conditions.	53
6.2	Cylinder and Dirichlet kernels.	55
6.3	The stress-energy tensor.	56
6.4	The total energy.	57
6.5	The boundary forces.	57
6.6	Dirichlet boundary conditions.	58
6.7	Dirichlet-Neumann boundary conditions.	59
6.8	Neumann boundary conditions.	61
6.9	Periodic boundary conditions.	62
A	Appendix. On the form (2.11) for the stress-energy tensor	63
B	Appendix. Smoothness properties of some integral kernels	66
C	Appendix. A $(d+1)$-dimensional Green function and its relation with the cylinder kernel	73
D	Appendix. Derivation of Eq. (3.90)	75
E	Appendix. Derivation of Eq.s (3.112-3.114)	76
F	Appendix. Some results on boundary forces	78
F.1	Derivation of Eq. (4.25) for the pressure.	78
F.2	Derivation of Eq. (4.29).	79
G	Appendix. Derivation of Eq. (5.21)	80

1 Introduction

As well known, zeta regularization treats divergent quantities in quantum field theory introducing a complex parameter, with the role of a regulator, and defining the renormalized observables in terms of analytic continuation with respect to the regulator. This construction was first proposed by Dowker and Critchley [30], Hawking [58] and Wald [101] to renormalize local observables, such as the vacuum expectation value (VEV) of the stress-energy tensor; the ultimate purpose was the semiclassical treatment of quantum effects in general relativity (e.g., using the stress-energy VEV as a source in Einstein's equations). After the previously cited pioneers, a number of authors championed the zeta approach to treat local observables; let us mention, in particular, Cognola, Zerbini, Elizalde [25, 26], and Moretti [17, 60, 75, 76, 77, 78] who worked in curved spacetimes, and Actor, Svaiter et al. [2, 3, 4, 86, 87] who worked on spatial domains with boundaries in flat spacetime.

The zeta strategy can be as well applied to global observables, such as the VEV of the total energy; in this version, it has in fact become more popular than its local counterpart. The literature on global zeta regularization is enormous; here we only cite the classical papers [88, 89] by Zimmerman et al., [13] by Blau, Visser, and Wipf and the monographies of Elizalde et al. [17, 33, 34], of Bordag et al. [14, 15] and of Kirsten [62] (see also [21, 44, 45, 57, 66, 63]).

Both in the local and in the global version, zeta methods provide a very natural approach to study the effects of quantum vacuum; in comparison with the original derivation of these effects by Casimir [20], and with other methods such as point-splitting [12, 16, 72] (in particular, see [55, 76] for a comparison between point-splitting and the zeta regularization approach) and the algebraic, microlocal approach (see [8, 27, 46, 81, 83] and citations therein), the zeta strategy is competitive and, perhaps, more elegant.

The present series of papers (formed by this work and by the subsequent Parts II,III,IV [38, 39, 40]) considers both the local and the global zeta approach, developing the viewpoint proposed in a special case by our previous work [36]. We are mainly interested in the theory of a quantized neutral scalar field on flat Minkowski spacetime, of arbitrary dimension $d+1$. The field is confined within a d -dimensional spatial domain Ω and arbitrary boundary conditions are prescribed for it on the boundary $\partial\Omega$; besides, we also admit the presence of a classical external potential, which could be meant to describe, in some sense, an effective theory of interacting fields. (We will only occasionally mention the possibility of replacing the flat domain Ω with a Riemannian manifold, or an open subset of it with prescribed boundary conditions; this amounts to replace Minkowski spacetime with a non-flat ultrastatic spacetime [17].)

Our attention is mainly focused on the VEV of the stress-energy tensor, of which we consider both the conformal and the non-conformal parts; however, we also

determine the total energy and boundary forces.

Most of the works on local zeta regularization cited above consider Euclidean quantum fields; this makes a difference with respect to our formalism, based on canonical quantization in a genuinely Lorentzian framework. Moreover, each one of the previously mentioned works [2, 3, 4, 36, 86, 87] about local zeta regularization for flat domains deals with a specific spatial configuration (e.g. a strip between parallel planes, configurations with perpendicular planes, a rectangular wave guide or a rectangular box). In the present series of papers we are trying to develop a more systematic approach to the zeta strategy, to be applied nearly automatically in each specific case.

In order to set up the general formalism we use with more generality the following fact, emerging from the previous literature: the analytic continuations involved in the zeta approach are closely related to certain integral kernels determined by the basic elliptic operator $\mathcal{A}$ which governs the spatial part of the field equation; among these kernels one could mention, in particular, the Dirichlet and heat kernels corresponding, respectively, to complex powers of $\mathcal{A}$ and to the exponential $e^{-t\mathcal{A}}$. Our papers I-IV emphasize as far as possible the basic role of these and other kernels in view of zeta regularization. Our aim is to write down few very general rules to construct the required analytic continuation via integral kernels; these prescriptions can be applied in an almost mechanical way to treat specific configurations, as shown by several applications.

The present Part I is mainly devoted to the general formulation, and in particular to the set of rules mentioned before; at the end of this paper we discuss, as a first application, the very simple case of a massless field on a segment (i.e., in spatial dimension $d = 1$) for several types of boundary conditions. In the subsequent Part II [38] we will show how our general schemes work in a number of explicitly solvable cases (the field between parallel or perpendicular planes, or inside a wedge, with arbitrary boundary conditions and arbitrary choices of the spatial dimension and of the conformal parameter). In Parts III [39] and IV [40] we will consider two cases in which the implementation of our general rules requires a mixture of analytic and numerical calculations; more precisely, Part III will discuss a field in presence of a background harmonic potential and Part IV a field confined within a rectangular box. In all these cases we will illustrate connections with the past literature including, when they exist, previous computations based on alternative approaches such as point-splitting.

Let us describe in more detail the contents of the present Part I. In Section 2 and in the related Appendix A, we introduce the general framework for a neutral scalar field on a d -dimensional spatial domain Ω , including a brief discussion on the stress-energy tensor with its conformal and non-conformal parts. This is an occasion to fix the attention on the basic elliptic operator $\mathcal{A} := -\Delta + V(\mathbf{x})$ on Ω , where Δ is the Laplacian and $V(\mathbf{x})$ the external potential. Always in Section

2, we introduce zeta regularization in the formulation already used in our previous work [36]; the basic idea is to replace the quantized field $\hat{\phi}$ with its regularized version $\hat{\phi}^u := (\mathcal{A}/\kappa^2)^{-u/4}\hat{\phi}$, where $u \in \mathbb{C}$ is the regulator and $\kappa > 0$ is a mass parameter. Formally, $\hat{\phi}^u$ becomes $\hat{\phi}$ for $u = 0$; the zeta approach implements this idea in terms of analytic continuation. This means the following: for any local or global observable (say, the VEV of either the stress-energy tensor or of the total energy), after introducing a regularized version of the observable based on $\hat{\phi}^u$, we define the renormalized value as the analytic continuation at $u = 0$. We distinguish between two versions of this prescription: the restricted zeta approach, in which the observable is analytic at $u = 0$, and the extended approach, where a singularity appears at $u = 0$ and is eliminated removing the negative powers of u in the Laurent expansion.

In Section 3 (and in the related Appendices C, D and E) we present a number of integral kernels associated to the operator $\mathcal{A} := -\Delta + V(\mathbf{x})$. The Dirichlet kernel $D_s(\mathbf{x}, \mathbf{y}) := \langle \delta_{\mathbf{x}} | \mathcal{A}^{-s} \delta_{\mathbf{y}} \rangle$ (s in a complex domain, $\mathbf{x}, \mathbf{y} \in \Omega$) is closely related to the stress-energy VEV. More precisely, the regularized stress-energy VEV, built from $\hat{\phi}^u$, is determined by the Dirichlet kernel $D_s(\mathbf{x}, \mathbf{y})$ (and by its spatial derivatives) at $\mathbf{y} = \mathbf{x}$, $s = (u \pm 1)/2$; so, the renormalized version of this VEV is determined by the analytic continuation of D_s near $s = \pm 1/2$. As shown in the cited section, the continuation of the Dirichlet kernel can be determined algorithmically relating it to the heat kernel $K(\mathbf{t}; \mathbf{x}, \mathbf{y}) := \langle \delta_{\mathbf{x}} | e^{-\mathbf{t}\mathcal{A}} \delta_{\mathbf{y}} \rangle$ or to other kernels (among them the so-called cylinder kernel $T(\mathbf{t}; \mathbf{x}, \mathbf{y}) := \langle \delta_{\mathbf{x}} | e^{-\mathbf{t}\sqrt{\mathcal{A}}} \delta_{\mathbf{y}} \rangle$, considered by Fulling [35, 47]). This procedure ultimately gives the set of mechanical rules for zeta regularization mentioned previously in this Introduction.

In Section 4 and in Appendix F we relate to the previous framework the total energy, the pressure and the total forces on the boundary. We also take the occasion to prove (at the regularized level) the equivalence between two alternative definitions of the pressure, often assumed without proof in the literature: pressure as the action of the stress-energy VEV on the normal to the boundary, and pressure as the functional derivative of the bulk energy with respect to deformations of the spatial domain Ω . In Section 5 and in Appendix G we consider some variations of the general framework of Sections 2 and 3 concerning the following situations: i) the case where 0 is either an isolated or non-isolated point of the spectrum of the fundamental operator $\mathcal{A} := -\Delta + V(\mathbf{x})$; ii) the case where the flat spatial domain Ω is described via curvilinear coordinates or, more generally, the case where Ω is a (possibly non-flat) Riemannian manifold equipped with arbitrary coordinates (or an open subset of it, with prescribed boundary conditions). This suggests the possibility to apply our formalism to the case of a non-flat ultrastatic spacetime where the line element reads $ds^2 = -dt^2 + d\ell^2$, with $d\ell^2$ the Riemannian line element of Ω .

The final Section 6 presents a first application of our formalism; this concerns the

case in which $d = 1$ and Ω is the interval $(0, a)$, with suitable boundary conditions. This configuration is very simple: we take it in consideration just to show in action our mechanical procedures for analytic continuation. Comparison with the existing literature on this simple case is performed. As already mentioned, more sophisticated applications of our approach will appear in Parts II-IV.

2 Zeta regularization for a scalar field

2.1 General setting. In this section we summarize the zeta regularization method for the propagator and the vacuum expectation value (VEV) of the stress-energy tensor of a quantized scalar field, in the formulation presented in [36] (see also [37]). Here the scheme of [36] is slightly generalized, admitting the presence of a classical background potential V and arbitrary spacetime dimensions. Throughout the paper we use natural units, so that

$$c = 1, \quad \hbar = 1. \quad (2.1)$$

We work in $(d + 1)$ -dimensional Minkowski spacetime; this is identified with $\mathbf{R}^{d+1}$ using a set of inertial coordinates

$$x = (x^\mu)_{\mu=0,1,\dots,d} \equiv (x^0, \mathbf{x}) \equiv (t, \mathbf{x}), \quad (2.2)$$

in which the Minkowski metric has coefficients $(\eta_{\mu\nu}) = \text{diag}(-1, 1, \dots, 1)$. Let us fix a spatial domain $\Omega \subset \mathbf{R}^d$ where we consider a quantized neutral, scalar field $\hat{\phi}$ in presence of a classical background static potential V ; so, we have $V : \Omega \rightarrow \mathbf{R}$, $\mathbf{x} \mapsto V(\mathbf{x})$ and

$$\hat{\phi} : \mathbf{R} \times \Omega \rightarrow \mathcal{L}_{sa}(\mathfrak{F}) ; \quad 0 = (-\partial_{tt} + \Delta - V(\mathbf{x}))\hat{\phi}(\mathbf{x}, t), \quad (2.3)$$

(analogous settings are considered, e.g., in [14, 15, 17, 46]) ⁽²⁾. Here we are referring to the space $\mathcal{L}(\mathfrak{F})$ of linear operators on the Fock space $\mathfrak{F}$, and to the subset $\mathcal{L}_{sa}(\mathfrak{F})$ of selfadjoint operators; $\Delta := \sum_{i=1}^d \partial_{ii}$ is the d -dimensional Laplacian. We assume V to be smooth, and prescribe appropriate boundary conditions (e.g., the Dirichlet conditions $\hat{\phi}(t, \mathbf{x}) = 0$ for $\mathbf{x} \in \partial\Omega$). From here to the end of the paper we put

$$\mathcal{A} := -\Delta + V, \quad (2.4)$$

intending that the boundary conditions are accounted for in the above definition; we assume this framework to grant that $\mathcal{A}$ is a selfadjoint operator in the Hilbert

² Of course the notation $\hat{\phi} : \mathbf{R} \times \Omega \rightarrow \mathcal{L}_{sa}(\mathfrak{F})$, $(\mathbf{x}, t) \mapsto \hat{\phi}(\mathbf{x}, t)$ is used here in connection with a *generalized* operator valued function; in fact, as well known, $\hat{\phi}$ is an operator valued *distribution*. In the papers of this series we use the classical functional notations even for generalized functions.

space $L^2(\Omega)$, with the inner product

$$\langle f|g \rangle := \int_{\Omega} d\mathbf{x} \bar{f}(\mathbf{x}) g(\mathbf{x}) \quad (2.5)$$

($d\mathbf{x}$ denoting the standard Lebesgue measure on Ω) ⁽³⁾. Moreover, we assume $\mathcal{A}$ to be *strictly positive*, by which we mean that the *spectrum* $\sigma(\mathcal{A})$ is contained in $[\varepsilon^2, +\infty)$ for some $\varepsilon > 0$. Let us mention that $\mathcal{A}$ may have continuous spectrum, a fact typically occurring when Ω is unbounded; therefore, when we speak of the eigenvectors of $\mathcal{A}$ we always intend them in a generalized sense, including improper eigenfunctions.

Let us consider a complete orthonormal set $(F_k)_{k \in \mathcal{K}}$ of (generalized) eigenfunctions of $\mathcal{A}$ ⁽⁴⁾, indexed by an unspecified set of labels $\mathcal{K}$, and let us write the corresponding eigenvalues in the form $(\omega_k^2)_{k \in \mathcal{K}}$ ($\omega_k \geq \varepsilon$ for all $k \in \mathcal{K}$). Thus

$$\begin{aligned} F_k : \Omega &\rightarrow \mathbf{C}; & \mathcal{A}F_k &= \omega_k^2 F_k; \\ \langle F_k | F_h \rangle &= \delta(k, h) & \text{for all } k, h \in \mathcal{K}. \end{aligned} \quad (2.6)$$

Any eigenfunction label $k \in \mathcal{K}$ can include different parameters, both discrete and continuous. Besides, we generically write $\int_{\mathcal{K}} dk$ to indicate summation over all labels (i.e., literal summation over discrete parameters and integration over continuous parameters, with respect to a suitable measure); $\delta(h, k) = \delta(k, h)$ is the Dirac delta function for the label space $\mathcal{K}$ (this reduces to the Kröneckers symbol in the case of discrete parameters). Note that the condition $\omega_k \geq \varepsilon > 0$ excludes the presence of infrared divergences from all the sums over k appearing in the sequel.

The functions

$$f_k : \mathbf{R} \times \Omega \rightarrow \mathbf{C}, \quad x = (t, \mathbf{x}) \mapsto f_k(x) := e^{-i\omega_k t} F_k(\mathbf{x}) \quad (2.7)$$

fulfill $(-\partial_{tt} - \mathcal{A})f_k = 0$; they allow us to infer for the quantized field a normal modes expansion of the form

$$\widehat{\phi}(x) = \int_{\mathcal{K}} \frac{dk}{\sqrt{2\omega_k}} \left[\widehat{a}_k f_k(x) + \widehat{a}_k^\dagger \overline{f_k}(x) \right] \quad (2.8)$$

³In passing, we recall that the Hilbert space $\mathfrak{F}$ can be realized as the direct sum of all symmetrized tensor powers of $L^2(\Omega)$.

⁴For a fully rigorous discussion of generalized eigenfunctions, see Chapter IV of [52]. In the sequel, following the usual terminology, when speaking of functions (or distributions) on Ω we use the adjectives “proper” or “improper”, to distinguish between objects which actually belong to $L^2(\Omega)$ or not. In this spirit we speak of proper or improper eigenfunctions, and use the same terminology for the corresponding eigenvalues. In the sequel, the adjective “generalized” in relation to eigenfunctions is sometimes omitted.

(with † indicating the adjoint operator, and $\bar{}$ the complex conjugate). In the above we are considering the destruction and creation operators $\hat{a}_k, \hat{a}_k^\dagger \in \mathcal{L}(\mathfrak{F})$, which fulfill the canonical commutation relations

$$[\hat{a}_k, \hat{a}_h] = 0, \quad [\hat{a}_k, \hat{a}_h^\dagger] = \delta(h, k), \quad \hat{a}_k|0\rangle = 0, \quad (2.9)$$

where $|0\rangle \in \mathfrak{F}$ is the vacuum state (of unit norm). A relevant character for the sequel of our analysis will be the *propagator*, i.e., the vacuum expectation value (VEV)

$$\langle 0 | \hat{\phi}(x) \hat{\phi}(y) | 0 \rangle \quad (x, y \in \mathbf{R} \times \Omega). \quad (2.10)$$

Let us pass to the stress-energy tensor operator; this depends on a parameter $\xi \in \mathbf{R}$, and its components $\hat{T}_{\mu\nu} : \mathbf{R} \times \Omega \rightarrow \mathcal{L}_{sa}(\mathfrak{F})$, for $\mu, \nu \in \{0, 1, \dots, d\}$, are given by

$$\hat{T}_{\mu\nu} := (1 - 2\xi) \partial_\mu \hat{\phi} \circ \partial_\nu \hat{\phi} - \left(\frac{1}{2} - 2\xi \right) \eta_{\mu\nu} (\partial^\lambda \hat{\phi} \partial_\lambda \hat{\phi} + V \hat{\phi}^2) - 2\xi \hat{\phi} \circ \partial_{\mu\nu} \hat{\phi} \quad (2.11)$$

where we use the symmetrized operator product $\hat{A} \circ \hat{B} := (1/2)(\hat{A}\hat{B} + \hat{B}\hat{A})$ and all the bilinear terms in the field are evaluated on the diagonal (e.g., $\partial_\mu \hat{\phi} \circ \partial_\nu \hat{\phi}$ indicates the map $x \mapsto \partial_\mu \hat{\phi}(x) \circ \partial_\nu \hat{\phi}(x)$).

Eq. (2.11) provides a natural quantization of what is often called the “improved” stress-energy tensor; this is a well-known modification of the canonical stress-energy tensor with an additive term proportional to the parameter ξ , that does not alter its divergence. For certain boundary conditions, e.g. of Dirichlet type, this addition does not even alter the corresponding momentum vector. We refer to Appendix A for further details on this topic. Here we only recall that the improved stress-energy tensor was first proposed by Callan, Coleman and Jackiw [19] in order to deal with some pathologies appearing in perturbation theory; later on, this tensor was reinterpreted in terms of the Minkowskian limit for a scalar field coupled to gravity via the curvature scalar [12, 17, 30, 79, 81].

Needless to say, the vacuum expectation value of the stress-energy tensor (2.11) is the main character in the theory of the Casimir effect. It is evident from Eq. (2.11) that $\langle 0 | \hat{T}_{\mu\nu}(x) | 0 \rangle$ can be expressed formally in terms of the propagator (2.10) (and of its derivatives) evaluated on the diagonal $y = x$. On the other hand, the propagator is known to be plagued with ultraviolet divergences along the diagonal, so that $\langle 0 | \hat{T}_{\mu\nu}(x) | 0 \rangle$ is a merely formal expression for a divergent quantity; our purpose is to redefine the propagator and the stress-energy VEV via a suitable regularization, ultimately yielding finite values for these quantities.

2.2 Zeta regularization. Let $\kappa > 0$ denote a parameter, to which we attribute the dimension of a mass (or of an inverse length, since $\hbar = 1$). κ will be called the mass scale; it is introduced for dimensional reasons and plays the role of a normalization scale. See [13, 17, 33, 60, 75] for further comments regarding this parameter and its presence or absence in the renormalized observables of the field.

We will check that the final, renormalized results depend on κ only when singularities appear in the analytic continuations involved in the following construction.

The zeta strategy, in the version proposed in [36] to give meaning to the VEV of $\widehat{T}_{\mu\nu}$, relies on the powers

$$(\kappa^{-2}\mathcal{A})^{-u/4} = \kappa^{u/2}\mathcal{A}^{-u/4} \quad (2.12)$$

where $\mathcal{A}$ is the operator (2.4) and $u \in \mathbf{C}$; these are employed to define the *smeared*, or *zeta-regularized, field operator*

$$\widehat{\phi}^u := (\kappa^{-2}\mathcal{A})^{-u/4} \widehat{\phi}, \quad (2.13)$$

depending on the complex parameter u and coinciding with the usual field operator $\widehat{\phi}$ for $u = 0$. If $(F_k)_{k \in \mathcal{K}}$ is a complete orthonormal set of eigenfunctions of $\mathcal{A}$ with eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$, we have $(\kappa^{-2}\mathcal{A})^{-u/4} F_k = \kappa^{u/2} \omega_k^{-u/2} F_k$, for any $k \in \mathcal{K}$; so, the functions $f_k(t, \mathbf{x}) = e^{-i\omega_k t} F_k(\mathbf{x})$ fulfill $(\kappa^{-2}\mathcal{A})^{-u/4} f_k = \kappa^{u/2} \omega_k^{-u/2} f_k$, and the expansion (2.8) for $\widehat{\phi}$ becomes, after application of $(\kappa^{-2}\mathcal{A})^{-u/4}$,

$$\widehat{\phi}^u(x) = \kappa^{u/2} \int_{\mathcal{K}} \frac{dk}{\sqrt{2} \omega_k^{1/2+u/2}} \left[\widehat{a}_k f_k(x) + \widehat{a}_k^\dagger \overline{f_k}(x) \right]. \quad (2.14)$$

Note that, in the limit $\omega_k \rightarrow +\infty$, the term $1/\omega_k^{1/2+u/2}$ in the above integral vanishes rapidly if $\Re u$ is large; this is a manifestation of the regularizing effect of the operator $(\kappa^{-2}\mathcal{A})^{-u/4}$ for large $\Re u$, a fact we will describe much more precisely in the sequel. Using $\widehat{\phi}^u$, we can define a *regularized propagator*

$$\langle 0 | \widehat{\phi}^u(x) \widehat{\phi}^u(y) | 0 \rangle \quad (x, y \in \mathbf{R} \times \Omega) \quad (2.15)$$

and a *zeta regularized stress-energy tensor*

$$\widehat{T}_{\mu\nu}^u := (1-2\xi) \partial_\mu \widehat{\phi}^u \circ \partial_\nu \widehat{\phi}^u - \left(\frac{1}{2} - 2\xi \right) \eta_{\mu\nu} \left(\partial^\lambda \widehat{\phi}^u \partial_\lambda \widehat{\phi}^u + V(\widehat{\phi}^u)^2 \right) - 2\xi \widehat{\phi}^u \circ \partial_{\mu\nu} \widehat{\phi}^u, \quad (2.16)$$

where, as in Eq. (2.11), all the bilinear terms in the field are evaluated on the diagonal.

We are interested in the VEV of this regularized stress-energy tensor, which formally gives $\langle 0 | \widehat{T}_{\mu\nu}^u(x) | 0 \rangle$ in the limit $u \rightarrow 0$. Of course, we can relate the VEV of $\widehat{T}_{\mu\nu}^u(x)$ to the regularized propagator (2.15) in the following way:

$$\begin{aligned} \langle 0 | \widehat{T}_{\mu\nu}^u(x) | 0 \rangle &= \\ &= \left(\frac{1}{2} - \xi \right) (\partial_{x^\mu y^\nu} + \partial_{x^\nu y^\mu}) - \left(\frac{1}{2} - 2\xi \right) \eta_{\mu\nu} \left(\partial^{x^\lambda} \partial_{y^\lambda} + V(\mathbf{x}) \right) - \xi (\partial_{x^\mu x^\nu} + \partial_{y^\mu y^\nu}) \Big|_{y=x} \cdot \\ &\quad \cdot \langle 0 | \widehat{\phi}^u(x) \widehat{\phi}^u(y) | 0 \rangle. \end{aligned} \quad (2.17)$$

We will return later on this equation and on its use for the actual computation of the above VEV.

Typically, the regularized propagator and the VEV of $\hat{T}_{\mu\nu}^u(x)$ are analytic functions of u , for $\Re u$ sufficiently large; the same can be said of many related observables (including global object, such as the total energy, which is related to the space integral of the $(0,0)$ component of the stress-energy tensor). Let us consider any one of these (local or global) observables, and denote with $\mathcal{F}(u)$ its zeta-regularized version, based on Eq. (2.13) (see, e.g., Eq. (2.15) or Eq. (2.16)); we assume that the function $u \mapsto \mathcal{F}(u)$ is well defined and analytic for u in a suitable domain $\mathcal{U}_0$ of the complex plane. The zeta approach to renormalization can be formulated in either a “restricted” or an “extended” version, both described hereafter.

i) *Zeta approach, restricted version.* Assume that the function $\mathcal{U}_0 \rightarrow \mathbf{C}, u \mapsto \mathcal{F}(u)$ can be analytically continued to a larger open subset $\mathcal{U}$ of $\mathbf{C}$ such that $0 \in \mathcal{U}$; let us use the notation $u \mapsto \mathcal{F}(u)$ even for this extension ⁽⁵⁾. In this case, making reference to the analytic continuation, we define the renormalized value of the observable under consideration as

$$\mathcal{F}_{ren} := \mathcal{F}(0) . \quad (2.18)$$

ii) *Zeta approach, extended version.* Assume that there is an open subset $\mathcal{U}$ of $\mathbf{C}$, larger than $\mathcal{U}_0$, such that $0 \in \mathcal{U}$ and the function $u \in \mathcal{U}_0 \mapsto \mathcal{F}(u)$ has an analytic continuation to $\mathcal{U} \setminus \{0\}$, still indicated with $\mathcal{F}$. In this case, since there is an isolated singularity at $u = 0$, in a neighborhood of this point we have the Laurent expansion $\mathcal{F}(u) = \sum_{k=-\infty}^{+\infty} \mathcal{F}_k u^k$. Let us consider the *regular part*

$$(RP \mathcal{F})(u) := \sum_{k=0}^{+\infty} \mathcal{F}_k u^k ; \quad (2.19)$$

we define the renormalized value of the given observable as

$$\mathcal{F}_{ren} := (RP \mathcal{F})(0) \quad (2.20)$$

(i.e., $\mathcal{F}_{ren} = \mathcal{F}_0$). In most applications $\mathcal{F}$ is *meromorphic* close to $u = 0$, which means that it has a *pole* at this point; in this case the previous Laurent expansion has the form $\mathcal{F}(u) = \sum_{k=-N}^{+\infty} \mathcal{F}_k u^k$, where $N \in \{1, 2, 3, \dots\}$ is the order of the pole. Let us stress that the prescription (2.20) is a quite straightforward generalization of the approach considered in [13, 33], where $\mathcal{F}$ was assumed to possess a simple pole in $u = 0$ (i.e., it was assumed that $N = 1$).

⁵In the style of our previous work [36] we should write $\mathcal{F} : \mathcal{U}_0 \rightarrow \mathbf{C}$ for the initially given function and $AC \mathcal{F} : \mathcal{U} \rightarrow \mathbf{C}$ for its analytic continuation; here, we choose to simplify the notation, writing $\mathcal{F}$ for both functions. Note that the analogue of Eq. (2.18) in the style of [36] would be $\mathcal{F}_{ren} := (AC \mathcal{F})(0)$.

Of course, the restricted zeta approach of item (i) is equivalent to a special case of the extended approach, in which $\mathcal{F}(u)$ has a removable singularity at $u = 0$ and the Laurent expansion at this point is the usual power series expansion.

A large part of our subsequent work will be devoted to the application of the previous scheme to the VEV of the stress-energy tensor. In general, we define the renormalized version of the latter as

$$\langle 0|\widehat{T}_{\mu\nu}(x)|0\rangle_{ren} := RP\Big|_{u=0} \langle 0|\widehat{T}_{\mu\nu}^u(x)|0\rangle ; \quad (2.21)$$

when no singularity appears at $u = 0$, according to the restricted approach (i) the above definition reduces to

$$\langle 0|\widehat{T}_{\mu\nu}(x)|0\rangle_{ren} := \langle 0|\widehat{T}_{\mu\nu}^u(x)|0\rangle\Big|_{u=0} . \quad (2.22)$$

In [36], we only considered the prescription (2.22) in the special case of a Dirichlet field (with $V = 0$) between two parallel planes, i.e., in the configuration corresponding to the standard theory of the Casimir effect. In that case the approach (2.22) was implemented via a direct computation of the analytic continuation appearing therein; as already stressed, here we aim to much more generality.

2.3 A remark. In the sequel, while performing zeta regularization and the consequent renormalization, it is sometimes natural to consider, in place of u , some complex parameter s related to u by a simple transformation. In view of such situations, it is convenient to generalize some notations of the previous subsection in the following way:

- i) Consider an analytic function $\mathcal{S}_0 \rightarrow \mathbf{C}$, $s \mapsto \mathcal{F}(s)$, where $\mathcal{S}_0$ is an open subset of $\mathbf{C}$; if this admits an analytic continuation to a larger open subset $\mathcal{S}$, the latter will be still denoted with $s \mapsto \mathcal{F}(s)$.
- ii) Suppose the analytic function $\mathcal{S}_0 \rightarrow \mathbf{C}$, $s \mapsto \mathcal{F}(s)$ has an analytic extension to $\mathcal{S} \setminus \{s_0\}$, where $\mathcal{S}$ is an open subset of $\mathbf{C}$ and $s_0 \in \mathcal{S}$. Then, the Laurent expansion $\mathcal{F}(s) = \sum_{k=-\infty}^{+\infty} \mathcal{F}_k(s-s_0)^k$ will be used to define the regular part (near s_0) of this analytic continuation as $(RP \mathcal{F})(s) := \sum_{k=0}^{+\infty} \mathcal{F}_k(s-s_0)^k$; of course, this implies $(RP \mathcal{F})(s_0) = \mathcal{F}_0$.

2.4 Staticity features of the VEV of $\widehat{T}_{\mu\nu}$. Let us return to the regularized stress-energy tensor; for $x = (t, \mathbf{x}) \in \mathbf{R} \times \Omega$, we claim that

$$\begin{aligned} \langle 0|\widehat{T}_{\mu\nu}^u(x)|0\rangle & \text{ is independent of } t , \\ \langle 0|\widehat{T}_{0i}^u(x)|0\rangle & = \langle 0|\widehat{T}_{i0}^u(x)|0\rangle = 0 \quad \text{ for } i \in \{1, \dots, d\} . \end{aligned} \quad (2.23)$$

These statements are not surprising, due to the staticity of the general framework considered in the present paper; a formal proof will be given in subsection 3.8 (see Eq.s (3.31-3.33) and the considerations which follow them). Of course the features of Eq. (2.23) are preserved by analytic continuation, so that an analogue of this equation holds for the renormalized VEV $\langle 0|\widehat{T}_{\mu\nu}(x)|0\rangle_{ren}$ as well.

2.5 Conformal and non-conformal parts of the stress-energy tensor. In the literature (see, e.g., [12, 17, 102]) it is customary to write the stress-energy tensor (here to be intended as one of the operators $\hat{T}_{\mu\nu}$, $\hat{T}_{\mu\nu}^u$, or either one of the VEVs $\langle 0|\hat{T}_{\mu\nu}^u|0\rangle$, $\langle 0|\hat{T}_{\mu\nu}|0\rangle_{ren}$) as the sum of a *conformal* and a *non-conformal* part. In order to define these quantities, let us consider for ξ the critical value

$$\xi_d := \frac{d-1}{4d} ; \quad (2.24)$$

it is known that, when coupling of the scalar field to gravity is taken into account, the theory is invariant (for $V = 0$) under conformal transformations of the spacetime line element if ξ has the above critical value (see, e.g., [102], p.447). In the sequel we adopt the notations

$$\diamond \equiv \text{conformal} , \quad \blacksquare \equiv \text{non-conformal} \quad (2.25)$$

and we define, for example, the conformal and non-conformal parts of the renormalized stress-energy VEV $\langle 0|\hat{T}_{\mu\nu}|0\rangle_{ren}$ in the following way:

$$\langle 0|\hat{T}_{\mu\nu}^{(\diamond)}|0\rangle_{ren} := \langle 0|\hat{T}_{\mu\nu}|0\rangle_{ren} \Big|_{\xi=\xi_d} , \quad (2.26)$$

$$\langle 0|\hat{T}_{\mu\nu}^{(\blacksquare)}|0\rangle_{ren} := \frac{1}{\xi - \xi_d} \left(\langle 0|\hat{T}_{\mu\nu}|0\rangle_{ren} - \langle 0|\hat{T}_{\mu\nu}^{(\diamond)}|0\rangle_{ren} \right) . \quad (2.27)$$

Of course, this implies

$$\langle 0|\hat{T}_{\mu\nu}|0\rangle_{ren} = \langle 0|\hat{T}_{\mu\nu}^{(\diamond)}|0\rangle_{ren} + (\xi - \xi_d) \langle 0|\hat{T}_{\mu\nu}^{(\blacksquare)}|0\rangle_{ren} . \quad (2.28)$$

In the applications to be considered in Section 6 and in the subsequent Parts II,III and IV, when presenting our final results for the renormalized stress-energy VEV, we will either write them in the form (2.28) or give separately the conformal and non-conformal parts (2.26) (2.27).

2.6 Total energy and pressure on the boundary. The *total energy* is, by definition, the integral of $\langle 0|\hat{T}_{00}(x)|0\rangle$ over the spatial domain Ω . We defer the discussion of this topic to Section 4; therein we will describe the representation of the total energy as the sum of a bulk term and a boundary term, in the framework of zeta regularization.

In the same section we will use zeta regularization to treat the *pressure* on the boundary $\partial\Omega$ of the spatial domain; this quantity can be defined in terms of the VEV of the spatial components $\hat{T}_{ij}$. There is an alternative characterization of the pressure in terms of the variation of the bulk energy (see Eq. (4.6) for the definition) with respect to deformations of the spatial domain Ω ; the equivalence of this definition with the previous one has often been assumed uncritically in the literature, so we think it can be useful to produce a formal proof (see Section 4).

3 Expressions of the zeta regularized stress-energy VEV in terms of integral kernels

In this section Ω always denotes a spatial domain in $\mathbf{R}^d$, and we often consider the Hilbert space $L^2(\Omega)$ of the square integrable complex-valued functions on Ω .

3.2 Basics on integral kernels. Let us consider a linear operator $\mathcal{B}$ acting on $L^2(\Omega)$. The *integral kernel* of $\mathcal{B}$ is the (generalized) function

$$\mathcal{B}(\cdot, \cdot) : \Omega \times \Omega \rightarrow \mathbf{C}, \quad (\mathbf{x}, \mathbf{y}) \mapsto \mathcal{B}(\mathbf{x}, \mathbf{y}) := \langle \delta_{\mathbf{x}} | \mathcal{B} \delta_{\mathbf{y}} \rangle \quad (3.1)$$

where $\delta_{\mathbf{x}}$ and $\delta_{\mathbf{y}}$ are the Dirac delta functions centered at $\mathbf{x}$ and $\mathbf{y}$, respectively, here viewed as improper vectors of the Hilbert space $L^2(\Omega)$ ⁽⁶⁾. Equivalently, the integral kernel of the operator $\mathcal{B}$ can be defined as the unique (generalized) function $\mathcal{B}(\cdot, \cdot) : \Omega \times \Omega \rightarrow \mathbf{C}$ such that

$$(\mathcal{B}\psi)(\mathbf{x}) = \int_{\Omega} d\mathbf{y} \mathcal{B}(\mathbf{x}, \mathbf{y}) \psi(\mathbf{y}), \quad (3.2)$$

for all sufficiently regular $\psi : \Omega \rightarrow \mathbf{C}$. If $\mathcal{B}$ possesses a complete orthonormal set of (generalized) eigenfunctions $(F_k)_{k \in \mathcal{K}}$ with corresponding eigenvalues $\beta_k \in \mathbf{C}$

⁶Let us repeat what we have declared in the footnote of page 6: in these papers we use functional notations even for objects which are not ordinary functions. Making reference to the theory of Schwartz distributions, we can understand the statement “ $\mathcal{B}(\cdot, \cdot) : \Omega \times \Omega \rightarrow \mathbf{C}$ ” as a way to indicate that we are considering a (complex) distribution on $\Omega \times \Omega$; let us describe how to intend Eq. (3.1) from this viewpoint. We refer to the space of test functions $D(\Omega)$, formed by the C^∞ , compactly supported functions $\varphi : \Omega \rightarrow \mathbf{C}$ and equipped with its standard inductive limit topology; $D(\Omega \times \Omega)$ has a similar meaning. We wish to define rigorously $(\mathbf{x}, \mathbf{y}) \mapsto \langle \delta_{\mathbf{x}} | \mathcal{B} \delta_{\mathbf{y}} \rangle$ as a distribution on $\Omega \times \Omega$, i.e., as a continuous linear form on $D(\Omega \times \Omega)$. To this purpose we consider $\varphi, \psi \in D(\Omega)$ and note that the identities $\varphi = \int_{\Omega} d\mathbf{x} \varphi(\mathbf{x}) \delta_{\mathbf{x}}$ and $\psi = \int_{\Omega} d\mathbf{x} \psi(\mathbf{x}) \delta_{\mathbf{x}}$ give formally

$$\int_{\Omega \times \Omega} d\mathbf{x} d\mathbf{y} \overline{\varphi}(\mathbf{x}) \langle \delta_{\mathbf{x}} | \mathcal{B} \delta_{\mathbf{y}} \rangle \psi(\mathbf{y}) = \langle \varphi | \mathcal{B} \psi \rangle.$$

On the other hand, if the domain of the operator $\mathcal{B}$ contains $D(\Omega)$ and the sesquilinear map $D(\Omega) \times D(\Omega) \rightarrow \mathbf{C}$, $(\psi, \varphi) \mapsto \langle \varphi | \mathcal{B} \psi \rangle$ is continuous, using the nuclear theorem of Schwartz (see [92], Thm. II) one proves rigorously the existence of a unique distribution on $\Omega \times \Omega$, denoted with $(\mathbf{x}, \mathbf{y}) \mapsto \langle \delta_{\mathbf{x}} | \mathcal{B} \delta_{\mathbf{y}} \rangle$, such that the above relation holds for all $\varphi, \psi \in D(\Omega)$, intending the integral in the left hand side as the action of this distribution on the test function $(\mathbf{x}, \mathbf{y}) \mapsto \overline{\varphi}(\mathbf{x}) \psi(\mathbf{y})$. One can use similar considerations to give a distributional meaning to Eq. (3.2) (see again [92]). Throughout the papers of this series, whenever we speak of the integral kernel of an operator $\mathcal{B}$ we implicitly assume $\mathcal{B}$ to possess the regularity features mentioned before, so that $\langle \delta_{\mathbf{x}} | \mathcal{B} \delta_{\mathbf{y}} \rangle$ makes sense at least as a distribution. In some cases described explicitly in the sequel, stronger assumptions on $\mathcal{B}$ ensure that $\langle \delta_{\mathbf{x}} | \mathcal{B} \delta_{\mathbf{y}} \rangle$ can be understood as an ordinary function, continuous with its derivatives up to a certain order; this is the situation outlined in Appendix B, often mentioned in the sequel in relation to several kernels of interest for us.

$(\mathcal{B}F_k = \beta_k F_k)$, then

$$\mathcal{B}(\mathbf{x}, \mathbf{y}) = \int_{\mathcal{K}} dk \beta_k F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) \quad (3.3)$$

(since the function in the right-hand side fulfills equation (3.2) for all ψ). The precise sense in which the eigenfunction expansion (3.3) converges depends on the specific features of the operator $\mathcal{B}$; we generally assume distributional convergence⁽⁷⁾, leaving to future subsections the consideration of special cases where convergence can be intended in a stronger sense.

Incidentally, let us mention the relation existing between the kernel $\mathcal{B}(\cdot, \cdot)$ and the trace of $\mathcal{B}$; the latter, if it exists, is the number $\text{Tr } \mathcal{B} := \int_{\mathcal{K}} dk \langle F_k | \mathcal{B} F_k \rangle \in \mathbf{C}$, where $(F_k)_{k \in \mathcal{K}}$ is any complete orthonormal set of $L^2(\Omega)$. The right-hand side does not depend on the choice of $(F_k)_{k \in \mathcal{K}}$; in particular, if $\mathcal{B}$ has purely discrete spectrum, $(F_k)_{k \in \mathcal{K}}$ is a complete orthonormal set of proper eigenfunctions labelled by a countable set $\mathcal{K}$ and $(\beta_k)_{k \in \mathcal{K}}$ are the corresponding eigenvalues, we have $\text{Tr } \mathcal{B} = \sum_{k \in \mathcal{K}} \beta_k$, if this series converges. Returning to definition (3.1) of the kernel $\mathcal{B}(\cdot, \cdot)$, we see that

$$\text{Tr } \mathcal{B} = \int_{\Omega} d\mathbf{x} \mathcal{B}(\mathbf{x}, \mathbf{x}) \quad (3.4)$$

(since $(\delta_{\mathbf{x}})_{\mathbf{x} \in \Omega}$ is a generalized complete orthonormal set)⁽⁸⁾.

Let us move on and note that the boundary conditions possibly involved in the definition of $\mathcal{B}$ have implications for the kernel $\mathcal{B}(\cdot, \cdot)$; for example, if boundary conditions of the Dirichlet type are involved, the eigenfunctions $(F_k)_{k \in \mathcal{K}}$ in Eq. (3.3) vanish on $\partial\Omega$, thus yielding $\mathcal{B}(\mathbf{x}, \mathbf{y}) = 0$ for $\mathbf{x} \in \partial\Omega$ or $\mathbf{y} \in \partial\Omega$.

Let us also mention that from Eq. (3.1) one infers

$$\mathcal{B}^\dagger(\mathbf{x}, \mathbf{y}) = \overline{\mathcal{B}(\mathbf{y}, \mathbf{x})}, \quad \overline{\mathcal{B}}(\mathbf{x}, \mathbf{y}) = \overline{\mathcal{B}(\mathbf{x}, \mathbf{y})} \quad (3.5)$$

where $\mathcal{B}^\dagger$ is the adjoint operator of $\mathcal{B}$ with respect to the inner product of $L^2(\Omega)$ while $\overline{\mathcal{B}}$ is the complex conjugate operator, such that $\overline{\mathcal{B}\psi} = \overline{\mathcal{B}}\overline{\psi}$ for all ψ . These facts imply

$$\mathcal{B}(\mathbf{y}, \mathbf{x}) = \mathcal{B}(\mathbf{x}, \mathbf{y}) \quad \text{if } \mathcal{B}^\dagger = \overline{\mathcal{B}}. \quad (3.6)$$

3.3 The operator $\mathcal{A}$. Most of the integral kernels considered in the sequel will be related to a selfadjoint operator $\mathcal{A}$ acting in $L^2(\Omega)$; for example, they will be the kernels associated to some function of the operator $\mathcal{A}$ (say, a power $\mathcal{A}^s$). To treat these kernels, precise assumptions on $\mathcal{A}$ will be specified whenever necessary; in any

⁷Meaning that $\int_{\Omega \times \Omega} d\mathbf{x} d\mathbf{y} \mathcal{B}(\mathbf{x}, \mathbf{y}) \varphi(\mathbf{x}, \mathbf{y}) = \int_{\mathcal{K}} dk \beta_k \int_{\Omega \times \Omega} d\mathbf{x} d\mathbf{y} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) \varphi(\mathbf{x}, \mathbf{y})$ for all test functions φ on $\Omega \times \Omega$.

⁸In the sequel, the adjective “generalized” in relation to complete orthonormal sets is sometimes omitted.

case, we will typically consider three situations of decreasing generality, described by the forthcoming Eqs (3.7)(3.8)(3.9).

In the first situation, we will simply assume that

$$\mathcal{A} \text{ is a strictly positive, selfadjoint operator in } L^2(\Omega) \quad (3.7)$$

(let us recall that strict positivity means $\sigma(\mathcal{A}) \subset [\varepsilon^2, +\infty)$ for some $\varepsilon > 0$).

In the second situation, the assumptions are the following:

$$\begin{aligned} \mathcal{A} &= -\Delta + V \text{ in } \Omega, \text{ with } V \text{ a real } C^\infty \text{ potential on } \Omega \text{ and} \\ &\text{boundary conditions such that } \mathcal{A} \text{ be selfadjoint and strictly positive.} \end{aligned} \quad (3.8)$$

In the third situation, the assumptions are:

$$\begin{aligned} \mathcal{A} &= -\Delta + V \text{ on a bounded domain } \Omega \text{ with } \partial\Omega \text{ of class } C^\infty \text{ and Dirichlet} \\ &\text{boundary conditions, } V \text{ a real } C^\infty \text{ potential on } \overline{\Omega} \text{ and } V(\mathbf{x}) \geq 0 \text{ for all } \mathbf{x} \in \overline{\Omega}. \end{aligned} \quad (3.9)$$

Here and in the sequel $\overline{\Omega} = \Omega \cup \partial\Omega$ is the closure of Ω . The assumptions (3.9) grant selfadjointness and strict positivity of $\mathcal{A}$; moreover, they imply that $\mathcal{A}$ has a purely discrete spectrum and one can build a complete orthonormal system of proper eigenfunctions F_k with eigenvalues ω_k^2 labelled by $\mathcal{K} = \{1, 2, 3, \dots\}$ in such a way that $0 < \omega_1 \leq \omega_2 \leq \omega_3 \leq \dots$ (with the possibility that some of these inequalities are equalities, to deal with the case of degenerate eigenvalues). It is well-known that the eigenvalues, when ordered in this manner, fulfill the Weyl asymptotic relation

$$\omega_k \sim C k^{1/d} \quad \text{for } k \rightarrow +\infty, \quad (3.10)$$

where $C := 2\sqrt{\pi} \Gamma(d/2 + 1)^{1/d} \text{Vol}(\Omega)^{-1/d}$ (see [71], Thm.5, page 189 and [32], § 8.2, pages 99-101 for elementary derivations). Appendix B contains a number of technical results concerning cases (3.7) (3.8) (3.9), that will be mentioned whenever necessary in the sequel of this section.

These results often refer to the spaces $C^j(\Omega)$ and $C^j(\Omega \times \Omega)$ or (in the case (3.9)) $C^j(\overline{\Omega})$ and $C^j(\overline{\Omega} \times \overline{\Omega})$, for $j \in \mathbf{N}$ or $j = \infty$; these are formed by the complex functions on $\Omega, \Omega \times \Omega$ and so on, which are continuous along with their partial derivatives of all orders $\leq j$. In particular, in the rest of the section we will present situations connected to cases (3.7) (3.8) (3.9), in which certain integral kernels $(\mathbf{x}, \mathbf{y}) \mapsto \mathcal{B}(\mathbf{x}, \mathbf{y})$ related to $\mathcal{A}$ are of class $C^j(\Omega \times \Omega)$ or $C^j(\overline{\Omega} \times \overline{\Omega})$, for suitable j .

The condition of strict positivity for $\mathcal{A}$ in Eqs (3.7) (3.8) will be occasionally relaxed, assuming only that the eigenvalues of $\mathcal{A}$ are non-negative (and declaring this explicitly, to avoid misunderstandings); see, e.g., subsection 3.10.

3.4 The Green function. Let $\mathcal{A}$ be a strictly positive selfadjoint operator in $L^2(\Omega)$; then, we can introduce the inverse operator $\mathcal{A}^{-1}$ and the corresponding kernel

$$G(\mathbf{x}, \mathbf{y}) := \mathcal{A}^{-1}(\mathbf{x}, \mathbf{y}), \quad (3.11)$$

which is called the *Green function* of $\mathcal{A}$. In terms of this kernel, the identity $\mathcal{A}\mathcal{A}^{-1} = \mathbf{1}$ can be re-expressed as

$$\mathcal{A}_{\mathbf{x}} G(\mathbf{x}, \mathbf{y}) = \delta(\mathbf{x} - \mathbf{y}) , \quad (3.12)$$

where $\mathcal{A}_{\mathbf{x}}$ indicates the operator $\mathcal{A}$ acting on $G(\mathbf{x}, \mathbf{y})$ as a function of $\mathbf{x}$. Using a complete orthonormal system $(F_k)_{k \in \mathcal{K}}$ of eigenfunctions of $\mathcal{A}$ with corresponding eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$, we can express the Green function as

$$G(\mathbf{x}, \mathbf{y}) = \int_{\mathcal{K}} \frac{dk}{\omega_k^2} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) . \quad (3.13)$$

The Green function is among the most familiar integral kernels, especially when $\mathcal{A} = -\Delta + V$ with suitable boundary conditions (of the Dirichlet, Neumann or Robin type); in this case, uniqueness results are available for the Poisson equation (perturbed with an external potential), allowing to characterize the Green function $G(\mathbf{x}, \mathbf{y})$ as the unique solution of Eq. (3.12) fulfilling the prescribed boundary conditions for $\mathbf{x} \in \partial\Omega$ or $\mathbf{y} \in \partial\Omega$. The literature on this topic is enormous, and here we only mention some well-known monographies: Berezanskii [9], Krylov [64], Sauvigny [90] and Shimakura [96] give abstract and rigorous analyses, while Duffy [31], Kytke [65], Sommerfeld [99] and Stakgold and Holst [100] present more practical and explicit discussions.

3.5 A digression on complex powers. Throughout the present paper (and in Parts II-IV), the following conventions are employed:

- i) $\ln : (0, +\infty) \rightarrow \mathbf{R}$ is the elementary logarithm;
- ii) for any $\alpha \in \mathbf{C}$, we systematically refer to the standard definition

$$x^\alpha := e^{\alpha \ln x} \quad \text{for all } x \in (0, +\infty) ; \quad (3.14)$$

- iii) for $\alpha \in \mathbf{C}$ and z in a convenient subset $\mathbf{C}^\times$ of the complex plane, we define

$$z^\alpha := e^{\alpha \ln |z| + i\alpha \arg z} , \quad (3.15)$$

where $\arg : \mathbf{C}^\times \rightarrow \mathbf{R}$ is some determination of the argument; this determination depends on the domain $\mathbf{C}^\times$ and must be specified in each case of interest. In most applications considered hereafter we set

$$\begin{aligned} \mathbf{C}^\times &:= \mathbf{C} \setminus [0, +\infty) ; \\ \arg &:= \text{the unique determination of the argument with values in } (0, 2\pi) . \end{aligned} \quad (3.16)$$

3.6 The Dirichlet kernel. Let again $\mathcal{A}$ be a strictly positive selfadjoint operator in $L^2(\Omega)$. The power $\mathcal{A}^{-s}$ can be defined through the standard functional calculus for each $s \in \mathbf{C}$; the corresponding integral kernel

$$D_s(\mathbf{x}, \mathbf{y}) := \mathcal{A}^{-s}(\mathbf{x}, \mathbf{y}) \quad (3.17)$$

is called the s -th *Dirichlet kernel*. In passing, let us note that $D_{-1}(\mathbf{x}, \mathbf{y})$ coincides with the Green function $G(\mathbf{x}, \mathbf{y})$ considered in subsection 3.4.

If $(F_k)_{k \in \mathcal{K}}$ is a complete orthonormal set of eigenfunctions of $\mathcal{A}$ with corresponding eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$ we have $\mathcal{A}^{-s} F_k = \omega_k^{-2s} F_k$, so that (by Eq. (3.3))

$$D_s(\mathbf{x}, \mathbf{y}) = \int_{\mathcal{K}} \frac{dk}{\omega_k^{2s}} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) . \quad (3.18)$$

The denomination of “Dirichlet kernel” employed for D_s is suggested by the similarity between the above expansion and the Dirichlet series, considered in [56, 73, 74, 95].

For our purposes, it is important to give sufficient conditions under which $D_s(\mathbf{x}, \mathbf{y})$ is a regular function of $(\mathbf{x}, \mathbf{y})$ (even on the diagonal $\mathbf{y} = \mathbf{x}$, of special interest in the sequel); we are also interested in cases in which the expansion (3.18) converges in a stronger sense, in comparison with the distributional sense that we are generically ascribing to kernel expansions. Let us present some results of this kind, which are proved in Appendix B.

Let $\mathcal{A} = -\Delta + V$ be as in Eq. (3.8); then

$$D_s \in C^j(\Omega \times \Omega) \quad \text{for } s \in \mathbf{C}, j \in \mathbf{N} \text{ such that } \Re s > \frac{d}{2} + \frac{j}{2} . \quad (3.19)$$

With the stronger assumptions (3.9), we have

$$D_s \in C^j(\overline{\Omega} \times \overline{\Omega}) \quad \text{for } s \in \mathbf{C}, j \in \mathbf{N} \text{ such that } \Re s > \frac{d}{2} + \frac{j}{2} . \quad (3.20)$$

Again with the assumptions (3.9), $\mathcal{A}$ possesses a complete orthonormal set of proper eigenfunctions $(F_k)_{k=1,2,3,\dots}$ (see the comments after the cited equation), and the expansion (3.18) reads

$$D_s(\mathbf{x}, \mathbf{y}) = \sum_{k=1}^{+\infty} \frac{1}{\omega_k^{2s}} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) . \quad (3.21)$$

In this case one has $F_k \in C^\infty(\overline{\Omega})$ for each k ; moreover, for $\Re s > d+j/2$ the expansion (3.21) is absolutely and uniformly convergent on $\overline{\Omega} \times \overline{\Omega}$, with all its derivatives up to order j : see Appendix B (especially Eq. (B.26)) for more details.

To conclude the present subsection let us mention that the general statement of Eq. (3.4), here applied with $\mathcal{B} = \mathcal{A}^{-s}$, yields

$$\int_{\Omega} d\mathbf{x} D_s(\mathbf{x}, \mathbf{x}) = \text{Tr } \mathcal{A}^{-s} , \quad (3.22)$$

provided that the above trace exists. With the assumptions (3.9) for $\mathcal{A} = -\Delta + V$, using the Weyl estimates (3.10) for $\omega_1 \leq \omega_2 \leq \dots$ we find

$$\mathrm{Tr} \mathcal{A}^{-s} = \sum_{k=1}^{+\infty} \frac{1}{\omega_k^{2s}} \quad \text{is finite for } \Re s > \frac{d}{2} . \quad (3.23)$$

3.7 Some remarks concerning the Dirichlet kernel and its derivatives.

Let us consider again a strictly positive selfadjoint operator $\mathcal{A}$ in $L^2(\Omega)$; moreover, assume this operator to be real, in the sense that $\overline{\mathcal{A}} = \mathcal{A}$ (i.e., $\overline{A\psi} = A\overline{\psi}$ for all ψ). If $(F_k)_{k \in \mathcal{K}}$ is a complete orthonormal set of eigenfunctions of $\mathcal{A}$ with related eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$, then the conjugate system $(\overline{F_k})_{k \in \mathcal{K}}$ is as well a complete orthonormal set of eigenfunctions of $\mathcal{A}$ with the same eigenvalues; so, besides Eq. (3.18) we have an alternative representation for the Dirichlet kernel $D_s(\mathbf{x}, \mathbf{y})$, based on this conjugate system. From here, we easily infer that, for $s \in \mathbf{C}$ with complex conjugate $\bar{s}$,

$$D_s(\mathbf{x}, \mathbf{y}) = D_s(\mathbf{y}, \mathbf{x}) \quad \text{and} \quad \overline{D_s(\mathbf{x}, \mathbf{y})} = D_{\bar{s}}(\mathbf{x}, \mathbf{y}) . \quad (3.24)$$

To go on we claim that, for any pair of multi-indexes α, β ,

$$\partial_{\mathbf{x}}^{\alpha} \partial_{\mathbf{y}}^{\beta} D_s(\mathbf{x}, \mathbf{y}) \big|_{\mathbf{y}=\mathbf{x}} = \partial_{\mathbf{x}}^{\beta} \partial_{\mathbf{y}}^{\alpha} D_s(\mathbf{x}, \mathbf{y}) \big|_{\mathbf{y}=\mathbf{x}} . \quad (3.25)$$

Indeed, due to the first identity in Eq. (3.24), we have $\partial_{\mathbf{x}}^{\alpha} \partial_{\mathbf{y}}^{\beta} D_s(\mathbf{x}, \mathbf{y}) = \partial_{\mathbf{x}}^{\alpha} \partial_{\mathbf{y}}^{\beta} D_s(\mathbf{y}, \mathbf{x})$; when evaluating the right-hand side of this equality on the diagonal $\mathbf{y} = \mathbf{x}$, the variables can be relabeled to yield $\partial_{\mathbf{x}}^{\alpha} \partial_{\mathbf{y}}^{\beta} D_s(\mathbf{y}, \mathbf{x}) \big|_{\mathbf{y}=\mathbf{x}} = \partial_{\mathbf{y}}^{\alpha} \partial_{\mathbf{x}}^{\beta} D_s(\mathbf{x}, \mathbf{y}) \big|_{\mathbf{x}=\mathbf{y}}$, thus proving Eq. (3.25).

All the above results can be applied to the (real) operator $\mathcal{A} := -\Delta + V(\mathbf{x})$; the symmetry properties outlined here for the corresponding Dirichlet kernel will be relevant in connection with the results of the next section.

3.8 The regularized propagator and stress-energy VEV: connections with the Dirichlet kernel.

Let us refer to the framework of the previous section, where the operator $A = -\Delta + V$ in $L^2(\Omega)$ has been considered in connection with a quantized scalar field. In the sequel $x = (x^0, \mathbf{x}), y = (y^0, \mathbf{y}) \in \mathbf{R} \times \Omega$; if we use the expansion (2.14) for the regularized field $\widehat{\phi}^u$ in terms of creation and destruction operators we obtain for the regularized propagator the expression

$$\begin{aligned} & \langle 0 | \widehat{\phi}^u(x) \widehat{\phi}^u(y) | 0 \rangle = \\ & = \kappa^u \int_{\mathcal{K} \times \mathcal{K}} \frac{dk dh}{2(\omega_k \omega_h)^{\frac{u+1}{2}}} \langle 0 | \left[\widehat{a}_k f_k(x) + \widehat{a}_k^{\dagger} \overline{f_k}(x) \right] \left[\widehat{a}_h f_h(y) + \widehat{a}_h^{\dagger} \overline{f_h}(y) \right] | 0 \rangle . \end{aligned} \quad (3.26)$$

This relation, along with the identities $\langle 0|\widehat{a}_k\widehat{a}_h|0\rangle = \langle 0|\widehat{a}_k^\dagger\widehat{a}_h^\dagger|0\rangle = \langle 0|\widehat{a}_k^\dagger\widehat{a}_h|0\rangle = 0$ and $\langle 0|\widehat{a}_k\widehat{a}_h^\dagger|0\rangle = \delta(k, h)$, gives

$$\begin{aligned}\langle 0|\widehat{\phi}^u(x)\widehat{\phi}^u(y)|0\rangle &= \kappa^u \int_{\mathcal{K}} \frac{dk}{2\omega_k^{1+u}} f_k(x)\overline{f_k}(y) = \\ &= \kappa^u \int_{\mathcal{K}} \frac{dk}{2\omega_k^{1+u}} F_k(\mathbf{x})\overline{F_k}(\mathbf{y}) e^{-i\omega_k(x^0-y^0)} .\end{aligned}\quad (3.27)$$

From here we can easily obtain the derivatives of the propagator; for example, for $j \in \{1, \dots, d\}$, we have

$$\partial_{x^0 y^j} \langle 0|\widehat{\phi}^u(x)\widehat{\phi}^u(y)|0\rangle = -i\kappa^u \int_{\mathcal{K}} \frac{dk}{2\omega_k^u} F_k(\mathbf{x})(\partial_{y^j}\overline{F_k})(\mathbf{y}) e^{-i\omega_k(x^0-y^0)} . \quad (3.28)$$

In particular, if we apply Eq.s (3.27) (3.28) with $y = x$ and compare with the eigenfunction expansion (3.18) of the Dirichlet kernel, we get

$$\langle 0|\widehat{\phi}^u(x)\widehat{\phi}^u(y)|0\rangle \Big|_{y=x} = \frac{\kappa^u}{2} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} ; \quad (3.29)$$

$$\partial_{x^0 y^j} \langle 0|\widehat{\phi}^u(x)\widehat{\phi}^u(y)|0\rangle \Big|_{y=x} = -i\kappa^u \partial_{y^j} D_{\frac{u}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} . \quad (3.30)$$

One can express similarly all the derivatives with $y = x$ appearing in Eq. (2.17) for the regularized stress-energy VEV. In this way (and using as well the identity (3.25)) we obtain the following results, where i, j, ℓ are spatial indexes ranging in $\{1, \dots, d\}$ ⁽⁹⁾:

$$\langle 0|\widehat{T}_{00}^u(t, \mathbf{x})|0\rangle = \kappa^u \left[\left(\frac{1}{4} + \xi \right) D_{\frac{u-1}{2}}(\mathbf{x}, \mathbf{y}) + \left(\frac{1}{4} - \xi \right) (\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x})) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right] \Big|_{\mathbf{y}=\mathbf{x}} , \quad (3.31)$$

$$\langle 0|\widehat{T}_{0j}^u(t, \mathbf{x})|0\rangle = \langle 0|\widehat{T}_{j0}^u(t, \mathbf{x})|0\rangle = 0 , \quad (3.32)$$

⁹To prove Eq. (3.32), note that

$$\langle 0|\widehat{T}_{0j}^u(t, \mathbf{x})|0\rangle = -\langle 0|\widehat{T}_{0j}^u(t, \mathbf{x})|0\rangle = -\frac{i\kappa^u}{2} \left(\partial_{y^j} D_{\frac{u}{2}}(\mathbf{x}, \mathbf{y}) - \partial_{x^j} D_{\frac{u}{2}}(\mathbf{x}, \mathbf{y}) \right) \Big|_{\mathbf{y}=\mathbf{x}}$$

and that the last expression vanishes due to identity (3.25). Besides, note that Eq. (3.33) is equivalent to the more symmetric expression

$$\begin{aligned}\langle 0|\widehat{T}_{ij}^u(t, \mathbf{x})|0\rangle &= \langle 0|\widehat{T}_{ji}^u(t, \mathbf{x})|0\rangle = \kappa^u \left[\left(\frac{1}{4} - \xi \right) \delta_{ij} \left(D_{\frac{u-1}{2}}(\mathbf{x}, \mathbf{y}) - (\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x})) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right) + \right. \\ &\quad \left. + \left(\left(\frac{1}{4} - \xi \right) (\partial_{x^i y^j} + \partial_{x^j y^i}) - \frac{\xi}{2} (\partial_{x^i x^j} + \partial_{y^i y^j}) \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right] \Big|_{\mathbf{y}=\mathbf{x}} .\end{aligned}$$

$$\begin{aligned}
\langle 0|\widehat{T}_{ij}^u(t, \mathbf{x})|0\rangle &= \langle 0|\widehat{T}_{ji}^u(t, \mathbf{x})|0\rangle = \\
&= \kappa^u \left[\left(\frac{1}{4} - \xi \right) \delta_{ij} \left(D_{\frac{u-1}{2}}(\mathbf{x}, \mathbf{y}) - (\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x})) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right) + \right. \\
&\quad \left. + \left(\left(\frac{1}{2} - \xi \right) \partial_{x^i y^j} - \xi \partial_{x^i x^j} \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}}.
\end{aligned} \tag{3.33}$$

The above equations indicate, amongst else, that $\langle 0|\widehat{T}_{\mu\nu}^u(t, \mathbf{x})|0\rangle$ does not depend on the time variable t ; this comes as no surprise at all, since our general framework is itself static (indeed, the spatial domain Ω and the potential V are time independent). These features of the regularized stress-energy VEV had been anticipated in subsection 2.4; due to them, in the rest of the paper we will use the notation ⁽¹⁰⁾

$$\langle 0|\widehat{T}_{\mu\nu}^u(\mathbf{x})|0\rangle \equiv \langle 0|\widehat{T}_{\mu\nu}^u(t, \mathbf{x})|0\rangle. \tag{3.34}$$

Up to now we have been working rather loosely, but it is not difficult to indicate the precise conditions for the validity of our manipulations. Indeed, Eq.s (3.31-3.33) for $\langle 0|\widehat{T}_{\mu\nu}^u(\mathbf{x})|0\rangle$ involve the Dirichlet kernels $D_{\frac{u-1}{2}}$ and $D_{\frac{u+1}{2}}$, with its second order derivatives, along the diagonal $\mathbf{y} = \mathbf{x}$. So, for these equations to be meaningful it is sufficient that $D_{\frac{u-1}{2}} \in C^0(\Omega \times \Omega)$ and $D_{\frac{u+1}{2}} \in C^2(\Omega \times \Omega)$; with the assumptions (3.8) on $\mathcal{A} = -\Delta + V$, recalling Eq. (3.19) we see that both conditions are fulfilled if

$$\Re u > d + 1. \tag{3.35}$$

For $\mathcal{A}$, u as in Eq.s (3.8) (3.35), the map $\mathbf{x} \mapsto \langle 0|\widehat{T}_{\mu\nu}^u(\mathbf{x})|0\rangle$ is in $C^0(\Omega)$ for all μ, ν . With the stronger assumptions (3.9) on $\mathcal{A}$, and again with u as in (3.35), we infer from (3.20) that $\mathbf{x} \mapsto \langle 0|\widehat{T}_{\mu\nu}^u(\mathbf{x})|0\rangle$ is in $C^0(\overline{\Omega})$ for all $\mu, \nu \in \{0, \dots, d\}$ (i.e., the regularized stress-energy VEV is continuous up to the boundary) ⁽¹¹⁾.

Now, let us fix a point $\mathbf{x} \in \Omega$ and, making the assumptions (3.8), consider the functions $u \mapsto \langle 0|\widehat{T}_{\mu\nu}^u(\mathbf{x})|0\rangle$; these are not only well defined, but even analytic on the half plane $\{u \in \mathbf{C} \mid \Re u > d + 1\}$ ⁽¹²⁾.

Let us recall that, according to Eq.s (2.18) (2.20), the analytic continuation of $\langle 0|\widehat{T}_{\mu\nu}^u(\mathbf{x})|0\rangle$ at $u = 0$ determines the zeta renormalized VEV of the stress-energy

¹⁰This is slightly abusive, since staticity occurs only *after* taking the VEV; the alternative notation $\langle 0|\widehat{T}_{\mu\nu}^u|0\rangle(\mathbf{x})$ is more precise, but graphically disturbing and will not be employed in the sequel.

¹¹Generalizing the previous considerations, one proves the following for any $\ell \in \mathbf{N}$: with the assumptions (3.8) (resp., (3.9)), if $\Re u > d + \ell + 1$ the function $\mathbf{x} \mapsto \langle 0|\widehat{T}_{\mu\nu}^u(\mathbf{x})|0\rangle$ is in $C^\ell(\Omega)$ (resp., in $C^\ell(\overline{\Omega})$).

¹²This fact can be established proving analyticity with respect to u of $D_{\frac{u-1}{2}}(\mathbf{x}, \mathbf{x})$, of $D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{x})$ and of the second order derivatives of $D_{\frac{u+1}{2}}$ at $(\mathbf{x}, \mathbf{x})$. We do not go into the details of the proofs, that will be given elsewhere [41] using rather obvious analyticity results for the operator functions $u \mapsto \mathcal{A}^{-(u \mp 1)/2}$.

tensor; of course, the latter does not depend on t as well and we will write

$$\langle 0|\widehat{T}_{\mu\nu}(\mathbf{x})|0\rangle_{ren} \equiv \langle 0|\widehat{T}_{\mu\nu}(t, \mathbf{x})|0\rangle_{ren} . \quad (3.36)$$

Of course, due to Eqs (3.31-3.33), the renormalized stress-energy VEV is determined by the “renormalized” functions

$$D_{\pm\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) := RP\Big|_{u=0} \left(\kappa^u D_{\frac{u\pm 1}{2}}(\mathbf{x}, \mathbf{y}) \right) , \quad (3.37)$$

$$\partial_{zw} D_{\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) := RP\Big|_{u=0} \left(\kappa^u \partial_{zw} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right) \quad (3.38)$$

(with z, w any two spatial variables), to be evaluated along the diagonal $\mathbf{y} = \mathbf{x}$. More precisely, we have

$$\langle 0|\widehat{T}_{00}(\mathbf{x})|0\rangle_{ren} = \left[\left(\frac{1}{4} + \xi \right) D_{-\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) + \left(\frac{1}{4} - \xi \right) (\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x})) D_{+\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} , \quad (3.39)$$

$$\langle 0|\widehat{T}_{0j}(\mathbf{x})|0\rangle_{ren} = \langle 0|\widehat{T}_{j0}(\mathbf{x})|0\rangle_{ren} = 0 , \quad (3.40)$$

$$\begin{aligned} \langle 0|\widehat{T}_{ij}(\mathbf{x})|0\rangle_{ren} &= \langle 0|\widehat{T}_{ji}(\mathbf{x})|0\rangle_{ren} = \\ &= \left[\left(\frac{1}{4} - \xi \right) \delta_{ij} \left(D_{-\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) - (\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x})) D_{+\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) \right) + \right. \\ &\quad \left. + \left(\left(\frac{1}{2} - \xi \right) \partial_{x^i y^j} - \xi \partial_{x^i x^j} \right) D_{+\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} . \end{aligned} \quad (3.41)$$

Let us remark that, if $D_{\frac{u\pm 1}{2}}(\mathbf{x}, \mathbf{y})$ and $\partial_{zw} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y})$ have analytic continuations regular at $u = 0$, indicated hereafter with $D_{\pm\frac{1}{2}}(\mathbf{x}, \mathbf{y})$ and $\partial_{zw} D_{\frac{1}{2}}(\mathbf{x}, \mathbf{y})$, one has

$$\begin{aligned} D_{\pm\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) &= D_{\pm\frac{1}{2}}(\mathbf{x}, \mathbf{y}) , \\ \partial_{zw} D_{\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) &= \partial_{zw} D_{\frac{1}{2}}(\mathbf{x}, \mathbf{y}) \end{aligned} \quad (3.42)$$

for any choice of the mass scale κ ; clearly, in this case the renormalized stress-energy VEV is independent of κ . On the contrary, an explicit dependence on κ appears if the analytic continuations of $D_{\frac{u\pm 1}{2}}(\mathbf{x}, \mathbf{x})$ or $\partial_{zw} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{x})$ (or both) have a singularity at $u = 0$; this will occur in some specific examples, to be considered in the subsequent Parts II and III.

A connection between a regularized stress-energy VEV and a Dirichlet-like kernel is mentioned by Cognola, Vanzo and Zerbini [24] in a slightly different framework, where the field theory becomes Euclidean after Wick rotation of the time coordinate, and our operator $\mathcal{A}$ (in the spatial variables $\mathbf{x}$) is replaced by the (spacetime) differential operator $-\partial_{tt} - \Delta + V$.

A variant of the previous results about the Dirichlet kernel and the regularized VEV $\langle 0 | \widehat{T}_{\mu\nu}^u(\mathbf{x}) | 0 \rangle$ can be formulated in the case of a *slab*. In this case $\Omega = \Omega_1 \times \mathbf{R}^{d_2}$, with Ω_1 a domain in $\mathbf{R}^{d_1}$ and $d_1 + d_2 = d$; moreover, the potential V depends only on the coordinates $\mathbf{x}_1 \in \Omega_1$. In this situation we can express $\langle 0 | \widehat{T}_{\mu\nu}^u(\mathbf{x}) | 0 \rangle$ in terms of the Dirichlet kernel associated to the operator $\mathcal{A}_1 := -\Delta_1 + V(\mathbf{x}_1)$ (we defer to subsection 3.18 a comprehensive discussion of slabs configurations).

In the following subsections we return to the case where Ω is an arbitrary domain in $\mathbf{R}^d$ and we connect the Dirichlet kernel D_s to other integral kernels, in order to shed light on the analytic continuation of D_s (of course, these connections will also be useful, in their d_1 -dimensional formulation, in the case of a slab).

3.9 The heat kernel, the cylinder kernel and some variations. Let us consider again a strictly positive selfadjoint operator $\mathcal{A}$ in $L^2(\Omega)$ (that will be $-\Delta + V$ in the subsequent applications). For all $\mathbf{t} \in [0, +\infty)$, using the standard functional calculus we can define the operators

$$e^{-\mathbf{t}\mathcal{A}}, \quad e^{-\mathbf{t}\sqrt{\mathcal{A}}} . \quad (3.43)$$

These fulfill the following conditions:

$$\left(\frac{d}{d\mathbf{t}} + \mathcal{A} \right) e^{-\mathbf{t}\mathcal{A}} = 0 , \quad e^{-\mathbf{t}\mathcal{A}}|_{\mathbf{t}=0} = \mathbf{1} , \quad (3.44)$$

$$\left(\frac{d^2}{d\mathbf{t}^2} - \mathcal{A} \right) e^{-\mathbf{t}\sqrt{\mathcal{A}}} = 0 , \quad e^{-\mathbf{t}\sqrt{\mathcal{A}}}|_{\mathbf{t}=0} = \mathbf{1} ; \quad (3.45)$$

moreover, due to the strict positivity of $\mathcal{A}$, $e^{-\mathbf{t}\mathcal{A}}$ and $e^{-\mathbf{t}\sqrt{\mathcal{A}}}$ are expected to vanish for $\mathbf{t} \rightarrow +\infty$, in some sense that can be made more precise in terms of integral kernels. Let us now pass to the kernels

$$K(\mathbf{t}; \mathbf{x}, \mathbf{y}) := e^{-\mathbf{t}\mathcal{A}}(\mathbf{x}, \mathbf{y}) , \quad T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = e^{-\mathbf{t}\sqrt{\mathcal{A}}}(\mathbf{x}, \mathbf{y}) , \quad (3.46)$$

which can be expressed as follows in terms of a complete orthonormal set $(F_k)_{k \in \mathcal{K}}$ of eigenfunctions of $\mathcal{A}$ and of the corresponding eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$:

$$K(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \int_{\mathcal{K}} dk \, e^{-\mathbf{t}\omega_k^2} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) , \quad (3.47)$$

$$T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \int_{\mathcal{K}} dk \, e^{-\mathbf{t}\omega_k} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) . \quad (3.48)$$

We note that

$$(\partial_{\mathbf{t}} + \mathcal{A}_{\mathbf{x}}) K(\mathbf{t}; \mathbf{x}, \mathbf{y}) = 0 , \quad K(0; \mathbf{x}, \mathbf{y}) = \delta(\mathbf{x} - \mathbf{y}) ; \quad (3.49)$$

$$(\partial_{\mathbf{t}} - \mathcal{A}_{\mathbf{x}}) T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = 0 , \quad T(0; \mathbf{x}, \mathbf{y}) = \delta(\mathbf{x} - \mathbf{y}) . \quad (3.50)$$

In the above $\mathcal{A}_{\mathbf{x}}$ indicates the operator $\mathcal{A}$ acting on $K(\mathbf{t}; \mathbf{x}, \mathbf{y})$ and $T(\mathbf{t}; \mathbf{x}, \mathbf{y})$ as functions of the $\mathbf{x}$ variable; Eq.s (3.49)(3.50) follow, respectively, from Eq.s (3.44) (3.45). Besides, under minimal supplementary conditions one can prove that $K(\mathbf{t}; \mathbf{x}, \mathbf{y})$ and $T(\mathbf{t}; \mathbf{x}, \mathbf{y})$ vanish exponentially for fixed $\mathbf{x}, \mathbf{y} \in \Omega$ and $\mathbf{t} \rightarrow +\infty$; we shall return on this in subsection 3.12.

In case (3.8) (i.e., $\mathcal{A} = -\Delta + V$ with V smooth) we have $\partial_{\mathbf{t}} + \mathcal{A}_{\mathbf{x}} = \partial_{\mathbf{t}} - \Delta_{\mathbf{x}} + V(\mathbf{x})$ and $\partial_{\mathbf{t}} - \mathcal{A}_{\mathbf{x}} = \partial_{\mathbf{t}} + \Delta_{\mathbf{x}} - V(\mathbf{x})$, so Eq. (3.49) contains a heat equation and Eq. (3.50) a $(d+1)$ -dimensional Laplace equation (with an external potential); note as well that $K(\mathbf{t}; \mathbf{x}, \mathbf{y})$ and $T(\mathbf{t}; \mathbf{x}, \mathbf{y})$ fulfill the boundary conditions in the definition of $\mathcal{A}$ for $\mathbf{x}$ or $\mathbf{y}$ in $\partial\Omega$. For obvious reasons, K is called the *heat kernel* of $\mathcal{A}$ (even in cases where $\mathcal{A}$ is not of the form $-\Delta+V$); T is called by Fulling [48] the *cylinder kernel* of $\mathcal{A}$. It should be mentioned that, under the assumptions (3.8), there are rigorous proofs that

$$(\mathbf{t}, \mathbf{x}, \mathbf{y}) \mapsto K(\mathbf{t}, \mathbf{x}, \mathbf{y}), (\mathbf{t}, \mathbf{x}, \mathbf{y}) \mapsto T(\mathbf{t}, \mathbf{x}, \mathbf{y}) \text{ are in } C^\infty((0, +\infty) \times \Omega \times \Omega); \quad (3.51)$$

with the stronger assumptions (3.9) there is a result similar to (3.51), with Ω replaced by $\bar{\Omega}$. (These results can be proved by a slight generalization of Thm. 5.2.1 in [28]. For some related statements, see also Appendix B and [41].)

Again with the assumptions (3.9), $\mathcal{A}$ has pure point spectrum and possesses a complete orthonormal set of proper eigenfunctions $(F_k)_{k=1,2,3,\dots}$, for which the expansions (3.47) (3.48) hold with $\int_{\mathcal{K}} dk = \sum_{k=1}^{+\infty}$. One can prove that, for each $\mathbf{t} > 0$, these expansions converge absolutely and uniformly on $\bar{\Omega} \times \bar{\Omega}$ with their derivatives of any order (see again Appendix B, especially Eq. (B.35)).

With the more general assumptions (3.8) we have $\mathcal{A}^\dagger = \bar{\mathcal{A}} = \mathcal{A}$, which in turn implies similar relations for $e^{-\mathbf{t}\mathcal{A}}$, $e^{-\mathbf{t}\sqrt{\mathcal{A}}}$; due to Eq. (3.6), this gives

$$K(\mathbf{t}; \mathbf{y}, \mathbf{x}) = K(\mathbf{t}; \mathbf{x}, \mathbf{y}), \quad T(\mathbf{t}; \mathbf{y}, \mathbf{x}) = T(\mathbf{t}; \mathbf{x}, \mathbf{y}). \quad (3.52)$$

Needless to say, the heat kernel has been the object of intensive and detailed studies, even in a much more general framework than the one considered in the present paper; exhaustive analyses have been given, for example, by Berline et al. [11], Calin et al. [18], Chavel [22], Davies [28], Gilkey [53] and Grigor'yan [54]. On the contrary, the cylinder kernel is a less popular object; it has mainly been investigated by Fulling and co-authors [47, 48, 50]. Some considerations of Fulling (see, e.g., [51]) also involve the operator $\sqrt{\mathcal{A}}^{-1} e^{-\mathbf{t}\sqrt{\mathcal{A}}}$ and the associated kernel

$$\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) := (\sqrt{\mathcal{A}}^{-1} e^{-\mathbf{t}\sqrt{\mathcal{A}}})(\mathbf{x}, \mathbf{y}) = \int_{\mathcal{K}} \frac{dk}{\omega_k} e^{-\mathbf{t}\omega_k} F_k(\mathbf{x}) \bar{F}_k(\mathbf{y}), \quad (3.53)$$

which we will refer to as the *modified cylinder kernel*, for reasons which become apparent hereafter (see Eq. (3.54)). Let us observe that the trivial relation $e^{-\mathbf{t}\sqrt{\mathcal{A}}} =$

$-\frac{d}{dt}(\sqrt{\mathcal{A}}^{-1}e^{-t\sqrt{\mathcal{A}}})$ can be reformulated in terms of integral kernels as

$$T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = -\partial_{\mathbf{t}}\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) ; \quad (3.54)$$

conversely, $\tilde{T}$ can be determined as the primitive of $-T$ which vanishes for $\mathbf{t} \rightarrow +\infty$, that is

$$\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \int_{\mathbf{t}}^{+\infty} d\mathbf{t}' T(\mathbf{t}'; \mathbf{x}, \mathbf{y}) . \quad (3.55)$$

In some cases $\tilde{T}$ is easier to compute than T , and some identities relating the cylinder kernel T to the Dirichlet kernel D_s can be applied more efficiently if they are rephrased in terms of $\tilde{T}$ (this situation will be exemplified in the case of a wedge domain, to be discussed in the subsequent Part II; see Section 5 therein).

With the assumptions (3.8) or (3.9), one could give for $\tilde{T}$ some regularity results very similar to the ones illustrated previously for T (e.g.: statement (3.51) holds as well for $\tilde{T}$ under the conditions (3.8)).

Before moving on, let us consider the *heat* and *cylinder traces*; these are respectively defined, for $\mathbf{t} \in (0, +\infty)$, as

$$K(\mathbf{t}) := \text{Tr } e^{-\mathbf{t}\mathcal{A}} , \quad T(\mathbf{t}) := \text{Tr } e^{-\mathbf{t}\sqrt{\mathcal{A}}} . \quad (3.56)$$

Assuming the above traces to exist, the general identity (3.4) for the trace of an operator $\mathcal{B}$ (here applied with either $\mathcal{B} = e^{-\mathbf{t}\mathcal{A}}$ or $\mathcal{B} = e^{-\mathbf{t}\sqrt{\mathcal{A}}}$) yields respectively

$$K(\mathbf{t}) = \int_{\Omega} d\mathbf{x} K(\mathbf{t}; \mathbf{x}, \mathbf{x}) , \quad T(\mathbf{t}) = \int_{\Omega} d\mathbf{x} T(\mathbf{t}; \mathbf{x}, \mathbf{x}) . \quad (3.57)$$

In particular, in the case (3.9) where the eigenvalues of $\mathcal{A}$ are labelled by $\mathcal{K} = \{1, 2, 3, \dots\}$ and the Weyl estimate (3.10) holds, we have:

$$K(\mathbf{t}) = \sum_{k=1}^{+\infty} e^{-\mathbf{t}\omega_k^2} < +\infty , \quad T(\mathbf{t}) = \sum_{k=1}^{+\infty} e^{-\mathbf{t}\omega_k} < +\infty \quad \text{for all } \mathbf{t} > 0 . \quad (3.58)$$

Clearly enough, an analogous discussion could be made for the space integral of the diagonal, modified cylinder kernel $\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{x})$; we omit this discussion for brevity.

3.10 The case of a non-negative $\mathcal{A}$. The heat and cylinder kernels. Let us remark that the heat and cylinder kernels can both be defined even in case $\mathcal{A}$ is *non-negative*, without requiring strict positivity; by this we mean that $\sigma(\mathcal{A}) \subset [0, +\infty)$, and that we are not assuming $\sigma(\mathcal{A}) \subset [\varepsilon^2, +\infty)$ for any $\varepsilon > 0$. The non-negativity of $\mathcal{A}$ is equivalent to the existence of a complete orthonormal system $(F_k)_{k \in \mathcal{K}}$ of (either proper or improper) eigenfunctions with corresponding non-negative eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$ ($\omega_k \geq 0$). Most of the considerations of the previous subsection still hold, in particular Eqs (3.47) (3.48).

For example, if $\mathcal{A} = -\Delta$ and $\Omega = \mathbf{R}^d$, then $\mathcal{A}$ is non-negative with eigenfunctions $F_{\mathbf{k}}(\mathbf{x}) = (2\pi)^{-d/2} e^{i\mathbf{k}\cdot\mathbf{x}}$ and eigenvalues $\omega_{\mathbf{k}}^2 = |\mathbf{k}|^2$, labelled by $\mathbf{k} \in \mathbf{R}^d$; the measure $d\mathbf{k}$ on the set of labels is the usual Lebesgue measure of $\mathbf{R}^d$. The eigenfunction expansion (3.47) of the heat kernel yields in this case the familiar result

$$K(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \frac{1}{(4\pi\mathbf{t})^{d/2}} e^{-\frac{|\mathbf{x}-\mathbf{y}|^2}{4\mathbf{t}}}; \quad (3.59)$$

moreover, the expansion (3.48) of the cylinder kernels gives the result

$$T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \frac{\Gamma(\frac{d+1}{2}) \mathbf{t}}{\pi^{\frac{d+1}{2}} (\mathbf{t}^2 + |\mathbf{x} - \mathbf{y}|^2)^{\frac{d+1}{2}}}, \quad (3.60)$$

which is a bit less popular and appears, e.g., in [49].

In some subcases with non-negative spectrum we can speak as well of the modified cylinder kernel $\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y})$. In fact, if 0 has zero spectral measure (a fact holding when 0 belongs to the continuous spectrum, but not holding when 0 is a proper eigenvalue), $\sqrt{\mathcal{A}}^{-1} e^{-\mathbf{t}\sqrt{\mathcal{A}}}$ can be defined through the standard functional calculus for selfadjoint operators. With minimal supplementary assumptions of regularity, the kernel $\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) := \langle \delta_{\mathbf{x}} | \sqrt{\mathcal{A}}^{-1} e^{-\mathbf{t}\sqrt{\mathcal{A}}} \delta_{\mathbf{y}} \rangle$ makes sense, as well as its expansion (3.53) in terms of a complete orthonormal set of (generalized) eigenfunctions F_k with eigenvalues ω_k^2 ($k \in \mathcal{K}$); note that the assumption of zero spectral measure for 0 is equivalent to the requirement that $\omega_k = 0$ only on a zero-measure subset of $\mathcal{K}$. In these situations we have again Eq. (3.55), describing the cylinder kernel T as the primitive of $-\tilde{T}$.

For example, let us return to the case where $\mathcal{A} = -\Delta$ and $\Omega = \mathbf{R}^d$, in which the spectrum $\sigma(\mathcal{A}) = [0, +\infty)$ is purely continuous. We have mentioned previously the eigenfunctions $F_{\mathbf{k}}$ and the eigenvalues $\omega_{\mathbf{k}}^2$ (labelled by $\mathbf{k} \in \mathbf{R}^d$), where $F_{\mathbf{k}}(\mathbf{x}) = (2\pi)^{-d/2} e^{i\mathbf{k}\cdot\mathbf{x}}$ and $\omega_{\mathbf{k}} = |\mathbf{k}|$; of course $\omega_{\mathbf{k}} = 0$ only on a set of zero Lebesgue measure (consisting of the unique point $\mathbf{k} = 0$). In this case, the expansion (3.53) (involving an integral in the Lebesgue measure $d\mathbf{k}$) gives the following for $d \geq 2$:⁽¹³⁾

$$\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \frac{\Gamma(\frac{d-1}{2})}{2\pi^{\frac{d+1}{2}} (\mathbf{t}^2 + |\mathbf{x} - \mathbf{y}|^2)^{\frac{d-1}{2}}}. \quad (3.61)$$

3.11 Connections between the cylinder kernel and a $(d+1)$ -dimensional Green function. Due to the limited popularity of T it can be useful to connect this kernel to a more familiar object, namely a Green function, even though this requires to pass to $d+1$ dimensions. All details of this construction are given in Appendix C where, as an example, this approach is used for a novel derivation of Eq. (3.60) not relying on the eigenfunction expansion (3.48).

¹³For $d = 1$ the right hand side of Eq.(3.53) for $\tilde{T}$ does not converge (not even distributionally); we take this as an indication that the modified cylinder kernel is ill-defined.

3.12 Behaviour of the heat and cylinder kernels for small and large $\mathfrak{t}$.

The small $\mathfrak{t}$ limit. The asymptotic expansion of the heat kernel of an operator $\mathcal{A} = -\Delta + V$ on an open set $\Omega \subset \mathbf{R}^d$ has been extensively studied (see, e.g., the work of Minakshisundaram and Pleijel [74], or the already cited monographies [11, 18, 22, 28, 53, 54] on the heat kernel). From here to the end of this paragraph, we discuss the $\mathfrak{t} \rightarrow 0^+$ behavior of the heat and cylinder kernels of $\mathcal{A} = -\Delta + V$ under the assumptions (3.9) (these could be generalized to get the same results, but we are not going to discuss this subject). As known from the previously cited references, there is a unique sequence of real functions $a_n : \Omega \times \Omega \rightarrow \mathbf{R}$ ($n = 1, 2, 3, \dots$), usually referred to as HMDS (Hadamard-Minakshisundaram-DeWitt-Seeley) coefficients, such that for any $N \in \{1, 2, 3, \dots\}$ one has

$$K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) = \frac{1}{(4\pi\mathfrak{t})^{d/2}} e^{-\frac{|\mathbf{x}-\mathbf{y}|^2}{4\mathfrak{t}}} \left[1 + \sum_{n=1}^N a_n(\mathbf{x}, \mathbf{y}) \mathfrak{t}^n + O(\mathfrak{t}^{N+1}) \right] \quad \text{for } \mathfrak{t} \rightarrow 0^+. \quad (3.62)$$

In the above equation notice the factor $K_0(\mathfrak{t}; \mathbf{x}, \mathbf{y}) := \frac{1}{(4\pi\mathfrak{t})^{d/2}} e^{-\frac{|\mathbf{x}-\mathbf{y}|^2}{4\mathfrak{t}}}$, which is just the heat kernel associated to $-\Delta$ on $\mathbf{R}^d$. In case $V = 0$, we have $a_n = 0$ for all n ; thus, $K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) = K_0(\mathfrak{t}; \mathbf{x}, \mathbf{y})[1 + O(\mathfrak{t}^\infty)]$ (where the last term indicates a remainder which is $O(\mathfrak{t}^N)$ for each $N \in \{1, 2, 3, \dots\}$).

Along the diagonal $\mathbf{y} = \mathbf{x}$ (for any V) Eq. (3.62) reduces to

$$K(\mathfrak{t}; \mathbf{x}, \mathbf{x}) = \frac{1}{(4\pi\mathfrak{t})^{d/2}} \left[1 + \sum_{n=1}^N a_n(\mathbf{x}) \mathfrak{t}^n + O(\mathfrak{t}^{N+1}) \right] \quad \text{for } \mathfrak{t} \rightarrow 0^+ \quad (3.63)$$

where $a_n(\mathbf{x})$ is shorthand for $a_n(\mathbf{x}, \mathbf{x})$. The small $\mathfrak{t}$ analysis of the cylinder kernel is more involved; however, Fulling proved (see, e.g., [47]) that its asymptotic behaviour along the diagonal $\mathbf{y} = \mathbf{x}$ is as follows: there exist functions $e_n, f_n : \Omega \rightarrow \mathbf{R}$ ($n = 0, 1, 2, \dots$) such that, for any $N \in \{0, 1, 2, \dots\}$,

$$T(\mathfrak{t}; \mathbf{x}, \mathbf{x}) = \frac{1}{\mathfrak{t}^d} \left[\sum_{n=0}^N e_n(\mathbf{x}) \mathfrak{t}^n + \sum_{\substack{n=d+1 \\ n-d \text{ odd}}}^N f_n(\mathbf{x}) \mathfrak{t}^n \ln \mathfrak{t} + O(\mathfrak{t}^{N+1} \ln \mathfrak{t}) \right] \quad \text{for } \mathfrak{t} \rightarrow 0^+. \quad (3.64)$$

As pointed out in [47], some of the functions e_n, f_n (but not all of them) can be expressed in terms of the diagonal HMDS coefficients $\mathbf{x} \mapsto a_n(\mathbf{x})$ mentioned before.

Before proceeding, let us remark that the heat and cylinder traces $K(\mathfrak{t}), T(\mathfrak{t})$ (see Eq.s (3.56) (3.57)) are well-known to admit small $\mathfrak{t}$ expansions analogous to those in Eq.s (3.63) (3.64); see once more the references cited above. In particular, assuming again Ω to be compact and V to be smooth and bounded below, for $\mathfrak{t} \rightarrow 0^+$ there hold

$$K(\mathfrak{t}) = \frac{1}{(4\pi\mathfrak{t})^{d/2}} \left[\text{Vol}(\Omega) + \sum_{n=1}^N A_n \mathfrak{t}^{n/2} + O(\mathfrak{t}^{\frac{N+1}{2}}) \right], \quad (3.65)$$

$$T(\mathfrak{t}) = \frac{1}{\mathfrak{t}^d} \left[\sum_{n=0}^N E_n \mathfrak{t}^n + \sum_{\substack{n=d+1 \\ n-d \text{ odd}}}^N F_n \mathfrak{t}^n \ln \mathfrak{t} + O(\mathfrak{t}^{N+1} \ln \mathfrak{t}) \right] \quad (3.66)$$

($Vol(\Omega)$ denotes the volume of the spatial domain Ω). Notice, in particular, that the expansion (3.65) for $K(\mathfrak{t})$ involves half-integer powers of $\mathfrak{t}$, whereas in expansions (3.62) (3.63) for the local heat kernel $K(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ only integer powers of $\mathfrak{t}$ appear; besides, let us stress that the real coefficients A_n, E_n, F_n in Eq.s (3.65) (3.66) are not just the integrals over the spatial domain Ω of the functions $a_n(\mathbf{x}), e_n(\mathbf{x}), f_n(\mathbf{x})$ of Eq.s (3.63) (3.64), since boundary contributions arise as well.

The large $\mathfrak{t}$ behavior. Let us move on and note that, as anticipated in subsection 3.9, both the heat and the cylinder kernel (along with their traces) vanish exponentially for large $\mathfrak{t}$, under minimal regularity conditions. As an example, we will prove the exponential decay of the cylinder kernel with the following assumptions:

- a) $\mathcal{A}$ is a strictly positive, selfadjoint operator in $L^2(\Omega)$, possessing a generalized complete orthonormal set $(F_k)_{k \in \mathcal{K}}$ of eigenfunctions with eigenvalues ω_k^2 ($\omega_k > \varepsilon > 0$); the F_k 's are continuous functions on Ω .
- b) for all points $\mathbf{x}, \mathbf{y} \in \Omega$ and $\mathfrak{t} > 0$, one has

$$\hat{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y}) := \int_{\mathcal{K}} dk e^{-\omega_k \mathfrak{t}} |F_k(\mathbf{x})| |F_k(\mathbf{y})| < +\infty. \quad (3.67)$$

In this case, starting from the eigenfunction expansion (3.18) we find

$$|T(\mathfrak{t}; \mathbf{x}, \mathbf{y})| \leq \hat{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y}) \quad (3.68)$$

at all points $\mathbf{x}, \mathbf{y} \in \Omega$. On the other hand, after fixing $\tau > 0$ we see that

$$\hat{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y}) \leq e^{-\varepsilon(\mathfrak{t}-\tau)} \hat{T}(\tau; \mathbf{x}, \mathbf{y}) \quad \text{for all } \mathfrak{t} \geq \tau \quad (3.69)$$

(this follows from the definition (3.67) of $\hat{T}$, noting that the inequalities $\omega_k \geq \varepsilon > 0$ and $\mathfrak{t} - \tau \geq 0$ imply $e^{-\omega_k \mathfrak{t}} = e^{-\omega_k(\mathfrak{t}-\tau)} e^{-\omega_k \tau} \leq e^{-\varepsilon(\mathfrak{t}-\tau)} e^{-\omega_k \tau}$). The relations (3.68) and (3.69) give the desired result of exponential decay for T , since at all points $\mathbf{x}, \mathbf{y} \in \Omega$ they imply

$$|T(\mathfrak{t}; \mathbf{x}, \mathbf{y})| \leq e^{-\varepsilon(\mathfrak{t}-\tau)} \hat{T}(\tau; \mathbf{x}, \mathbf{y}) \quad \text{for all } \mathfrak{t} \geq \tau. \quad (3.70)$$

In the above argument, condition (3.67) plays a crucial role. This condition is fulfilled, for example, if $\mathcal{A} = -\Delta + V$ and the assumptions (3.9) are satisfied. In this case we have a uniform bound $\hat{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y}) \leq \check{T}(\mathfrak{t}) < +\infty$ for all $\mathbf{x}, \mathbf{y} \in \overline{\Omega}$ and $\mathfrak{t} > 0$ (see Appendix B, Eq. (B.37)), so that Eq. (3.70) implies

$$|T(\mathfrak{t}; \mathbf{x}, \mathbf{y})| \leq e^{-\varepsilon(\mathfrak{t}-\tau)} \check{T}(\tau) \quad \text{for all } \mathbf{x}, \mathbf{y} \in \overline{\Omega} \text{ and } \mathfrak{t} \geq \tau. \quad (3.71)$$

The exponential decay of the heat kernel and of the traces of both the heat and cylinder kernels can be derived by similar considerations. For alternative approaches, see [28] or [54].

3.13 The Dirichlet kernel as the Mellin transform of the heat or cylinder kernel. The results reported in this subsection are well-known; they were derived by Dowker and Critchley [30], Hawking [58] and Wald [101] and later reconsidered by Moretti et al. (see [17, 77] and citations therein).

The representations of D_s mentioned in the title are useful in view of the analytic continuation with respect to s ; they can be derived starting from the well-known relation (see [80], p.139, Eq.5.9.1)

$$\frac{1}{z^s} = \frac{1}{\Gamma(s)} \int_0^{+\infty} dt \, t^{s-1} e^{-zt} \quad \text{for all } z \in (0, +\infty), s \in \mathbf{C} \text{ with } \Re s > 0. \quad (3.72)$$

This identity, along with the eigenfunction expansions (3.18), (3.47) and (3.48) of the Dirichlet, heat and cylinder kernels, yields

$$D_s(\mathbf{x}, \mathbf{y}) = \frac{1}{\Gamma(s)} \int_0^{+\infty} dt \, t^{s-1} K(t; \mathbf{x}, \mathbf{y}), \quad (3.73)$$

$$D_s(\mathbf{x}, \mathbf{y}) = \frac{1}{\Gamma(2s)} \int_0^{+\infty} dt \, t^{2s-1} T(t; \mathbf{x}, \mathbf{y}). \quad (3.74)$$

For example, Eq. (3.73) is derived via the following chain of equalities:

$$\begin{aligned} D_s(\mathbf{x}, \mathbf{y}) &= \int_{\mathcal{K}} \frac{dk}{\omega_k^{2s}} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) = \int_{\mathcal{K}} dk \frac{1}{\Gamma(s)} \int_0^{+\infty} dt \, t^{s-1} e^{-\omega_k^2 t} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) = \\ &= \frac{1}{\Gamma(s)} \int_0^{+\infty} dt \, t^{s-1} \int_{\mathcal{K}} dk e^{-\omega_k^2 t} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) = \frac{1}{\Gamma(s)} \int_0^{+\infty} dt \, t^{s-1} K(t; \mathbf{x}, \mathbf{y}). \end{aligned} \quad (3.75)$$

In the second passage above, we used Eq. (3.72) with $z = \omega_k^2$; in the third passage, the exchange in the order of integration is justified with arguments similar to those in [73, 74]. The derivation of Eq. (3.74) is similar; in this case one has to resort to Eq. (3.72) with $z = \omega_k$ and s replaced by $2s$.

Eq.s (3.73) (3.74) state that the Dirichlet kernel D_s can be represented as the *Mellin transform* of either the heat or the cylinder kernel; there are analogous relations for the derivatives of the Dirichlet kernel, involving the corresponding derivatives of the heat and cylinder kernels.

In the sequel we are especially interested in the case in which the integral representations (3.73) and (3.74) hold *pointwisely*, at a given pair of points $\mathbf{x}, \mathbf{y} \in \Omega$ (including the case $\mathbf{y} = \mathbf{x}$). This occurs under the following three conditions:

- a) $D_s(\mathbf{x}, \mathbf{y})$ is well defined at the given points $\mathbf{x}, \mathbf{y}$;
- b) $K(t; \mathbf{x}, \mathbf{y})$ and $T(t; \mathbf{x}, \mathbf{y})$ are also well-defined at these points, for all $t > 0$.
- c) the integrals in Eq.s (3.73) (3.74) converge.

With the assumptions (3.9), a) b) c) hold at all points $\mathbf{x}, \mathbf{y} \in \Omega$ if $\Re s > d/2$. In fact: for $\Re s > d/2$, D_s is continuous on $\overline{\Omega} \times \overline{\Omega}$ (see Eq. (3.20)), so there is no problem

with its pointwise evaluation; for each $\mathfrak{t} > 0$, the functions $K(\mathfrak{t}; \cdot, \cdot)$ and $T(\mathfrak{t}; \cdot, \cdot)$ are continuous (in fact C^∞) on $\overline{\Omega} \times \overline{\Omega}$, and again their pointwise evaluation is not a problem; due to the exponential decay of $K(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ and $T(\mathfrak{t}; \mathbf{x}, \mathbf{y})$, the integrals (3.73) (3.74) have no convergence problems for large $\mathfrak{t}$; for any $\mathbf{x}, \mathbf{y} \in \Omega$, due to the $\mathfrak{t} \rightarrow 0^+$ asymptotic expansions (3.62) (3.64), the same integrals are both convergent for $\mathfrak{t}$ close to zero if $\Re s > d/2$ ⁽¹⁴⁾.

The pointwise representations (3.73) (3.74) can be used as a starting point to build the analytic continuation of the function $s \mapsto D_s(\mathbf{x}, \mathbf{y})$ to a larger domain than the one where they are initially granted to hold (e.g., larger than the half plane $\{s \in \mathbb{C} \mid \Re s > d/2\}$ of case (3.9)); we return to this point in the next two subsections. One could make similar considerations for the derivatives of D_s . For example, Eq. (3.73) gives

$$\partial_{\mathbf{x}}^\alpha \partial_{\mathbf{y}}^\beta D_s(\mathbf{x}, \mathbf{y}) = \frac{1}{\Gamma(s)} \int_0^{+\infty} dt \, \mathfrak{t}^{s-1} \partial_{\mathbf{x}}^\alpha \partial_{\mathbf{y}}^\beta K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; \quad (3.76)$$

for any pair of multi-indices α, β . If we make the assumptions (3.9) on $\mathcal{A} = -\Delta + V$ and admit that the $\mathfrak{t} \rightarrow 0^+$ asymptotic expansion (3.62) can be derived term by term, Eq. (3.76) holds pointwisely at all $\mathbf{x}, \mathbf{y} \in \Omega$ if $\Re s > d/2 + j/2$, where $j := |\alpha| + |\beta|$ ⁽¹⁵⁾.

To conclude this subsection let us notice that, setting $\mathbf{y} = \mathbf{x}$ in Eq.s (3.73) (3.74) and integrating over the spatial domain Ω ⁽¹⁶⁾, the relations (3.22) for the trace $\text{Tr } \mathcal{A}^{-s}$ and (3.56) (3.57) for the heat and cylinder traces $K(\mathfrak{t}), T(\mathfrak{t})$ allow us to infer

$$\text{Tr } \mathcal{A}^{-s} = \frac{1}{\Gamma(s)} \int_0^{+\infty} dt \, \mathfrak{t}^{s-1} K(\mathfrak{t}) ; \quad (3.77)$$

$$\text{Tr } \mathcal{A}^{-s} = \frac{1}{\Gamma(2s)} \int_0^{+\infty} dt \, \mathfrak{t}^{2s-1} T(\mathfrak{t}) . \quad (3.78)$$

Eq.s (3.77) (3.78) can be used to continue analytically the function $s \mapsto \text{Tr } (\mathcal{A}^{-s})$; the situation is similar to the one outlined previously for the local counterparts of these equations, and will be reconsidered in the next two subsections.

¹⁴For example, from (3.62) it is clear that, in the limit $\mathfrak{t} \rightarrow 0^+$, one has $K(\mathfrak{t}, \mathbf{x}, \mathbf{y}) = O(1)$ for $\mathbf{y} \neq \mathbf{x}$ and $K(\mathfrak{t}, \mathbf{x}, \mathbf{x}) = O(1/\mathfrak{t}^{d/2})$; so the integral in (3.73) converges for all $\mathbf{x}, \mathbf{y}$ if $\Re s > d/2$.

¹⁵In fact: for $\Re s > d/2 + j/2$, D_s is C^j on $\overline{\Omega} \times \overline{\Omega}$ (see Eq. (3.20)), so there is no problem with pointwise evaluation of its derivatives up to order j ; for each $\mathfrak{t} > 0$, the function $K(\mathfrak{t}; \cdot, \cdot)$ is C^∞ on $\overline{\Omega} \times \overline{\Omega}$, and again the pointwise evaluation of its derivatives is not a problem; the derivatives of $\partial_{\mathbf{x}}^\alpha \partial_{\mathbf{y}}^\beta K(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ are easily seen to decay exponentially, so the integral (3.76) has no convergence problem for large $\mathfrak{t}$; deriving term by term the asymptotic expansion (3.62) we see that, for $\mathfrak{t} \rightarrow 0^+$, $\partial_{\mathbf{x}}^\alpha \partial_{\mathbf{y}}^\beta K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) = O(1)$ for $\mathbf{y} \neq \mathbf{x}$ and $\partial_{\mathbf{x}}^\alpha \partial_{\mathbf{y}}^\beta K(\mathfrak{t}; \mathbf{x}, \mathbf{x}) = O(1/\mathfrak{t}^{d/2 + [j/2]})$ which implies that, for $\Re s > d/2 + j/2$, the integral (3.76) is in any case convergent.

¹⁶Assuming that the order of integration can be interchanged for s is a suitable complex domain.

3.14 Analytic continuation of Mellin transforms via integration by parts.

In the first part of this subsection, the analytic continuation via integration by parts will be presented for an arbitrary Mellin transform; in the second part, we will connect this general construction to the representation of the Dirichlet kernel (resp., of $\text{Tr } \mathcal{A}^{-s}$) as the Mellin transform of either the heat or the cylinder kernel (resp., of their traces).

Let $\mathcal{F} : (0, +\infty) \rightarrow \mathbf{C}$ be a function of the form

$$\mathcal{F}(\mathfrak{t}) = \frac{1}{\mathfrak{t}^\rho} \mathcal{H}(\mathfrak{t}) \quad (3.79)$$

for some $\rho \in \mathbf{C}$ and some smooth function $\mathcal{H} : [0, +\infty) \rightarrow \mathbf{C}$, vanishing exponentially for $\mathfrak{t} \rightarrow +\infty$; consider the Mellin transform of $\mathcal{F}$, i.e., the function

$$\mathfrak{M}(\sigma) := \int_0^{+\infty} d\mathfrak{t} \mathfrak{t}^{\sigma-1} \mathcal{F}(\mathfrak{t}) , \quad (3.80)$$

defined for appropriate $\sigma \in \mathbf{C}$. Due to the hypotheses on $\mathcal{F}$, the integral in Eq. (3.80) converges only for $\sigma \in \mathbf{C}$ with $\Re \sigma > \Re \rho$ and gives an analytic function of σ in this region. However, integrating by parts n times (for any $n \in \{1, 2, 3, \dots\}$) and noting that the boundary terms vanish (for $\Re \sigma > \Re \rho$), we obtain

$$\mathfrak{M}(\sigma) = \frac{(-1)^n}{(\sigma - \rho) \dots (\sigma - \rho + n - 1)} \int_0^{+\infty} d\mathfrak{t} \mathfrak{t}^{\sigma - \rho + n - 1} \frac{d^n \mathcal{H}}{d\mathfrak{t}^n}(\mathfrak{t}) . \quad (3.81)$$

In consequence of the features of the function $\mathcal{H}$, the above integral converges for $\Re \sigma > \Re \rho - n$; thus, Eq. (3.81) yields the analytic continuation of the Mellin transform $\mathfrak{M}(\sigma)$ to the region

$$\{\sigma \in \mathbf{C} \mid \Re \sigma > \Re \rho - n\} \quad (3.82)$$

from which the zeros of the denominator in (3.81) must be removed; this gives a meromorphic function with (possibly) simple poles at the above zeros, which are the points

$$\sigma \in \{\rho, \rho - 1, \dots, \rho - n + 1\} . \quad (3.83)$$

Moreover, since the above results hold for any given $n \in \{1, 2, 3, \dots\}$, they actually allow to determine the analytic continuation of $\mathfrak{M}(\sigma)$ to the whole complex plane with simple poles at the points $\sigma \in \{\rho, \rho - 1, \rho - 2, \dots\}$.

As mentioned before, the above results can be employed to obtain the sought-for analytic continuation of the Dirichlet kernel D_s (treating $\mathbf{x}, \mathbf{y} \in \Omega$ as fixed parameters) starting from its representations (3.73) (3.74) in terms of the heat and cylinder kernel, respectively.

More precisely, consider the case in which the heat or the cylinder kernel is given by a smooth function of $\mathfrak{t}$ rapidly vanishing at infinity, divided by a power of $\mathfrak{t}$; by this we mean that

$$K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) = \frac{1}{\mathfrak{t}^p} H(\mathfrak{t}; \mathbf{x}, \mathbf{y}) \quad \text{or} \quad T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) = \frac{1}{\mathfrak{t}^q} J(\mathfrak{t}; \mathbf{x}, \mathbf{y}) , \quad (3.84)$$

where $p, q \in \mathbf{R}$, $H, J : [0, +\infty) \times \Omega \times \Omega \rightarrow \mathbf{R}$, and it is assumed that (for fixed $\mathbf{x}, \mathbf{y} \in \Omega$) the function $\mathfrak{t} \in [0, +\infty) \mapsto H(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ or $J(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ is smooth and rapidly vanishing for $\mathfrak{t} \rightarrow +\infty$. In these cases the integrals in the right-hand sides of Eq.s (3.73) and (3.74) converge for $\Re s > p$ and $\Re s > q/2$, respectively. Let us mention that, with the assumptions (3.9), the heat kernel of $\mathcal{A} = -\Delta + V$ is as in Eq. (3.84) with $p = d/2$ (recall the asymptotics (3.62)); again with the assumptions (3.9), the cylinder kernel of $\mathcal{A}$ is as in (3.84) with $q = d$, provided that no logarithmic terms appear in the asymptotic expansion (3.64).

Under the previous assumptions, Eq. (3.81), along with Eq.s (3.73) (3.74), gives the following for any $n \in \{1, 2, 3, \dots\}$ ⁽¹⁷⁾:

$$D_s(\mathbf{x}, \mathbf{y}) = \frac{(-1)^n}{\Gamma(s)(s-p)\dots(s-p+n-1)} \int_0^{+\infty} d\mathfrak{t} \mathfrak{t}^{s-p+n-1} \partial_{\mathfrak{t}}^n H(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; \quad (3.85)$$

$$D_s(\mathbf{x}, \mathbf{y}) = \frac{(-1)^n}{\Gamma(2s)(2s-q)\dots(2s-q+n-1)} \int_0^{+\infty} d\mathfrak{t} \mathfrak{t}^{2s-q+n-1} \partial_{\mathfrak{t}}^n J(\mathfrak{t}; \mathbf{x}, \mathbf{y}) . \quad (3.86)$$

Comments analogous to the ones below Eq. (3.81) can be done for the above representations. More in detail, on the one hand Eq. (3.85) gives the analytic continuation of the Dirichlet kernel D_s in the region $\{s \in \mathbf{C} \mid \Re s > p - n\}$ to a meromorphic function with simple poles at $s \in \{p, p-1, \dots, p-n+1\}$; on the other hand, Eq. (3.86) gives the analytic continuation of D_s to the region $\{s \in \mathbf{C} \mid \Re s > (q-n)/2\}$, with (possibly) simple poles at $s \in \{q/2, (q-1)/2, \dots, (q-n+1)/2\}$.

Of course, relations analogous to (3.85) and (3.86) hold as well for the spatial derivatives of the Dirichlet kernel.

In conclusion, let us stress that similar results can be deduced for the trace $\text{Tr } \mathcal{A}^{-s}$ (see Eq. (3.22)), giving its analytic continuation to wider regions in the complex plane. For example, assume the heat trace has the form (compare with the first relation in Eq. (3.84))

$$K(\mathfrak{t}) = \frac{1}{\mathfrak{t}^p} H(\mathfrak{t}) , \quad (3.87)$$

for some $p \in \mathbf{R}$ and some smooth function $H : [0, +\infty) \rightarrow \mathbf{R}$, rapidly vanishing for $\mathfrak{t} \rightarrow +\infty$; then, starting with Eq. (3.77) and using the relations (3.79-3.81), we

¹⁷To obtain Eq. (3.85) one uses Eq.s (3.79-3.81) with $\mathcal{F}(\mathfrak{t}) = K(\mathfrak{t}; \mathbf{x}, \mathbf{y})$, $\rho = p$, $\mathcal{H}(\mathfrak{t}) = H(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ and $\sigma = s$. To obtain Eq. (3.86) one uses Eq.s (3.79-3.81) with $\mathcal{F}(\mathfrak{t}) = T(\mathfrak{t}; \mathbf{x}, \mathbf{y})$, $\rho = q$, $\mathcal{H}(\mathfrak{t}) = J(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ and $\sigma = 2s$.

obtain the following, for $n \in \{1, 2, 3, \dots\}$:

$$\mathrm{Tr} \mathcal{A}^{-s} = \frac{(-1)^n}{\Gamma(s)(s-p)\dots(s-p+n-1)} \int_0^{+\infty} dt \, t^{s-p+n-1} \frac{d^n H}{dt^n}(t) . \quad (3.88)$$

The above relation gives the analytic continuation of $\mathrm{Tr} \mathcal{A}^{-s}$ to the region $\{s \in \mathbf{C} \mid \Re s > p - n\}$ to a meromorphic function with simple poles at $s \in \{p, p-1, \dots, p-n+1\}$. A similar result can be derived using the cylinder trace $T(t)$.

3.15 Analytic continuation of Mellin transforms via complex integration.

Another way to obtain the analytic continuation of the Mellin transform of a given function is available (assuming the latter to fulfill suitable conditions). Consider again the framework of the previous subsection; this time the idea is to re-express the integral in Eq. (3.80) as an integral along a suitable path in the complex plane. To this purpose, first consider the following identity concerning Mellin transforms. Let $t \mapsto h(t)$ be a complex-valued function, analytic in a complex neighborhood of $[0, +\infty)$ and exponentially vanishing for $\Re t \rightarrow +\infty$ in this neighborhood; then

$$\int_0^{+\infty} dt \, t^{s-1} h(t) = \frac{e^{-i\pi s}}{2i \sin(\pi s)} \int_{\mathfrak{H}} dt \, t^{s-1} h(t) \quad \text{for } s \in \mathbf{C} \setminus \{1, 2, 3, \dots\}, \Re s > 0 , \quad (3.89)$$

where $\mathfrak{H}$ denotes the *Hankel contour*, that is a simple path in the complex plane that starts in the upper half-plane near $+\infty$, encircles the origin counterclockwise and returns to $+\infty$ in the lower half-plane (see Fig. 1 below).




Figure 1: The Hankel contour $\mathfrak{H}$.

In the right-hand side of Eq. (3.89), the complex power t^{s-1} is defined making reference to Eq.s (3.15) (3.16); in the left-hand side, since $t \in (0, +\infty)$, we define t^{s-1} according to the standard convention (3.14). See Appendix D for the derivation of Eq. (3.89).

Assume now $\mathcal{F}$ is as in Eq. (3.79) with $\mathcal{H}$ a complex function, analytic in a neighborhood of $[0, +\infty)$ and exponentially vanishing for $\Re t \rightarrow +\infty$; then, considering

the Mellin transform $\mathfrak{M}(\sigma)$ in Eq. (3.80) and using Eq. (3.89) with $s = \sigma - \rho$ and $h = \mathcal{H}$, we obtain

$$\mathfrak{M}(\sigma) = \frac{e^{-i\pi(\sigma-\rho)}}{2i \sin(\pi(\sigma-\rho))} \int_{\mathfrak{H}} d\mathfrak{t} \, \mathfrak{t}^{\sigma-1} \mathcal{F}(\mathfrak{t}) . \quad (3.90)$$

In principle, Eq. (3.90) holds under certain conditions: to ensure existence of $\mathfrak{M}(\sigma)$ as defined by Eq. (3.80) we must require $\Re\sigma > \Re\rho$, and the denominator $\sin(\pi(\sigma-\rho))$ must be nonzero. However, the integral in Eq. (3.90) converges for any $\sigma \in \mathbf{C}$, so this equation yields the analytic continuation of the Mellin transform $\mathcal{M}(\sigma)$ to the whole complex plane, possibly with simple poles for

$$\sigma \in \{\rho, \rho - 1, \rho - 2, \dots\} , \quad (3.91)$$

due to the vanishing of the sine function in the denominator ⁽¹⁸⁾.

Recall once more that, according to Eq.s (3.73) (3.74), the Dirichlet kernel can be expressed as the Mellin transform of either the heat or the cylinder kernel; so, the above results on the analytic continuation via contour integration can be applied to $D_s(\mathbf{x}, \mathbf{y})$ (for fixed $\mathbf{x}, \mathbf{y} \in \Omega$). More precisely, suppose that either the heat or the cylinder kernel has the form (3.84):

$$K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) = \frac{1}{\mathfrak{t}^p} H(\mathfrak{t}; \mathbf{x}, \mathbf{y}) \quad \text{or} \quad T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) = \frac{1}{\mathfrak{t}^q} J(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ,$$

with $p, q \in \mathbf{R}$ and suitable functions $H, J : [0, +\infty) \times \Omega \times \Omega \rightarrow \mathbf{R}$. Assume these functions to have extensions $H, J : \mathcal{U}([0, +\infty)) \times \Omega \times \Omega \rightarrow \mathbf{C}$, where $\mathcal{U}([0, +\infty)) \subset \mathbf{C}$ is an open neighbourhood of the interval $[0, +\infty)$ and, for fixed $\mathbf{x}, \mathbf{y} \in \Omega$, the function $\mathfrak{t} \in \mathcal{U}([0, +\infty)) \mapsto H(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ or $J(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ is analytic and exponentially vanishing for $\Re\mathfrak{t} \rightarrow +\infty$. Making these hypotheses and using Eq. (3.90) along with Eq.s (3.73) (3.74), we obtain ⁽¹⁹⁾

$$D_s(\mathbf{x}, \mathbf{y}) = \frac{e^{-i\pi(s-p)}}{2i \Gamma(s) \sin(\pi(s-p))} \int_{\mathfrak{H}} d\mathfrak{t} \, \mathfrak{t}^{s-1} K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; \quad (3.92)$$

$$D_s(\mathbf{x}, \mathbf{y}) = \frac{e^{-i\pi(2s-q)}}{2i \Gamma(2s) \sin(\pi(2s-q))} \int_{\mathfrak{H}} d\mathfrak{t} \, \mathfrak{t}^{2s-1} T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) . \quad (3.93)$$

¹⁸By inspection of the denominator, it would seem that also the points $\sigma \in \{\rho+1, \rho+2, \rho+3, \dots\}$ are singular, but we know this is not the case since the original expression (3.80) for $\mathfrak{M}(\sigma)$ is regular at these points. The reason for the apparent contradiction lies in the fact that the integral over the Hankel contour in Eq. (3.90) vanishes for the above mentioned values of σ (as can be easily checked via the residue theorem recalling the properties of $\mathcal{F}$), thus yielding an indeterminate form $\infty \cdot 0$.

¹⁹One proceeds as in Footnote 17, using Eq. (3.90) in place of Eq. (3.81).

Due to the comments after Eq. (3.90), both the above identities yield the analytic continuation of the Dirichlet kernel to a meromorphic function on the whole complex plane; more precisely, the analytic continuations obtained via Eq.s (3.92) and (3.93) may have simple poles respectively for $s \in \{p, p-1, p-2, \dots\} \setminus \{0, -1, -2, \dots\}$ and $s \in \{q/2, (q-1)/2, (q-2)/2, \dots\} \setminus \{0, -1/2, -1, -3/2, \dots\}$ (as readily understood analysing the denominators in the right-hand sides of the cited equations).

In the subcases where $p, q \in \mathbf{Z} = \{0, \pm 1, \pm 2, \dots\}$, using trivial trigonometric identities and recalling that $\Gamma(s)\Gamma(1-s)\sin(\pi s) = \pi$ for any $s \in \mathbf{C}$ (see [80], p.138, Eq.5.5.3), Eq.s (3.92) (3.93) can be rephrased as

$$D_s(\mathbf{x}, \mathbf{y}) = \frac{e^{-i\pi s} \Gamma(1-s)}{2\pi i} \int_{\mathfrak{H}} d\mathfrak{t} \mathfrak{t}^{s-1} K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; \quad (3.94)$$

$$D_s(\mathbf{x}, \mathbf{y}) = \frac{e^{-2i\pi s} \Gamma(1-2s)}{2\pi i} \int_{\mathfrak{H}} d\mathfrak{t} \mathfrak{t}^{2s-1} T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) . \quad (3.95)$$

In these subcases the integrals along the Hankel contour can be computed straightforwardly for integer and half-integer values of s , respectively, by means of the residue theorem; for example, for $s = -n/2$ and $n \in \{0, 1, 2, \dots\}$, Eq. (3.95) yields

$$D_{-\frac{n}{2}}(\mathbf{x}, \mathbf{y}) = (-1)^n \Gamma(n+1) \operatorname{Res}\left(\mathfrak{t}^{-(n+1)} T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; 0\right) . \quad (3.96)$$

We can obtain a variant of Eq. (3.95), giving the analytic continuation of the Dirichlet kernel D_s in terms of the modified cylinder kernel $\tilde{T}$ (recall Eq.s (3.53) (3.55)). To this purpose, we assume $\tilde{T}$ to admit a meromorphic extension in $\mathfrak{t}$ to a neighborhood of $[0, +\infty)$, having a pole in $\mathfrak{t} = 0$ and rapidly vanishing for $\Re \mathfrak{t} \rightarrow +\infty$; then, expressing the cylinder kernel $T(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ in Eq. (3.95) as $-\partial_{\mathfrak{t}} \tilde{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ and integrating by parts, we obtain

$$D_s(\mathbf{x}, \mathbf{y}) = - \frac{e^{-2i\pi s} \Gamma(2-2s)}{2\pi i} \int_{\mathfrak{H}} d\mathfrak{t} \mathfrak{t}^{2s-2} \tilde{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y}) \quad (3.97)$$

(note that no boundary contribution arises, due to the rapid vanishing of $\tilde{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ for $\Re \mathfrak{t} \rightarrow +\infty$). Again, for half-integer values of s we can compute explicitly the analytic continuation (3.97) by means of the residue theorem; to be more precise, for $s = -n/2$ and $n \in \{-1, 0, 1, 2, \dots\}$ we have

$$D_{-\frac{n}{2}}(\mathbf{x}, \mathbf{y}) = (-1)^{n+1} \Gamma(n+2) \operatorname{Res}\left(\mathfrak{t}^{-(n+2)} \tilde{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; 0\right) . \quad (3.98)$$

Relations similar to the ones obtained above hold for the spatial derivatives of the Dirichlet kernel, allowing in turn to determine their analytic continuations.

To conclude, let us mention that similar results hold as well for the trace $\operatorname{Tr} \mathcal{A}^{-s}$; these are obtained using the representations (3.77) (3.78) in terms of the heat and

cylinder trace $K(\mathfrak{t}), T(\mathfrak{t})$, and assuming suitable features for the latters. In particular, if the map $\mathfrak{t} \mapsto T(\mathfrak{t})$ admits a meromorphic extension to a neighborhood of $[0, +\infty)$ which only has a pole singularity at $\mathfrak{t} = 0$ and vanishes exponentially for $\Re \mathfrak{t} \rightarrow +\infty$, for $n \in \{0, 1, 2, \dots\}$, we have

$$\mathrm{Tr} \mathcal{A}^{n/2} = (-1)^n \Gamma(n+1) \mathrm{Res} \left(\mathfrak{t}^{-(n+1)} T(\mathfrak{t}); 0 \right). \quad (3.99)$$

3.16 Other kernels, and their relations with D_s . In this section we are considering a number of integral kernels connected with the zeta regularization of the stress-energy VEV; attention is mainly focused on the kernels used in the subsequent applications (including the subsequent Parts II, III and IV). However, it would be against the spirit of this section to ignore completely the *resolvent kernel*, i.e., the kernel of the operator $(\mathcal{A} - \lambda)^{-1}$, where $\mathcal{A}$ is a selfadjoint operator and $\lambda \in \mathbf{C}$ is outside the spectrum $\sigma(\mathcal{A})$. Under appropriate conditions (in particular, the strict positivity of $\mathcal{A}$), the powers $\mathcal{A}^{-s}$ can be related to suitable contour integrals involving the resolvent [95]; this fact can be restated in terms of a relation between the corresponding kernels. This possibility will not be considered here and in Parts II-IV, but we plan to recover it elsewhere.

3.17 The case of product domains. Factorization of the heat kernel. Let us consider the case where $\mathcal{A} = -\Delta + V$ and the spatial domain $\Omega \subset \mathbf{R}^d$ has the form

$$\Omega = \Omega_1 \times \Omega_2 \quad (3.100)$$

with Ω_a , for $a \in \{1, 2\}$, indicating an open subset of $\mathbf{R}^{d_a}$ ($d_1 + d_2 = d$); in this case, points of Ω will be written as

$$\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2), \quad \mathbf{y} = (\mathbf{y}_1, \mathbf{y}_2) \quad (3.101)$$

etc., where $\mathbf{x}_a, \mathbf{y}_a \in \Omega_a$ ($a \in \{1, 2\}$). In addition to Eq. (3.100), we assume that the external potential has the form

$$V(\mathbf{x}) = V_1(\mathbf{x}_1) + V_2(\mathbf{x}_2) \quad (3.102)$$

and that the boundary conditions specified on $\partial\Omega = (\partial\Omega_1 \times \Omega_2) \cup (\Omega_1 \times \partial\Omega_2)$ arise from suitable boundary conditions prescribed separately on $\partial\Omega_1$ and $\partial\Omega_2$, in such a way that, for $a \in \{1, 2\}$, the operator

$$\mathcal{A}_a := -\Delta_a + V(\mathbf{x}_a) \quad (3.103)$$

(Δ_a the Laplacian on Ω_a) is selfadjoint in $L^2(\Omega_a)$. Moreover, each one of these operators is assumed to be strictly positive or, at least, non-negative.

In the situation described above, the Hilbert space $L^2(\Omega)$ and the fundamental operator $\mathcal{A} := -\Delta + V(\mathbf{x})$ acting therein can be represented, respectively, as

$$L^2(\Omega) = L^2(\Omega_1) \otimes L^2(\Omega_2) , \quad \mathcal{A} = \mathcal{A}_1 \otimes \mathbf{1} + \mathbf{1} \otimes \mathcal{A}_2 . \quad (3.104)$$

Because of the assumptions we have made, each of the two operators $\mathcal{A}_a$ ($a \in \{1, 2\}$) possesses a complete orthonormal system of eigenfunctions $(F_{a,k_a})_{k_a \in \mathcal{K}_a}$ with eigenvalues ω_{a,k_a}^2 ; as for the fundamental operator $\mathcal{A}$, we see that it has a complete orthonormal set of eigenfunctions of the form

$$F_k(\mathbf{x}) := F_{1,k_1}(\mathbf{x}_1) F_{2,k_2}(\mathbf{x}_2) \quad \text{for } k = (k_1, k_2) \in \mathcal{K}_1 \times \mathcal{K}_2 \quad (3.105)$$

and that

$$\mathcal{A} F_k = \omega_k^2 F_k , \quad \omega_k^2 = \omega_{1,k_1}^2 + \omega_{2,k_2}^2 . \quad (3.106)$$

Of course $\sigma(\mathcal{A}) = \sigma(\mathcal{A}_1) + \sigma(\mathcal{A}_2)$, so that $\mathcal{A}$ is non-negative; besides, $\mathcal{A}$ is strictly positive if so is one at least between $\mathcal{A}_1$ and $\mathcal{A}_2$.

In the product case under analysis, a number of interesting facts occurs for the integral kernels associated to $\mathcal{A}$ and $\mathcal{A}_1, \mathcal{A}_2$. The most elementary of these facts is the factorization of the heat kernel; by this we mean that the kernels

$$K(\mathbf{t}; \mathbf{x}, \mathbf{y}) := (e^{-\mathbf{t}\mathcal{A}})(\mathbf{x}, \mathbf{y}) , \quad K_a(\mathbf{t}; \mathbf{x}_a, \mathbf{y}_a) := (e^{-\mathbf{t}\mathcal{A}_a})(\mathbf{x}_a, \mathbf{y}_a) \quad (a \in \{1, 2\}) \quad (3.107)$$

are related by

$$K(\mathbf{t}; \mathbf{x}, \mathbf{y}) = K_1(\mathbf{t}; \mathbf{x}_1, \mathbf{y}_1) K_2(\mathbf{t}; \mathbf{x}_2, \mathbf{y}_2) , \quad (3.108)$$

a fact that is made apparent by the eigenfunction expansion (3.47) and by Eq.s (3.105) (3.106).

In passing, let also mention that an analogous relation can be easily derived for the heat trace; writing $K(\mathbf{t})$, $K_a(\mathbf{t})$ for the heat traces of $\mathcal{A}$ and $\mathcal{A}_a$ ($a \in \{1, 2\}$), respectively, we obtain ⁽²⁰⁾

$$K(\mathbf{t}) = K_1(\mathbf{t}) K_2(\mathbf{t}) . \quad (3.109)$$

In the present subsection we have analysed the case of a product configuration with two factors; as a straightforward generalization, one can consider a product with an arbitrary number of factors. Examples of such multiple products will appear in Parts III and IV.

²⁰In fact, Eq.s (3.100) (3.108) and the relations (3.56) (3.57) allow us to infer the following chain of equalities:

$$K(\mathbf{t}) = \int_{\Omega} d\mathbf{x} K(\mathbf{t}; \mathbf{x}, \mathbf{x}) = \int_{\Omega_1} d\mathbf{x}_1 K_1(\mathbf{t}; \mathbf{x}_1, \mathbf{x}_1) \int_{\Omega_2} d\mathbf{x}_2 K_2(\mathbf{t}; \mathbf{x}_2, \mathbf{x}_2) = K_1(\mathbf{t}) K_2(\mathbf{t}) .$$

3.18 The case of a slab: reduction to a lower-dimensional problem. By definition, we have a slab if

$$\Omega = \Omega_1 \times \mathbf{R}^{d_2}, \quad V = V(\mathbf{x}_1) \quad (3.110)$$

with Ω_1 a domain in $\mathbf{R}^{d_1}$ ($d_1 + d_2 = d$), and if the boundary conditions prescribed for the field refer to its behaviour on $\partial\Omega_1 \times \mathbf{R}^{d_2}$. Clearly, a slab is a subcase of the general product case discussed in the previous subsection, with $\Omega_2 = \mathbf{R}^{d_2}$ and $V_2 = 0$. In this subcase the relevant operators are $\mathcal{A} = -\Delta + V(\mathbf{x}_1)$ acting in $L^2(\Omega)$,

$$\mathcal{A}_1 := -\Delta_1 + V(\mathbf{x}_1) \quad (3.111)$$

acting in $L^2(\Omega_1)$, and $\mathcal{A}_2 := -\Delta_2$ acting in $L^2(\mathbf{R}^{d_2})$.

The operator $\mathcal{A}_1$ has its own eigenfunctions $F_{1,k_1}(\mathbf{x}_1) \equiv \mathfrak{F}_{k_1}(\mathbf{x}_1)$ and eigenvalues $\omega_{1,k_1}^2 \equiv \varpi_{k_1}^2$ ($k_1 \in \mathcal{K}_1$); we assume $\mathcal{A}_1$ to be strictly positive, so that $\varpi_{k_1} \geq \varepsilon$ for some $\varepsilon > 0$. Of course, $-\Delta_2$ is non-negative with eigenfunctions $F_{2,\mathbf{k}_2}(\mathbf{x}_2) = (2\pi)^{-d_2/2} e^{i\mathbf{k}_2 \cdot \mathbf{x}_2}$ and eigenvalues $\omega_{2,\mathbf{k}_2}^2 = |\mathbf{k}_2|^2$, for $\mathbf{k}_2 \in \mathbf{R}^{d_2}$.

In the sequel we write $D_s(\mathbf{x}_1, \mathbf{x}_2; \mathbf{y}_1, \mathbf{y}_2)$ for the Dirichlet kernel of $\mathcal{A}$ at the points $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2)$ and $\mathbf{y} = (\mathbf{y}_1, \mathbf{y}_2)$; $D_s^{(1)}(\mathbf{x}_1, \mathbf{y}_1)$ will be the Dirichlet kernel of $\mathcal{A}_1$.

On the one hand, according to the general results (3.31-3.33), the regularized VEV $\langle 0 | \hat{T}_{\mu\nu}^u(\mathbf{x}) | 0 \rangle$ is determined by Dirichlet kernel $D_s(\mathbf{x}, \mathbf{y})$ and its derivatives evaluated on the diagonal $\mathbf{y} = \mathbf{x}$. The main intent of this subsection is to express D_s and its derivatives in terms of the reduced kernel $D_s^{(1)}$ at all points of the diagonal $\mathbf{y} = \mathbf{x}$ (and, in fact, on an even larger domain). The starting point towards this goal is the identity

$$D_s(\mathbf{x}_1, \mathbf{x}_2; \mathbf{y}_1, \mathbf{y}_2) = \hat{D}_s(\mathbf{x}_1, \mathbf{y}_1; |\mathbf{x}_2 - \mathbf{y}_2|^2), \quad (3.112)$$

involving a function $\hat{D}_s : \Omega_1 \times \Omega_1 \times [0, +\infty) \rightarrow \mathbf{C}$, $(\mathbf{x}_1, \mathbf{y}_1, q) \mapsto \hat{D}_s(\mathbf{x}_1, \mathbf{y}_1, q)$. This function and its partial derivatives with respect to q , at $q = 0$, are completely determined by the kernel $D_s^{(1)}$, according to the following rules:

$$\hat{D}_s(\mathbf{x}_1, \mathbf{y}_1; 0) = \frac{\Gamma(s - \frac{d_1}{2})}{(4\pi)^{d_1/2} \Gamma(s)} D_{s - \frac{d_1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \quad \text{for } s \in \mathbf{C}, \Re s > \frac{d_1}{2}; \quad (3.113)$$

$$\frac{\partial^n \hat{D}_s}{\partial q^n}(\mathbf{x}_1, \mathbf{y}_1; 0) = \frac{(-1)^n \Gamma(s - \frac{d_1}{2} - n)}{(4\pi)^{d_1/2} 4^n \Gamma(s)} D_{s - \frac{d_1}{2} - n}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \quad \text{for } s \in \mathbf{C}, \Re s > \frac{d_1}{2} + n. \quad (3.114)$$

We defer to Appendix E the derivation of Eq.s (3.112-3.114). In particular, for the derivatives involved in Eq.s (3.31-3.33) on $\langle 0 | \hat{T}_{\mu\nu}^u(\mathbf{x}) | 0 \rangle$ we obtain the following expressions (where, for simplicity of notation, we write $(\mathbf{x}, \mathbf{y})$ for $(\mathbf{x}_1, \mathbf{x}_2; \mathbf{y}_1, \mathbf{y}_2)$):

$$D_{\frac{u \pm 1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} = \frac{\Gamma(\frac{u-d_2 \pm 1}{2})}{(4\pi)^{d_2/2} \Gamma(\frac{u \pm 1}{2})} D_{\frac{u-d_2 \pm 1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \Big|_{\mathbf{y}_1=\mathbf{x}_1}; \quad (3.115)$$

$$\partial_{x_a^i y_b^j} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} = \partial_{x_a^i x_b^j} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} = \partial_{y_a^i y_b^j} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} = 0 \quad (3.116)$$

for $(a, b) = (1, 2)$ or $(a, b) = (2, 1)$ and $i \in \{1, \dots, d_a\}$, $j \in \{1, \dots, d_b\}$;

$$\partial_{z_1^i w_1^j} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} = \frac{\Gamma(\frac{u-d_2+1}{2})}{(4\pi)^{d_2/2} \Gamma(\frac{u+1}{2})} \partial_{z_1^i w_1^j} D_{\frac{u-d_2+1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \Big|_{\mathbf{y}_1=\mathbf{x}_1} \quad (3.117)$$

for $z, w \in \{x, y\}$ and $i, j \in \{1, \dots, d_1\}$;

$$\begin{aligned} \partial_{x_2^i y_2^j} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} &= -\partial_{x_2^i x_2^j} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} = -\partial_{y_2^i y_2^j} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} = \\ &= \delta_{ij} \frac{\Gamma(\frac{u-d_2-1}{2})}{(4\pi)^{d_2/2} 2 \Gamma(\frac{u+1}{2})} D_{\frac{u-d_2-1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \Big|_{\mathbf{y}_1=\mathbf{x}_1} \quad \text{for } i, j \in \{1, \dots, d_2\} . \end{aligned} \quad (3.118)$$

Relations (3.115-3.118) are derived assuming $\Re u > d_2 + 1$, but it follows from them that the analytic continuations in u of the Dirichlet kernel, of its reduced analogue and of their derivatives fulfill the very same relations.

Let us remark that the left-hand sides of the above equations depend in principle on $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2)$, while the right-hand sides only contain $\mathbf{x}_1$; this confirms the expectation that the stress-energy VEV ought to be independent of the variable $\mathbf{x}_2$, due to the symmetry of the slab configuration under translations regarding this variable alone. Finally, using Eq.s (3.115-3.118) and (3.31-3.33), it can be easily checked that the components of the regularized stress-energy tensor VEV also fulfill

$$\begin{aligned} \langle 0 | \widehat{T}_{ij}^u(\mathbf{x}) | 0 \rangle &= 0 \quad \text{for } i, j \in \{d_1 + 1, \dots, d\}, i \neq j ; \\ \langle 0 | \widehat{T}_{ij}^u(\mathbf{x}) | 0 \rangle &= \langle 0 | \widehat{T}_{ji}^u(\mathbf{x}) | 0 \rangle = 0 \quad \text{for } i \in \{1, \dots, d_1\}, j \in \{d_1 + 1, \dots, d\} . \end{aligned} \quad (3.119)$$

4 Total energy and forces on the boundary

We refer again to the general framework of Section 2, where a quantized scalar field on a spatial domain Ω and the associated stress-energy tensor VEV are considered; we recall that $\mathcal{A}$ indicates the fundamental operator $-\Delta + V(\mathbf{x})$ acting in $L^2(\Omega)$. From now on, we indicate with da the area element on the boundary $\partial\Omega$, and use for Ω the standard Lebesgue measure $d\mathbf{x}$. We also write $\mathbf{n}(\mathbf{x}) \equiv (n^\ell(\mathbf{x}))_{\ell=1, \dots, d}$ for the outer unit normal at a point $\mathbf{x} \in \partial\Omega$; this is assumed to exist everywhere (which happens if $\partial\Omega$ is globally smooth) or almost everywhere (which happens if $\partial\Omega$ has edges or corners).

4.1 The total energy. As anticipated in subsection 2.6, the zeta-regularized total energy can be defined as

$$\mathcal{E}^u := \int_{\Omega} d\mathbf{x} \langle 0 | \widehat{T}_{00}^u(\mathbf{x}) | 0 \rangle , \quad (4.1)$$

provided that the above integral converges for u in a suitable complex domain; using Eq. (3.31), the above definition yields

$$\begin{aligned} \mathcal{E}^u &= \kappa^u \left(\frac{1}{4} + \xi \right) \int_{\Omega} d\mathbf{x} D_{\frac{u-1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} + \\ &+ \kappa^u \left(\frac{1}{4} - \xi \right) \int_{\Omega} d\mathbf{x} \left[\left(\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x}) \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} . \end{aligned} \quad (4.2)$$

On the other hand, the eigenfunction expansion (3.18) for the Dirichlet kernel (here used with $s = \frac{u+1}{2}$) gives

$$\begin{aligned} &\int_{\Omega} d\mathbf{x} \left[\left(\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x}) \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} = \\ &= \int_{\mathcal{K}} \frac{dk}{\omega_k^{u+1}} \int_{\Omega} d\mathbf{x} \left(\partial^\ell F_k(\mathbf{x}) \partial_\ell \overline{F_k}(\mathbf{x}) + V(\mathbf{x}) F_k(\mathbf{x}) \overline{F_k}(\mathbf{x}) \right) = \\ &= \int_{\mathcal{K}} \frac{dk}{\omega_k^{u+1}} \left(\int_{\Omega} d\mathbf{x} \left(F_k(\mathbf{x}) (-\partial^\ell \partial_\ell + V(\mathbf{x})) \overline{F_k}(\mathbf{x}) \right) + \int_{\partial\Omega} da(\mathbf{x}) F_k(\mathbf{x}) n^\ell(\mathbf{x}) \partial_\ell \overline{F_k}(\mathbf{x}) \right) \end{aligned} \quad (4.3)$$

where, in the last step, we have integrated by parts ⁽²¹⁾.

To go on we note that $(-\partial^\ell \partial_\ell + V) \overline{F_k} = \overline{\mathcal{A} F_k} = \omega_k^2 \overline{F_k}$ and $F_k(\mathbf{x}) n^\ell(\mathbf{x}) \partial_\ell \overline{F_k}(\mathbf{x}) = F_k(\mathbf{x}) \frac{\partial \overline{F_k}}{\partial n}(\mathbf{x}) = F_k(\mathbf{x}) \frac{\partial \overline{F_k}(\mathbf{y})}{\partial n_{\mathbf{y}}} \Big|_{\mathbf{y}=\mathbf{x}}$, where we have introduced the normal derivative $\partial/\partial n := n^\ell \partial_{x^\ell}$. Substituting into Eq. (4.3) and summing over $k \in \mathcal{K}$, we obtain

$$\begin{aligned} &\int_{\Omega} d\mathbf{x} \left[\left(\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x}) \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} = \\ &= \int_{\Omega} d\mathbf{x} D_{\frac{u-1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} + \int_{\partial\Omega} da(\mathbf{x}) \frac{\partial}{\partial n_{\mathbf{y}}} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{x}} ; \end{aligned} \quad (4.4)$$

inserting this result into Eq. (4.2), we conclude

$$\mathcal{E}^u = E^u + B^u , \quad (4.5)$$

where we have introduced the *regularized bulk* and *boundary energies*

$$E^u := \frac{\kappa^u}{2} \int_{\Omega} d\mathbf{x} D_{\frac{u-1}{2}}(\mathbf{x}, \mathbf{x}) = \frac{\kappa^u}{2} \text{Tr } \mathcal{A}^{\frac{1-u}{2}} , \quad (4.6)$$

²¹ Here and in similar situations, whenever we speak of an integration by parts we refer to the identity

$$\int_{\Omega} d\mathbf{x} (\partial_\ell f) g = \int_{\partial\Omega} da f g n_\ell - \int_{\Omega} d\mathbf{x} f \partial_\ell g ,$$

holding for all sufficiently smooth functions f, g .

$$B^u := \kappa^u \left(\frac{1}{4} - \xi \right) \int_{\partial\Omega} da(\mathbf{x}) \left. \frac{\partial}{\partial n_{\mathbf{y}}} D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right|_{\mathbf{y}=\mathbf{x}} . \quad (4.7)$$

The derivation of the above result is a bit formal since, in general, one cannot grant convergence of the integrals defining E^u and B^u , for suitable values $u \in \mathbf{C}$.

As for the bulk energy, it is easy to give an example in which finiteness is granted for appropriate u . To this purpose let us consider the case (3.9) (involving a bounded domain with Dirichlet boundary conditions); in this case, recalling the Weyl estimates (3.10), we conclude

$$E^u \text{ is finite if } \Re u > d+1 . \quad (4.8)$$

As for the regularized boundary energy B^u let us mention that, for Ω bounded,

$$B^u = 0 \quad \text{under Dirichlet or Neumann boundary conditions on } \partial\Omega \quad (4.9)$$

(since in the Dirichlet case we have $D_s(\mathbf{x}, \mathbf{y}) = 0$ for $\mathbf{x} \in \partial\Omega$ and all $\mathbf{y}$, while in the Neumann case $\frac{\partial}{\partial n_{\mathbf{y}}} D_s(\mathbf{x}, \mathbf{y}) = 0$ for $\mathbf{y} \in \partial\Omega$ and all $\mathbf{x}$).

Applying the above considerations in the case of an unbounded domain requires much caution. On the one hand, E^u can be infinite for all $u \in \mathbf{C}$; on the other hand, in the definition (4.7) of B^u it might be necessary to intend the integral $\int_{\partial\Omega} da$ as $\lim_{\ell \rightarrow +\infty} \int_{\partial\Omega_\ell} da$, where $(\Omega_\ell)_{\ell=0,1,2,\dots}$ is a sequence of bounded subdomains such that $\Omega_\ell \subset \Omega_{\ell+1}$ (for any $\ell \in \{0, 1, 2, \dots\}$) and $\cup_{\ell=0}^{+\infty} \Omega_\ell = \Omega$ (note that this limit could either be infinite or even fail to exist).

If E^u exists finite for u belonging to a suitable open subset of $\mathbf{C}$ and it is an analytic function of u on this domain, a renormalization by analytic continuation can be implemented; in general, following the extended version of the zeta approach, we define the renormalized bulk energy as

$$E^{ren} := RP \Big|_{u=0} E^u = RP \Big|_{u=0} \left(\frac{\kappa^u}{2} \text{Tr } \mathcal{A}^{\frac{1-u}{2}} \right) . \quad (4.10)$$

When the analytic continuation of $\text{Tr } \mathcal{A}^{\frac{1-u}{2}}$ is regular up to $u = 0$ the above prescription is reduced to

$$E^{ren} := E^u \Big|_{u=0} = \frac{1}{2} \text{Tr } \mathcal{A}^{1/2} \quad (4.11)$$

(of course $\text{Tr } \mathcal{A}^{1/2}$ indicates the analytic continuation of $\text{Tr } \mathcal{A}^{\frac{1-u}{2}}$ at $u = 0$).

In a similar way one can define the renormalized boundary and total energies as

$$B^{ren} := RP \Big|_{u=0} B^u ; \quad (4.12)$$

$$\mathcal{E}^{ren} := RP \Big|_{u=0} \mathcal{E}^u . \quad (4.13)$$

An alternative definition of the renormalized total energy could be

$$\mathcal{E}^{ren} := \int_{\Omega} d\mathbf{x} \langle 0 | \widehat{T}_{00}(\mathbf{x}) | 0 \rangle_{ren} . \quad (4.14)$$

This possibility, which is considered rarely in this series of papers, is not granted to be equivalent to (4.13); for example, it may happen that the integral in the right-hand side of Eq. (4.14) diverges, while the prescription (4.13) always gives a finite result by construction. For a comparison between the alternatives (4.13)–(4.14), see the final lines of subsection 6.4 (dealing with a field on a segment, for several types of boundary conditions).

4.1.1 Reduced energy for a slab configuration. Let us consider the slab configuration introduced in subsection 3.18, so that $\Omega := \Omega_1 \times \mathbf{R}^{d_2}$, the potential V depends only on $\mathbf{x}_1 \in \Omega_1$, and the boundary conditions regard only $\partial\Omega_1 \times \Omega_2$. We already observed in the mentioned subsection that the regularized VEV $\langle 0 | \widehat{T}_{\mu\nu} | 0 \rangle$ depends only on $\mathbf{x}_1 \in \Omega_1$ (and not on $\mathbf{x}_2 \in \mathbf{R}^{d_2}$); so, the integral in Eq. (4.1) defining the total energy diverges due to an infinite volume factor.

As a matter of fact, when dealing with a slab configuration one usually considers in place of the total energy $\mathcal{E}^u$ the *reduced total energy* $\mathcal{E}_1^s$; this is the total energy per unit volume in the “free” dimensions, i.e.,

$$\mathcal{E}_1^u := \int_{\Omega_1} d\mathbf{x}_1 \langle 0 | \widehat{T}_{00}^u | 0 \rangle . \quad (4.15)$$

Recalling Eq.s (3.115–3.118) and using some well-known identities regarding the gamma function, we infer

$$\begin{aligned} \mathcal{E}_1^u = & \frac{\kappa^u \Gamma(\frac{u-d_2+1}{2})}{(4\pi)^{d_2/2} \Gamma(\frac{u+1}{2})} \left\{ \left(\frac{u-1+d_2}{4(u-1-d_2)} + \xi \right) \int_{\Omega_1} d\mathbf{x}_1 D_{\frac{u-d_2-1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \Big|_{\mathbf{y}_1=\mathbf{x}_1} + \right. \\ & \left. + \left(\frac{1}{4} - \xi \right) \int_{\Omega_1} d\mathbf{x}_1 \left[\left(\partial_{x_1^\ell} \partial_{y_1^\ell} + V(\mathbf{x}_1) \right) D_{\frac{u-d_2+1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \right] \Big|_{\mathbf{y}_1=\mathbf{x}_1} \right\} . \end{aligned} \quad (4.16)$$

Concerning the second term above, we can express the reduced Dirichlet kernel $D_{\frac{u-d_2+1}{2}}^{(1)}$ in terms of the eigenfunctions $(\mathfrak{F}_{k_1}(\mathbf{x}_1))_{k_1 \in \mathcal{K}_1}$ and the eigenvalues $(\varpi_{k_1})_{k_1 \in \mathcal{K}_1}$ of the reduced operator $\mathcal{A}_1 = -\Delta_1 + V(\mathbf{x}_1)$ and integrate by parts as in the general setting; working as in the derivation of Eq.s (4.3)–(4.4) and keeping in mind that $\mathcal{A}_1 \mathfrak{F}_{k_1} = \varpi_{k_1}^2 \mathfrak{F}_{k_1}$, we obtain

$$\begin{aligned} & \int_{\Omega_1} d\mathbf{x}_1 \left[\left(\partial_{x_1^\ell} \partial_{y_1^\ell} + V(\mathbf{x}_1) \right) D_{\frac{u-d_2+1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \right] \Big|_{\mathbf{y}_1=\mathbf{x}_1} = \\ & = \int_{\Omega_1} d\mathbf{x}_1 D_{\frac{u-d_2-1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \Big|_{\mathbf{y}_1=\mathbf{x}_1} + \int_{\partial\Omega_1} da(\mathbf{x}_1) \frac{\partial}{\partial n_{\mathbf{y}_1}} D_{\frac{u-d_2+1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \Big|_{\mathbf{y}_1=\mathbf{x}_1} . \end{aligned} \quad (4.17)$$

In conclusion, we have a result similar to Eq. (4.5):

$$\mathcal{E}_1^u = E_1^u + B_1^u , \quad (4.18)$$

where we have introduced the *regularized reduced bulk* and *boundary energies*

$$\begin{aligned} E_1^u &:= \frac{\kappa^u \Gamma(\frac{u-d_2-1}{2})}{2(4\pi)^{d_2/2} \Gamma(\frac{u-1}{2})} \int_{\Omega_1} d\mathbf{x}_1 D_{\frac{u-d_2-1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{x}_1) = \\ &= \frac{\kappa^u \Gamma(\frac{u-d_2-1}{2})}{2(4\pi)^{d_2/2} \Gamma(\frac{u-1}{2})} \text{Tr} \mathcal{A}_1^{\frac{d_2+1-u}{2}} , \end{aligned} \quad (4.19)$$

$$B_1^u := \frac{\kappa^u \Gamma(\frac{u-d_2+1}{2})}{(4\pi)^{d_2/2} \Gamma(\frac{u+1}{2})} \left(\frac{1}{4} - \xi \right) \int_{\partial\Omega_1} da(\mathbf{x}_1) \frac{\partial}{\partial n_{\mathbf{y}_1}} D_{\frac{u-d_2+1}{2}}^{(1)}(\mathbf{x}_1, \mathbf{y}_1) \Big|_{\mathbf{y}_1=\mathbf{x}_1} . \quad (4.20)$$

The considerations of the previous subsection about convergence of the bulk and boundary energies E^u , B^u have obvious analogues for the reduced energies E_1^u, B_1^u . Of course, the reduced bulk and boundary energies are renormalized in terms of the analytic continuation (or, possibly, of its regular part) at $u = 0$.

4.2 Pressure on the boundary. In this subsection we are interested in the *pressure* $\mathbf{p}(\mathbf{x}) \equiv (p_i(\mathbf{x}))_{i=1,\dots,d}$, i.e., the force per unit area produced by the field inside Ω at a point $\mathbf{x}$ on the boundary $\partial\Omega$. A possible characterization is the following: we first introduce, for $\Re u$ large, the *regularized pressure* $\mathbf{p}^u(\mathbf{x})$ of components

$$p_i^u(\mathbf{x}) := \langle 0 | \widehat{T}_{ij}^u(\mathbf{x}) | 0 \rangle n^j(\mathbf{x}) \quad \text{for } i \in \{1, \dots, d\} ; \quad (4.21)$$

then, we define the *renormalized pressure* at $\mathbf{x}$ setting

$$p_i^{ren}(\mathbf{x}) := RP \Big|_{u=0} p_i^u(\mathbf{x}) \quad (4.22)$$

where $RP|_{u=0}$ indicates the regular of the analytic continuation evaluated at $u = 0$ (of course, if the mentioned continuation is regular up to $u = 0$, the above prescription reduces to $p_i^{ren}(\mathbf{x}) := p_i^u(\mathbf{x})|_{u=0}$, meaning that the analytic continuation at $u = 0$ has to be considered).

It is important to point out that this is not the only reasonable definition for the renormalized pressure at a point $\mathbf{x} \in \Omega$; another possibility is

$$p_i^{ren}(\mathbf{x}) := \left(\lim_{\mathbf{x}' \in \Omega, \mathbf{x}' \rightarrow \mathbf{x}} \langle 0 | \widehat{T}_{ij}(\mathbf{x}') | 0 \rangle_{ren} \right) n^j(\mathbf{x}) . \quad (4.23)$$

In few words: in the approach (4.21-4.22), one *stays at a point on the boundary*, and performs therein the renormalization; in the approach (4.23), one renormalizes *at points inside* Ω , and then moves towards the boundary. Notice that both approaches

require the existence of the normal $\mathbf{n}(\mathbf{x})$ (and thus lose meaning on edges and corner points of $\partial\Omega$).

As a matter of fact, the prescriptions (4.21-4.22) and (4.23) do not always agree. The approach (4.22) (possibly, in the restricted version) gives by construction a finite pressure; on the contrary, this is not granted for the alternative prescription (4.23). As an example, in Part II of this series of papers we discuss the case where the spatial domain Ω is a wedge; in this case at all boundary points not in the edge, where the normal is well defined, the pressure defined according to Eq. (4.23) diverges. We conjecture that, in general, at points $\mathbf{x} \in \partial\Omega$ where the normal is well defined and the approach (4.23) gives a finite pressure, the result obtained according to the latter prescription agrees with the renormalized pressure defined by Eq. (4.22); in fact, this happens in all the examples analysed in this series of papers.

In the rest of the present section, our analysis of the boundary forces will mainly refer to the approach (4.21-4.22).

In applications, one often considers a situation where a quantized field is present both inside Ω and in the complementary region $\Omega^c := \mathbf{R}^d \setminus \Omega$. In this setting the force per unit area acting on the boundary is the resultant of the pressure produced by the field inside Ω , on the one hand, and by the field inside Ω^c , on the other; the renormalized versions of both these observables can be computed, separately, using either one of the two approaches mentioned before.

4.3 Explicit expression for the (regularized) pressure. Let us stick to the viewpoint (4.21-4.22); in order to implement it, we use Eq. (3.33) for the regularized stress-energy tensor that gives the following, for $\mathbf{x} \in \partial\Omega$ (and $\mathbf{n}(\mathbf{x})$ well defined):

$$p_i^u(\mathbf{x}) = \kappa^u \left[\left(\frac{1}{4} - \xi \right) \delta_{ij} \left(D_{\frac{u-1}{2}}(\mathbf{x}, \mathbf{y}) - (\partial^{x^\ell} \partial_{y^\ell} + V(\mathbf{x})) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right) + \right. \\ \left. + \left(\left(\frac{1}{2} - \xi \right) \partial_{x^i y^j} - \xi \partial_{x^i x^j} \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} n^j(\mathbf{x}) . \quad (4.24)$$

To go on, let us restrict the attention to the case of *Dirichlet boundary conditions*; then, only the terms involving mixed derivatives (both with respect to $\mathbf{x}$ and $\mathbf{y}$) of the Dirichlet kernel yield non-vanishing contributions on the boundary $\partial\Omega$. Moreover, the terms proportional to ξ in Eq. (4.24) can be shown to vanish, so that

$$p_i^u(\mathbf{x}) = \kappa^u \left[\left(-\frac{1}{4} \delta_{ij} \partial^{x^\ell} \partial_{y^\ell} + \frac{1}{2} \partial_{x^i y^j} \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} n^j(\mathbf{x}) ; \quad (4.25)$$

see Appendix F for the proof. As a final step, the analytic continuation of $\mathbf{p}^u$ at $u = 0$ must be considered.

4.4 An equivalent characterization of boundary forces. In the literature, forces on $\partial\Omega$ are often characterized by a different approach, which does not require

the knowledge of the full stress-energy tensor; see, e.g., the monographies by Bordag et al. [14, 15], Milton [72] and Plunien et al. [82]. In this approach one considers a variation of the spatial domain Ω controlled by a real parameter, and defines the pressure in terms of the derivative of the bulk energy with respect to the mentioned parameter. For example, if Ω is a parallelepiped $(0, a) \times (0, b) \times (0, c)$ one could consider the variation of the length of any one of its sides, say a ; it is customary to define the total force on the face $\{x^1 = a\}$ as the derivative of the bulk energy with respect to a , with the sign changed.

The idea that boundary forces are related to the variation of Ω has been typically presented in simple examples like the previous one; it can be of interest to propose a general formulation of this idea, and to compare it with the characterization of boundary forces given in subsection 4.2 via the stress-energy tensor.

For the sake of definiteness, let us consider the case where Ω is a bounded domain in $\mathbf{R}^d$ and Dirichlet boundary conditions are prescribed on $\partial\Omega$; besides, let $\mathfrak{S} : \mathbf{R}^d \rightarrow \mathbf{R}^d$ be a vector field. We assume Ω , its boundary $\partial\Omega$ and $\mathfrak{S}$ are regular enough to permit the subsequent calculations.

First of all, consider the family of diffeomorphism

$$\mathbf{S}_\epsilon : \mathbf{R}^d \rightarrow \mathbf{R}^d, \quad \mathbf{x} \mapsto \mathbf{S}_\epsilon(\mathbf{x}) := \mathbf{x} + \epsilon \mathfrak{S}(\mathbf{x}), \quad (4.26)$$

labelled by a small parameter $\epsilon > 0$, that will be ultimately sent to zero. For any ϵ , the spatial domain

$$\Omega_\epsilon := \mathbf{S}_\epsilon(\Omega) \quad (\subset \mathbf{R}^d) \quad (4.27)$$

can be regarded as a deformation of the initial domain Ω .

Of course, a relation analogous to (4.6) holds for the regularized bulk energy E_ϵ^u associated to the deformed domain Ω_ϵ , i.e.,

$$E_\epsilon^u = \frac{\kappa^u}{2} \sum_{k \in \mathcal{K}} (\omega_{\epsilon,k}^2)^{\frac{1-u}{2}}; \quad (4.28)$$

in the above $(\omega_{\epsilon,k}^2)_{k \in \mathcal{K}}$ denote the eigenvalues of the fundamental operator $\mathcal{A}_\epsilon$, that is the operator $-\Delta + V$ acting on the Hilbert space $L^2(\Omega_\epsilon)$ (with Dirichlet boundary conditions on $\partial\Omega_\epsilon$) instead of $L^2(\Omega)$.

Let us now consider the expansion of E_ϵ^u to the first order in ϵ , which describes the variation of the regularized bulk energy under the deformation (4.26) (4.27) of the space domain. Due to Eq. (4.28), the calculation of this expansion can be reduced to the first order expansion of the eigenvalues $\omega_{\epsilon,k}$; this can be done by standard perturbation techniques, as well known from the classic work of Rellich [85]. As illustrated in Appendix F, the conclusion of this analysis is

$$E_\epsilon^u = E^u - \epsilon(1-u) \int_{\partial\Omega} da(\mathbf{x}) \mathfrak{S}^i(\mathbf{x}) p_i^u(\mathbf{x}) + O(\epsilon^2) \quad (4.29)$$

where $\mathbf{p}^u \equiv (p_i^u)$ is the regularized pressure, given by Eq. (F.11). Eq. (4.29) can be used for an alternative, but equivalent definition of the regularised pressure; from this viewpoint, the regularized pressure field $\mathbf{p}^u$ is the unique vector field on $\partial\Omega$ such that (4.29) holds for each one-parameter deformation of the form (4.26-4.27) for the domain Ω .

Let us now perform the analytic continuation up to $u = 0$, *assuming that no pole occurs at this point*; $E_\epsilon^u|_{u=0}$ and $\mathbf{p}^u|_{u=0}$ are the renormalized bulk energy and pressure, and Eq. (4.29) yields the relation

$$E_\epsilon^{ren} = E^{ren} - \epsilon \int_{\partial\Omega} da(\mathbf{x}) \mathfrak{S}^i(\mathbf{x}) p_i^{ren}(\mathbf{x}) + O(\epsilon^2) . \quad (4.30)$$

This is the result anticipated at the beginning of this subsection: a characterization of boundary forces in terms of the of the bulk energy variation under deformations of the domain. We already mentioned the frequent use of this idea in the literature, for particular choices of Ω .

4.5 Integrated forces on the boundary. Let us now discuss the evaluation of the integrated force $\mathfrak{F}_\mathfrak{D}$ acting on an arbitrary subset $\mathfrak{D}$ of the spatial boundary ($\mathfrak{D} \subset \partial\Omega$; possibly, $\mathfrak{D} = \partial\Omega$). As in the case of the pressure considered in subsection 4.2, we can give several alternative definitions of this quantity. As a first possibility, we introduce the regularized total force on $\mathfrak{D}$ (for large $\Re u$)

$$\mathfrak{F}_\mathfrak{D}^u := \int_{\mathfrak{D}} da(\mathbf{x}) \mathbf{p}^u(\mathbf{x}) , \quad (4.31)$$

where $\mathbf{p}^u \equiv (p_i^u(\mathbf{x}))$ indicates the regularized pressure (see Eq. (4.21)); then, we define the *renormalized total force* on $\mathfrak{D}$ as

$$\mathfrak{F}_\mathfrak{D}^{ren} := RP|_{u=0} \mathfrak{F}_\mathfrak{D}^u \quad (4.32)$$

(clearly, when there is no pole in $u = 0$, the above prescription reduces to $\mathfrak{F}_\mathfrak{D}^{ren} := \mathfrak{F}_\mathfrak{D}^u|_{u=0}$, meaning that the analytic continuation in $u = 0$ has to be considered).

Another possibility is to put

$$\mathfrak{F}_\mathfrak{D}^{ren} := \int_{\mathfrak{D}} da(\mathbf{x}) \mathbf{p}^{ren}(\mathbf{x}) , \quad (4.33)$$

where $\mathbf{p}^{ren}(\mathbf{x}) = (p_i^{ren}(\mathbf{x}))$ is the renormalized pressure, defined according either to Eq. (4.22) or to Eq. (4.23).

Similarly to what we pointed out in subsection 4.2 for the pressure, in general the two alternatives (4.32) (4.33) give different results; in fact, the prescription (4.32) always gives a finite result for $\mathfrak{F}_\mathfrak{D}^{ren}$, while (4.33) can give an infinite result.

In conclusion, let us stress that for the integrated force there hold comments analogous to the ones at the end of subsection 4.2, when a quantized field is present both inside Ω and in the complementary region $\Omega^c := \mathbf{R}^d \setminus \Omega$. In this case the total force on any subset $\mathfrak{D} \subset \partial\Omega$ is given by the resultant of the forces corresponding, respectively, to the field inside and outside the spatial domain Ω .

4.6 A comment on some previous “anomalies”. In the previous subsections we have pointed out that the renormalized versions of the total energy, of the pressure and of the integrated forces on the boundary can be defined according to different prescriptions, which in general are not equivalent (see Eq.s (4.13) (4.14), (4.22) (4.23), (4.32) (4.33) and the considerations in the corresponding subsections).

In consequence of this, there arise unavoidable ambiguities, or *anomalies*, when talking about the renormalized observables mentioned above. For example, we have mentioned previously the possible non-equivalence of the alternatives (4.13) (4.14) for the total energy $\mathcal{E}^{ren}$ and (4.22) (4.23) for the pressure $\mathbf{p}^{ren}$; recall that it may happen that the prescriptions (4.14) and (4.23) give infinite results for $\mathcal{E}^{ren}$ and $\mathbf{p}^{ren}$, due to boundary singularities of the stress-energy VEV which make divergent the integral $\int_{\Omega} d\mathbf{x} \langle 0 | \hat{T}_{00}(\mathbf{x}) | 0 \rangle_{ren}$ or the limit $\lim_{\mathbf{x}' \in \Omega, \mathbf{x}' \rightarrow \mathbf{x}} \langle 0 | \hat{T}_{ij}(\mathbf{x}') | 0 \rangle_{ren} n^j(\mathbf{x})$ ($\mathbf{x} \in \partial\Omega$, $i \in \{1, \dots, d\}$).

On the other hand, such boundary singularities of the renormalized stress-energy VEV are not a specific consequence of the zeta regularization; indeed, they also appear if one uses point-splitting, as indicated by the very systematic analysis of Deutsch and Candelas [29]. For the moment, the above mentioned anomalies must be accepted as a problematic aspect of the main regularization schemes; what we can do is just to record them when they appear, and hope that in the future they can be better understood ⁽²²⁾.

5 Some variations of the previous schemes

The variations mentioned in the title are essentially of three kinds, described hereafter in separate subsections. These variations will be mainly used in the applications of Parts II, III and IV; however, the first one will also be relevant for some subcases of the simple application proposed at the end of the present Part I (see subsections 6.8 and 6.9).

²²One should probably look for their origin in some excessive idealization of the physical model (for example, one could try to describe in a more realistic manner the boundaries of the spatial domain; these are “hard” and “deterministic” in the present formulation, but could perhaps be replaced with “soft” or “stochastic” [43] boundaries). For a different type of boundary anomaly, related to vacuum effects for charged fermions, and for its cure taking into account back reaction effects, see [91].

5.1 The basic Hilbert space when 0 is an isolated point of $\sigma(\mathcal{A})$; the case of Neumann and periodic boundary conditions. In this subsection we are going to consider a variation of the framework developed in Sections 2 and 3 to deal with the cases where the fundamental operator $\mathcal{A} = -\Delta + V$ acting on $L^2(\Omega)$ has its spectrum contained in $[0, +\infty)$, with 0 an isolated point; in other terms, $0 \in \sigma(\mathcal{A}) \subset \{0\} \cup [\varepsilon^2, +\infty)$, for some $\varepsilon > 0$. In this case 0 is a proper eigenvalue, as it always occurs for isolated points of the spectrum.

A standard approach employed in the physical literature to treat problems of this kind is to simply neglect the states of “zero energy”; see, e.g., [58, 60, 101]. According to the formulation considered in the present paper, this amounts to the following procedure: in place of $L^2(\Omega)$, we define the basic Hilbert space as the orthogonal complement in $L^2(\Omega)$ of the null eigenspace, that is

$$L_0^2(\Omega) := (\ker \mathcal{A})^\perp \quad (\subset L^2(\Omega)) . \quad (5.1)$$

It should be noted that the restriction of $\mathcal{A}$ to $L_0^2(\Omega)$ is a selfadjoint, strictly positive operator in $L_0^2(\Omega)$ with spectrum contained in $[\varepsilon^2, +\infty)$. In this situation, $L_0^2(\Omega)$ is the basic Hilbert space even from the viewpoint of field quantization ⁽²³⁾.

Let us recall the definition (3.1), giving the integral kernel associated to a given operator on $L^2(\Omega)$, and consider Eq.s (3.17), (3.46) and (3.53) for the Dirichlet, heat, cylinder and modified cylinder kernels associated to $\mathcal{A}$; if the latter operator is redefined as the restriction of $-\Delta + V$ to $L_0^2(\Omega)$, in the cited equations we should formally replace $\delta_{\mathbf{x}}$, $\delta_{\mathbf{y}}$ with $E_0\delta_{\mathbf{x}}$, $E_0\delta_{\mathbf{y}}$ where E_0 is the orthogonal projection onto $L_0^2(\Omega)$ (suitably extended to distributions, so that it can be applied to $\delta_{\mathbf{x}}, \delta_{\mathbf{y}}$). With this modification, the expansions (3.18), (3.47), (3.48) and (3.53) for the kernels mentioned above hold again, using the eigenfunctions of $\mathcal{A}$ in $L_0^2(\Omega)$ ⁽²⁴⁾.

Typical configurations of the above type are those where $\mathcal{A} = -\Delta$, the spatial domain Ω is bounded and the field fulfills either Neumann or periodic boundary conditions on $\partial\Omega$ ⁽²⁵⁾; indeed, in such cases the spectrum of $\mathcal{A}$ in $L^2(\Omega)$ is purely discrete, 0 is an eigenvalue and $\ker \mathcal{A}$ is formed by the constant functions. Therefore $L_0^2(\Omega)$, defined via Eq. (5.1), is formed by the functions with mean zero:

$$L_0^2(\Omega) = \left\{ f \in L^2(\Omega) \mid \int_{\Omega} d\mathbf{x} f(\mathbf{x}) = 0 \right\} . \quad (5.2)$$

²³By this, we mean that the Fock space $\mathfrak{F}$ of the quantized scalar field living in Ω is the direct sum of all symmetrized tensor powers of $L_0^2(\Omega)$.

²⁴As an example, in the case described by Eq. (5.2) the projection E_0 onto $L_0^2(\Omega)$ is given by $E_0 f = f - \frac{1}{\text{Vol}(\Omega)} \int_{\Omega} d\mathbf{x} f(\mathbf{x})$ ($\text{Vol}(\Omega)$ is the volume of Ω); the previous prescription makes sense as well for $f = \delta_{\mathbf{x}}$ and gives $E_0\delta_{\mathbf{x}} = \delta_{\mathbf{x}} - \frac{1}{\text{Vol}(\Omega)}$.

²⁵As will be observed in subsection 5.3, the case of periodic boundaries would be more properly formulated in terms of a free field on a torus, but this is cause of no concern for the present considerations.

Let us mention that an analogous framework can be considered for slab configurations where $\Omega = \Omega_1 \times \mathbf{R}^{d_2}$ and Neumann or periodic boundary conditions are prescribed on $\partial\Omega_1 \times \mathbf{R}^{d_2}$. In these cases one works with the reduced operator $\mathcal{A}_1$ acting in $L^2(\Omega_1)$; the latter must then be replaced with the Hilbert space

$$L_0^2(\Omega_1) := (\ker \mathcal{A}_1)^\perp = \left\{ f \in L^2(\Omega_1) \mid \int_{\Omega_1} d\mathbf{x}_1 f(\mathbf{x}_1) = 0 \right\} \quad (5.3)$$

and the basic Hilbert space for the full theory on Ω is $L_0^2(\Omega_1) \otimes L^2(\mathbf{R}^{d_2})$.

In the applications to be considered in the following, whenever 0 is an isolated point of the spectrum we will always assume that the fundamental operator $\mathcal{A}$ (resp. $\mathcal{A}_1$) has been redefined so that it acts on the Hilbert space $L_0^2(\Omega)$ of Eq. (5.1) (resp. $L_0^2(\Omega_1)$ of Eq. (5.3)).

5.2 The case where 0 is in the continuous spectrum of $\mathcal{A}$. Let us pass to the case where the fundamental operator $\mathcal{A} = -\Delta + V$ is non-negative ($\sigma(\mathcal{A}) \subset [0, +\infty)$), and 0 is in the continuous spectrum of $\mathcal{A}$; then 0 has zero spectral measure and is a non isolated point of the spectrum (otherwise, it would be a proper eigenvalue).

Here are two examples of this kind. To obtain them we consider the operator $\mathcal{A} := -\Delta$ in $L^2(\mathbf{R}^d)$, or the operator $\mathcal{A} := -\Delta$ in $L^2(\Omega)$ where Ω is the half-space $\{\mathbf{x} \in \mathbf{R}^d \mid x^1 > 0\}$, and suitable boundary conditions, say Dirichlet, are specified on $\partial\Omega = \{x^1 = 0\}$. In these cases $\mathcal{A}$ has a complete orthonormal system of (improper) eigenfunctions $(F_{\mathbf{k}})_{\mathbf{k} \in \mathcal{K}}$ with corresponding eigenvalues $\omega_{\mathbf{k}}^2$, where: in the first case, $\mathcal{K} = \mathbf{R}^d$ (with the Lebesgue measure $d\mathbf{k}$), $\mathcal{F}_{\mathbf{k}}(\mathbf{x}) := (2\pi)^{-d/2} e^{i\mathbf{k} \cdot \mathbf{x}}$, $\omega_{\mathbf{k}} := |\mathbf{k}|$; in the second case, $\mathcal{K} = (0, +\infty) \times \mathbf{R}^{d-1}$ (again, with the Lebesgue measure $d\mathbf{k}$, $\mathcal{F}_{\mathbf{k}}(\mathbf{x}) := \sqrt{2}(2\pi)^{-d/2} \sin(k^1 x^1) e^{ik^2 x^2 + \dots + k^d x^d}$ and, again, $\omega_{\mathbf{k}} := |\mathbf{k}|$). In both cases $\sigma(\mathcal{A}) = [0, +\infty)$ and the spectrum is purely continuous.

The case of $\mathcal{A}$ non-negative, with 0 in the continuous spectrum, cannot be treated with the approach of the previous subsection: there is no way to obtain a strictly positive operator by simply removing 0 from the spectrum. In a more physical language, infrared divergences cannot be simply ignored and we must devise a more sophisticated way to treat them, as we are currently doing for ultraviolet divergences. A natural approach to the problem is to represent $\mathcal{A}$ as a limit

$$\mathcal{A} := \lim_{\varepsilon \rightarrow 0^+} \mathcal{A}_\varepsilon \quad (5.4)$$

where $\mathcal{A}_\varepsilon$ is a selfadjoint operator, depending on a parameter $\varepsilon \in (0, \varepsilon_0)$ and such that the spectrum of $\mathcal{A}_\varepsilon$ is contained in $[\varepsilon^2, +\infty)$; the deformed operator $\mathcal{A}_\varepsilon$ is used everywhere in place of $\mathcal{A}$, and the limit $\varepsilon \rightarrow 0^+$ is performed only at the end, after zeta renormalization has been carried out. In particular, we define the *deformed smeared field operator*

$$\widehat{\phi}^{\varepsilon u} := (\kappa^{-2} \mathcal{A}_\varepsilon)^{-u/4} \widehat{\phi} \quad (5.5)$$

and the corresponding *deformed, regularized stress-energy tensor* $\widehat{T}_{\mu\nu}^{\varepsilon u}(x)$ whose VEV is given by

$$\begin{aligned} \langle 0 | \widehat{T}_{\mu\nu}^{\varepsilon u}(x) | 0 \rangle = & \left(\frac{1}{2} - \xi \right) (\partial_{x^\mu} y^\nu + \partial_{x^\nu} y^\mu) - \left(\frac{1}{2} - 2\xi \right) \eta_{\mu\nu} \left(\partial^{x^\lambda} \partial_{y^\lambda} + V \right) - \xi (\partial_{x^\mu} x^\nu + \partial_{y^\mu} y^\nu) \Big|_{y=x} \cdot \\ & \cdot \langle 0 | \widehat{\phi}^{\varepsilon u}(x) \widehat{\phi}^{\varepsilon u}(y) | 0 \rangle ; \end{aligned} \quad (5.6)$$

for the above VEV we have expression analogous to (3.31-3.33) in terms of the *deformed Dirichlet kernel*

$$D_s^\varepsilon(\mathbf{x}, \mathbf{y}) := \mathcal{A}_\varepsilon^{-s}(\mathbf{x}, \mathbf{y}) = \langle \delta_{\mathbf{x}} | \mathcal{A}_\varepsilon^{-s} \delta_{\mathbf{y}} \rangle , \quad (5.7)$$

with $s = (u \pm 1)/2$.

As mentioned above, in this generalized version of the local zeta regularization the limit $\varepsilon \rightarrow 0^+$ must be considered only after the analytic continuation has been performed; in particular, we define

$$\langle 0 | \widehat{T}_{\mu\nu}(x) | 0 \rangle_{ren} := \lim_{\varepsilon \rightarrow 0^+} RP \Big|_{u=0} \langle 0 | \widehat{T}_{\mu\nu}^{\varepsilon u}(x) | 0 \rangle . \quad (5.8)$$

The above renormalized VEV can be expressed in terms of the renormalized kernels $D_{\pm 1/2}^{(\kappa)}(\mathbf{x}, \mathbf{y})$ and $\partial_{zw} D_{1/2}^{(\kappa)}(\mathbf{x}, \mathbf{y})$, where

$$D_{\pm \frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) := \lim_{\varepsilon \rightarrow 0^+} RP \Big|_{u=0} \left(\kappa^u D_{\frac{u \pm 1}{2}}^\varepsilon(\mathbf{x}, \mathbf{y}) \right) = \lim_{\varepsilon \rightarrow 0^+} RP \Big|_{s=\pm \frac{1}{2}} \left(\kappa^{2s \mp 1} D_s^\varepsilon(\mathbf{x}, \mathbf{y}) \right) , \quad (5.9)$$

$$\partial_{zw} D_{\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) := \lim_{\varepsilon \rightarrow 0^+} RP \Big|_{u=0} \left(\kappa^u \partial_{zw} D_{\frac{u+1}{2}}^\varepsilon(\mathbf{x}, \mathbf{y}) \right) = \lim_{\varepsilon \rightarrow 0^+} RP \Big|_{s=\frac{1}{2}} \left(\kappa^{2s-1} \partial_{zw} D_s^\varepsilon(\mathbf{x}, \mathbf{y}) \right) ;$$

these functions play a role very similar to the ones introduced in Eq. (3.37) for a strictly positive $\mathcal{A}$, and allow to express the renormalized VEV (5.8) as in Eq.s (3.39-3.41).

In the sequel we will write K^ε and T^ε (or $\tilde{T}^\varepsilon$), respectively, for the heat and cylinder (or modified cylinder) kernel associated to $\mathcal{A}_\varepsilon$. Proceeding as in Section 3 we obtain, for $\Re s$ sufficiently large,

$$D_s^\varepsilon(\mathbf{x}, \mathbf{y}) = \frac{1}{\Gamma(s)} \int_0^{+\infty} dt \, t^{s-1} K^\varepsilon(t; \mathbf{x}, \mathbf{y}) ; \quad (5.10)$$

$$D_s^\varepsilon(\mathbf{x}, \mathbf{y}) = \frac{1}{\Gamma(2s)} \int_0^{+\infty} dt \, t^{2s-1} T^\varepsilon(t; \mathbf{x}, \mathbf{y}) ; \quad (5.11)$$

the above formulas are the starting point to discuss the analytic continuation in s of the Dirichlet kernel D_s^ε , for any fixed $\varepsilon \in (0, \varepsilon_0)$.

Let us associate to $\mathcal{A}$ to the “undeformed” fundamental operator $\mathcal{A}$ the heat and cylinder kernels

$$K(t; \mathbf{x}, \mathbf{y}) := e^{-t\mathcal{A}}(\mathbf{x}, \mathbf{y}) , \quad T(t; \mathbf{x}, \mathbf{y}) := e^{-t\sqrt{\mathcal{A}}}(\mathbf{x}, \mathbf{y}) , \quad (5.12)$$

as well as the modified cylinder kernel

$$\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) := (\sqrt{\mathcal{A}}^{-1} e^{-\mathbf{t}\sqrt{\mathcal{A}}})(\mathbf{x}, \mathbf{y}) \quad (5.13)$$

(see subsection 3.10); these can be represented as in Eq.s (3.47), (3.48) and (3.53) in terms of the eigenfunctions $(F_k)_{k \in \mathcal{K}}$ and eigenvalues $(\omega_k)_{k \in \mathcal{K}}$ of $\mathcal{A}$. We stress that the functions $D_{\pm 1/2}^{(\kappa)}$ of Eq. (5.9) do *not* possess integral representations of the form (5.10) (5.11) with $K^\varepsilon, T^\varepsilon$ replaced by K, T ; in fact, the corresponding integrals for K, T are typically divergent. In the sequel, we will present a more subtle way to obtain $D_{\pm 1/2}^{(\kappa)}$ from K or T (and $\tilde{T}$).

Up to now we have not specified any particular form for $\mathcal{A}_\varepsilon$. The following two choices seem to be natural:

$$\mathcal{A}_\varepsilon := \mathcal{A} + \varepsilon^2 , \quad (5.14)$$

$$\mathcal{A}_\varepsilon := (\sqrt{\mathcal{A}} + \varepsilon)^2 . \quad (5.15)$$

The first one corresponds to the idea, widespread in the physical literature, to treat infrared divergences adding a small mass ε [69, 93]; the second one is less familiar and is justified by the considerations that follow.

Assuming $\mathcal{A}_\varepsilon$ to have either the form (5.14) or (5.15), we readily obtain the following relations allowing to express the deformed kernels $K^\varepsilon, T^\varepsilon$ in terms of the analogous basic kernels K, T :

$$\mathcal{A}_\varepsilon := \mathcal{A} + \varepsilon^2 \quad \Rightarrow \quad K^\varepsilon(\mathbf{t}; \mathbf{x}, \mathbf{y}) = e^{-\varepsilon^2 \mathbf{t}} K(\mathbf{t}; \mathbf{x}, \mathbf{y}) ; \quad (5.16)$$

$$\mathcal{A}_\varepsilon := (\sqrt{\mathcal{A}} + \varepsilon)^2 \quad \Rightarrow \quad T^\varepsilon(\mathbf{t}; \mathbf{x}, \mathbf{y}) = e^{-\varepsilon \mathbf{t}} T(\mathbf{t}; \mathbf{x}, \mathbf{y}) . \quad (5.17)$$

In particular, assuming the kernels K, T to be meromorphic functions of $\mathbf{t}$, the above relations imply that the deformed kernels $K^\varepsilon, T^\varepsilon$ are meromorphic as well; in these cases, the deformed Dirichlet kernel D_s^ε admits integral representations analogous to (3.94) (3.95), involving the Hankel contour $\mathfrak{H}$. For example, one can write

$$D_s^\varepsilon(\mathbf{x}, \mathbf{y}) = \frac{e^{-2i\pi s} \Gamma(1-2s)}{2\pi i} \int_{\mathfrak{H}} d\mathbf{t} \, \mathbf{t}^{2s-1} T^\varepsilon(\mathbf{t}; \mathbf{x}, \mathbf{y}) . \quad (5.18)$$

Starting from the above representation we can derive explicit expressions for the renormalized functions $D_{\pm 1/2}^{(\kappa)}(\mathbf{x}, \mathbf{y})$, and, more generally, for

$$D_{-\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) := \lim_{\varepsilon \rightarrow 0} RP \Big|_{s=-\frac{n}{2}} \left(\kappa^{2s+n} D_s^\varepsilon(\mathbf{x}, \mathbf{y}) \right) \quad (n \in \{-1, 0, 1, 2, \dots\}) , \quad (5.19)$$

that could be called “renormalized Dirichlet kernels” of order $-n/2$. More precisely, let us assume the modified cylinder kernel $\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y})$ associated to the fundamental operator $\mathcal{A}$ to be a meromorphic function of $\mathbf{t}$ in the neighborhood of the positive real half-axis, fulfilling the bound

$$|\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y})| \leq C |\mathbf{t}|^{-a-n+1} \quad \text{for } \Re \mathbf{t} \rightarrow +\infty \text{ and some } C, a > 0. \quad (5.20)$$

Then, we obtain the following result, for $n = -1, 0, 1, 2, \dots$ (see Appendix G):

$$D_{-\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) = (-1)^{n+1} \Gamma(n+2) \text{Res} \left(\mathbf{t}^{-(n+2)} \tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}); 0 \right). \quad (5.21)$$

Let us remark that Eq. (5.21) has the same structure of Eq. (3.98), dealing with the Dirichlet kernel when $\mathcal{A}$ is strictly positive. In the strictly positive case, the cited result was derived rigorously (from Eq. (3.97)), with no need to introduce a regulating parameter ε ; in the present framework, instead, it would be impossible to establish (5.21) without using the regulator ε .

One could derive results similar to Eq. (5.21), involving the “renormalized derivatives”, e.g.,

$$\partial_{zw} D_{-\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) := \lim_{\varepsilon \rightarrow 0} RP \Big|_{s=-\frac{n}{2}} \left(\kappa^{2s+n} \partial_{zw} D_s^\varepsilon(\mathbf{x}, \mathbf{y}) \right), \quad (5.22)$$

where z, w are spatial variables; indeed, for $n = -1, 0, 1, 2, \dots$, we have

$$\partial_{zw} D_{-\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) = (-1)^{n+1} \Gamma(n+2) \text{Res} \left(\mathbf{t}^{-(n+2)} \partial_{zw} \tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}); 0 \right) \quad (5.23)$$

if $\partial_{zw} \tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y})$, as a function of $\mathbf{t}$, fulfills conditions of the form stipulated previously for $\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y})$ (see, in particular, Eq. (5.20)).

To conclude, let us discuss the pressure on the boundary in the present framework; the main point is the fact that, as in the case of strictly positive $\mathcal{A}$, there are two possible prescriptions for the renormalized pressure. The first alternative is to introduce, at each point $\mathbf{x} \in \partial\Omega$, a *deformed, regularized pressure* with components

$$p_i^{\varepsilon u}(\mathbf{x}) := \langle 0 | \hat{T}_{ij}^{\varepsilon u}(\mathbf{x}) | 0 \rangle n^j(\mathbf{x}) \quad (i \in \{1, \dots, d\}), \quad (5.24)$$

where $\mathbf{n}(\mathbf{x}) \equiv (n^j(\mathbf{x}))$ is the outer unit normal to the boundary; we then define the renormalized pressure at $\mathbf{x}$ as

$$p_i^{ren}(\mathbf{x}) := \lim_{\varepsilon \rightarrow 0^+} RP \Big|_{u=0} p_i^{\varepsilon u}(\mathbf{x}). \quad (5.25)$$

The second alternative is to put

$$p_i^{ren}(\mathbf{x}) := \left(\lim_{\mathbf{x}' \in \Omega, \mathbf{x}' \rightarrow \mathbf{x}} \langle 0 | \hat{T}_{ij}(\mathbf{x}') | 0 \rangle_{ren} \right) n^j(\mathbf{x}) \quad (5.26)$$

where $\langle 0 | \hat{T}_{ij}(\mathbf{x}') | 0 \rangle_{ren}$ is defined according to Eq. (5.8) at all interior points $\mathbf{x}'$ of the domain. The prescriptions (5.25) (5.26) can, in general, give different results: this happens, for example, in the case of a massless field on a wedge-shaped domain, to be discussed in Section 5 of Part II.

5.3 Some variations involving the spatial domain. In the literature, a scalar field fulfilling periodic boundary conditions is often considered. To give a rigorous description of this configuration, one should better give up to viewing Ω as an open subset of $\mathbf{R}^d$ and pass to a description in terms of tori. For example, it is customary to speak of a field on the hypercube $(0, a)^d$ with periodic boundary conditions, where $a > 0$ is some given length. In the most precise description of this configuration, Ω is not $(0, a)^d$ but rather the d -dimensional torus $\mathbf{T}_a^d := \mathbf{R}^d / (a\mathbf{Z})^d \simeq (\mathbf{R}/a\mathbf{Z})^d$ (where $\mathbf{Z}$ is the set of integers, so that $a\mathbf{Z} = \{\dots, -2a, -a, 0, a, 2a, \dots\}$)⁽²⁶⁾.

In some applications to be considered in the subsequent Parts II and III, the space domain Ω is an open subset of $\mathbf{R}^d$ but, in place of the Cartesian coordinates $\mathbf{x} \equiv (x^i)$, it is natural to use for it some curvilinear coordinates $(q^i)_{i=1, \dots, d} \equiv \mathbf{q}$; in these coordinates, the line element of $\mathbf{R}^d$ will have the form

$$d\ell^2 = a_{ij}(\mathbf{q}) dq^i dq^j . \quad (5.27)$$

The above spatial coordinates induce a set of spacetime coordinates $(q^\mu)_{\mu=0, \dots, d} \equiv q$ on $\mathbf{R} \times \Omega$ where $q^0 := t$ and the q^i 's are as before; clearly, the spacetime line element $ds^2 = -dt^2 + d\ell^2$ will have the form

$$ds^2 = g_{\mu\nu}(q) dq^\mu dq^\nu , \quad (5.28)$$

$$g_{00} := -1 , \quad g_{i0} = g_{0i} := 0 , \quad g_{ij}(q) := a_{ij}(\mathbf{q}) \quad \text{for } i, j \in \{1, \dots, d\} .$$

The analogue of Eq. (2.16) in the coordinate system (q^μ) is

$$\widehat{T}_{\mu\nu}^u := (1-2\xi)\partial_\mu \widehat{\phi}^u \circ \partial_\nu \widehat{\phi}^u - \left(\frac{1}{2} - 2\xi\right)\eta_{\mu\nu} \left(\partial^\lambda \widehat{\phi}^u \partial_\lambda \widehat{\phi}^u + V(\widehat{\phi}^u)^2\right) - 2\xi \widehat{\phi}^u \circ \nabla_{\mu\nu} \widehat{\phi}^u , \quad (5.29)$$

where ∇_μ is the covariant derivative induced by the (flat) spacetime metric (5.28). In principle, the covariant derivative ∇_μ should appear in place of any derivative ∂_μ ; however we are working with a *scalar* field and it is well-known that

$$\nabla_\mu f = \partial_\mu f \quad \text{if } f \text{ is a scalar function} . \quad (5.30)$$

The situation is different when we consider second order derivatives, which explains the appearing of $\nabla_{\mu\nu}$ in (5.29). Let us recall that

$$\nabla_{\mu\nu} f = \partial_{\mu\nu} f - \Gamma_{\mu\nu}^\lambda \partial_\lambda f \quad (= \nabla_{\nu\mu} f) \quad \text{if } f \text{ is a scalar function} , \quad (5.31)$$

²⁶The considerations of subsection 5.1 for the periodic case are easily rephrased in terms of the torus $\mathbf{T}_a^d$. The operator $\mathcal{A} := -\Delta$ acting in $L^2(\mathbf{T}_a^d)$ has 0 as an eigenvalue, with $\ker \mathcal{A}$ formed by the constant functions; again, 0 is eliminated viewing $\mathcal{A}$ as an operator acting in

$$L_0^2(\mathbf{T}_a^d) := (\ker \mathcal{A})^\perp = \left\{ f \in L^2(\mathbf{T}_a^d) \mid \int_{\mathbf{T}_a^d} d\mathbf{x} f(\mathbf{x}) = 0 \right\} .$$

where we are using the spacetime Christoffel symbols $\Gamma_{\mu\nu}^{\lambda} := \frac{1}{2}g^{\lambda\rho}(\partial_{\mu}g_{\rho\nu} + \partial_{\nu}g_{\mu\rho} - \partial_{\rho}g_{\mu\nu})$. The above computational rule is more efficiently implemented recalling that $q^0 = t$ and using the space covariant derivatives D_i corresponding to the line element (5.27); these rely on the Christoffel symbols $\gamma_{ij}^k := \frac{1}{2}a^{kh}(\partial_i a_{hj} + \partial_j a_{ih} - \partial_h a_{ij})$. From Eq. (5.28) one easily infers that $\Gamma_{ij}^k = \gamma_{ij}^k$ (for $i, j, k \in \{1, \dots, d\}$) are the only non-vanishing coefficients; so, Eq. (5.31) for a scalar function f on spacetime implies

$$\begin{aligned} \nabla_{ij}f &= D_{ij}f = \partial_{ij}f - \gamma_{ij}^k \partial_k f , \\ \nabla_{0i}f &= \partial_0(\partial_i f) = \partial_i(\partial_0 f) = \nabla_{i0}f , \quad \nabla_{00}f = \partial_{00}f . \end{aligned} \quad (5.32)$$

As a further variation of our schemes, we can stipulate the spatial domain Ω to be an arbitrary d -dimensional Riemannian manifold, possibly non flat; in any coordinate system $(q^i)_{i=1,\dots,d} \equiv \mathbf{q}$ of Ω , the Riemannian line element $d\ell^2$ will have a representation of the form (5.27). (Of course the position $\Omega = \mathbf{T}_a^d$, considered at the beginning of this paragraph in relation to periodic boundary conditions, amounts to choosing for Ω a very simple, flat Riemannian manifold). Given any Riemannian manifold Ω , we can associate to it the spacetime $\mathbf{R} \times \Omega$ equipped with the line element $ds^2 = -dt^2 + d\ell^2$; this takes the form (5.28) in coordinates $(t, q^i) \equiv (q^0, q^i) \equiv (q^{\mu})_{\mu=0,\dots,d}$.

As a final variation, we can assume the space domain Ω to be an open subset of a Riemannian manifold and prescribe boundary conditions on $\partial\Omega$.

Many results in Sections 2, 3 and 4 are readily adapted to the variations considered in this subsection for the space domain. An essential point in making these adaptations is to remember that, when an arbitrary coordinate system is employed, the second order derivatives of scalar functions must be intended in a covariant sense and the computational rules (5.32) must be applied.

6 The case of a massless field on the segment

6.1 Introducing the problem for arbitrary boundary conditions. To conclude the present Part I we present a simple application of our general formalism, namely a 1-dimensional model describing a massless scalar field living on a segment, with no background potential. This means that

$$d = 1 , \quad \Omega = (0, a) \quad (a > 0) , \quad \mathcal{A} = -\partial_{x^1 x^1} \quad (V = 0) \quad (6.1)$$

where $x^1 \in (0, a)$ is the standard Cartesian coordinate ⁽²⁷⁾. The field is assumed to fulfill Dirichlet, Neumann or periodic boundary conditions at the boundary $\partial\Omega = \{0\} \cup \{a\}$; we will deal with each one of these alternatives separately, in the subsequent subsections 6.6-6.9.

²⁷We could have used, in place of x^1 , the rescaled spatial variable

$$x_{\star}^1 := x^1/a \in (0, 1) ,$$

in terms of which, we would have obtained simpler expressions for the results reported in sequel. Yet, we choose not to employ this rescaled coordinate in order to make the comparison with known results more straightforward.

In passing, we note that the setting described above is the $d = 1$ case both for the configuration with two parallel hyperplanes and for the d -dimensional box (to be considered in Parts II and IV, respectively).

Let us make some comparison with the previous literature about the Casimir effect for a scalar field on a segment. First of all, we wish to mention the book of Bordag et al. [15] (see Chapter 2) and the work by Fulling et al. [50]; these authors derive the total bulk energy, for several boundary conditions, using regularization methods different from zeta approach. More precisely, [15] uses an exponential cut-off regularization followed by Abel-Plana resummation, while [50] employs essentially a point-splitting procedure. These authors also obtain the force acting on the end-points of the segment by differentiating the expression for the total energy with respect to the length of the segment (see the comments at the beginning of subsection 4.4). Let us also mention the paper [68] by Mamaev and Trunov deriving the stress-energy VEV for a massless scalar field on a segment in the case of periodic boundary conditions, via point-splitting regularization.

In all cases where the present section has an intersection with [15, 50, 68], our results are in agreement with these references ⁽²⁸⁾.

²⁸*Note added.* After this manuscript was written, we became aware of a very recent paper by Mera and Fulling [70] who consider a *massive* scalar field on a segment, regularized via an exponential cutoff, and compute the stress-energy VEV by the method of images, i.e., as a sum over infinitely many optical paths. (Even though different from zeta regularization, this approach is related to the cylinder kernel due to the cutoff structure.) The authors of [70] consider the zero cutoff limit, whose treatment can be understood as a renormalization, and also give a number of results on the zero mass limit. With a small addition to these results, presented hereafter, our renormalized stress-energy VEV can be shown to agree with the zero mass limit of the calculations in [70]. As an example let us consider the $(0, 0)$ component of the stress-energy VEV, i.e., the energy density, in the case of Neumann boundary conditions; in [70] this is expressed as the sum of a term diverging quadratically in the cutoff, and of other terms which have a finite limit when the cutoff is sent to zero. If one renormalizes removing the divergent term, one obtains from [70] that

$$\begin{aligned} & \langle 0 | T_{00}(x^1) | 0 \rangle_{ren} = \\ & = -\frac{m}{2\pi a} \sum_{n=1}^{+\infty} \frac{K_1(2mna)}{n} - \frac{m\xi}{\pi} \sum_{n=-\infty}^{+\infty} \frac{K_1(2m|x^1 + na|)}{|x^1 + na|} - \frac{2m^2}{\pi} \left(\xi - \frac{1}{4}\right) \sum_{n=-\infty}^{+\infty} K_0(2m|x^1 + na|) \end{aligned}$$

where m is the field mass and K_0, K_1 are modified Bessel functions of the second kind (see Eq.s (10) (15-17) and (51) of [70], here re-written with $L = a$, $\beta = \xi - 1/4$ and putting $r = l = 0$ to indicate the choice of Neumann boundary conditions). It is known that $K_0(z) = -\ln z + O(1)$ and $K_1(z) = 1/z + O(z \ln z)$ for $z \rightarrow 0^+$. So one expects that, for $m \rightarrow 0^+$, $m \sum_{n=1}^{+\infty} K_1(2mna)/n \rightarrow (\sum_{n=1}^{+\infty} 1/n^2)/(2a) = \pi^2/(12a)$ and $m^2 \sum_{n=-\infty}^{+\infty} K_0(2m|x^1 + na|) \rightarrow 0$; these two facts are in fact established in [70]. Let us add to these results the remark that, for $m \rightarrow 0^+$, $m \sum_{n=-\infty}^{+\infty} K_1(2m|x^1 + na|)/|x^1 + na| \rightarrow (1/2) \sum_{n=-\infty}^{+\infty} 1/(x^1 + na)^2 = \pi^2/(2a^2 \sin^2(\frac{\pi}{a}x^1))$ (for the computation of the last series by contour integral methods see [59], p. 268, Eq. 4.9-4). In view of these facts

$$\lim_{m \rightarrow 0^+} \langle 0 | T_{00}(x^1) | 0 \rangle_{ren} = -\frac{\pi}{24a^2} - \xi \frac{\pi}{2a^2 \sin^2(\frac{\pi}{a}x^1)} ,$$

in agreement with Eq. (6.40) of the present work. Let us emphasize that, differently from [70], our zeta approach gives the renormalized stress-energy VEV by mere analytic continuation, with no need to remove divergent terms.

6.2 Cylinder and Dirichlet kernels. For any one of the previously mentioned boundary conditions, we perform our analysis in the manner explained hereafter. First of all, we determine explicitly the cylinder kernel $T(\mathbf{t}; x^1, y^1)$ associated to the fundamental operator $\mathcal{A}$; to this purpose we consider a complete orthonormal set of eigenfunctions $(F_k)_{k \in \mathcal{K}}$ for $\mathcal{A}$ with eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$. The label set $\mathcal{K}$ is countable and $\int_{\mathcal{K}} dk$ means $\sum_{k \in \mathcal{K}}$; so, the eigenfunction expansion (3.48) for the cylinder kernel reads

$$T(\mathbf{t}; x^1, y^1) = \sum_{k \in \mathcal{K}} e^{-\omega_k \mathbf{t}} F_k(x^1) \overline{F_k}(y^1) . \quad (6.2)$$

Once the cylinder kernel has been computed explicitly by evaluating the above sum, we can proceed to determine the modified cylinder kernel $\tilde{T}(\mathbf{t}; x^1, y^1)$ as the primitive of $-T(\mathbf{t}; x^1, y^1)$ vanishing for $\mathbf{t} \rightarrow +\infty$ (see Eq. (3.55)).

In the next subsections T and $\tilde{T}$ will be computed explicitly, for several kinds of boundary conditions. In all cases, the cylinder kernel T and the *spatial derivatives* of the modified cylinder kernel $\tilde{T}$ will be found to have meromorphic extensions in the $\mathbf{t}$ variable to an open complex neighborhood of $[0, +\infty)$, with possible poles only at $\mathbf{t} = 0$, vanishing exponentially for $\Re \mathbf{t} \rightarrow +\infty$; thus, the framework of subsection 3.15 can be applied straightforwardly.

To be more precise: for $y^1 \neq x^1$, $T, \tilde{T}$ and their derivatives have analytic extensions in $\mathbf{t}$ to a neighborhood of $[0, +\infty)$. When evaluated on the diagonal $y^1 = x^1$, the modified cylinder kernel $\tilde{T}$ is found to have a logarithmic singularity in $\mathbf{t} = 0$, while the cylinder kernel T and the spatial derivatives of both T and $\tilde{T}$ are meromorphic in a neighborhood of $[0, +\infty)$ with only a pole singularity in $\mathbf{t} = 0$. Because of this, one can resort to Eq.s (3.95) and (3.97) to obtain the analytic continuation of the Dirichlet kernel and of its derivatives, required in order to determine the regularized VEV of the stress-energy tensor; explicitly, we have

$$D_{\frac{u-1}{2}}(x^1, y^1) \Big|_{y^1=x^1} = \frac{e^{-i\pi(u-1)} \Gamma(2-u)}{2\pi i} \int_{\mathfrak{H}} d\mathbf{t} \, \mathbf{t}^{u-2} T(\mathbf{t}; x^1, y^1) \Big|_{y^1=x^1} ; \quad (6.3)$$

$$\partial_{zw} D_{\frac{u+1}{2}}(x^1, y^1) \Big|_{y^1=x^1} = -\frac{e^{-i\pi(u+1)} \Gamma(1-u)}{2\pi i} \int_{\mathfrak{H}} d\mathbf{t} \, \mathbf{t}^{u-1} \partial_{zw} \tilde{T}(\mathbf{t}; x^1, y^1) \Big|_{y^1=x^1} \quad (6.4)$$

for $z, w \in \{x^1, y^1\}$.

In order to obtain the analytic continuations at $u = 0$ of the above functions, one can simply set $u = 0$ in the expressions on the right-hand sides of Eq.s (6.3) (6.4) and explicitly evaluate the remaining integrals along the Hankel contour via the

residue theorem ⁽²⁹⁾; as indicated in subsection 3.15, this gives

$$D_{-\frac{1}{2}}(x^1, y^1) \Big|_{y^1=x^1} = -\text{Res} \left(\mathfrak{t}^{-2} T(\mathfrak{t}; x^1, y^1) \Big|_{y^1=x^1}; 0 \right); \quad (6.5)$$

$$\partial_{zw} D_{\frac{1}{2}}(x^1, y^1) \Big|_{y^1=x^1} = \text{Res} \left(\mathfrak{t}^{-1} \partial_{zw} \tilde{T}(\mathfrak{t}; x^1, y^1) \Big|_{y^1=x^1}; 0 \right) \quad \text{for } z, w \in \{x^1, y^1\} \quad (6.6)$$

(use Eq. (3.96) and the analogue of Eq. (3.98) for the derivatives of D_s).

Before moving on, let us mention that analogous considerations can be made concerning the traces $\text{Tr } \mathcal{A}^{-s}$, $T(\mathfrak{t})$ (see Eq. (3.22) and Eq.s (3.56) (3.57), respectively). Indeed, we can compute the cylinder trace $T(\mathfrak{t})$ according to Eq. (3.58) that in the present case reads

$$T(\mathfrak{t}) = \sum_{k \in \mathcal{K}} e^{-\omega_k \mathfrak{t}}. \quad (6.7)$$

By explicit evaluation of the above sum for the boundary conditions considered in the sequel, it becomes apparent that $T(\mathfrak{t})$ possesses the same features of its local counterpart. Thus, we can resort again to the general framework of subsection 3.15 to obtain the analytic continuation of $\text{Tr } \mathcal{A}^{-s}$; in particular, due to Eq. (3.99), the continuation at $s = -1/2$ is

$$\text{Tr } \mathcal{A}^{1/2} = -\text{Res} \left(\mathfrak{t}^{-2} T(\mathfrak{t}); 0 \right). \quad (6.8)$$

6.3 The stress-energy tensor. We can now determine explicitly the renormalized VEV of the stress-energy tensor; in fact, since no singularity arises, Eq.s (3.39-3.41) and (3.42) imply

$$\langle 0 | \widehat{T}_{00}(x^1) | 0 \rangle_{ren} = \left[\left(\frac{1}{4} + \xi \right) D_{-\frac{1}{2}}(x^1, y^1) + \left(\frac{1}{4} - \xi \right) \partial_{x^1 y^1} D_{\frac{1}{2}}(x^1, y^1) \right]_{y^1=x^1}, \quad (6.9)$$

$$\langle 0 | \widehat{T}_{01}(x^1) | 0 \rangle_{ren} = \langle 0 | \widehat{T}_{10}(x^1) | 0 \rangle_{ren} = 0, \quad (6.10)$$

$$\begin{aligned} & \langle 0 | \widehat{T}_{11}(x^1) | 0 \rangle_{ren} = \\ & = \left[\left(\frac{1}{4} - \xi \right) D_{-\frac{1}{2}}(x^1, y^1) + \frac{1}{4} \partial_{x^1 y^1} D_{\frac{1}{2}}(x^1, y^1) - \xi \partial_{x^1 x^1} D_{\frac{1}{2}}(x^1, y^1) \right]_{y^1=x^1}. \end{aligned} \quad (6.11)$$

In the following, the scheme outlined above will be illustrated in detail, as an example, for the case of Dirichlet boundary conditions. For the other boundary conditions

²⁹Notice that the analogue of Eq. (6.3) in terms of $\tilde{T}$ is not so simple and straightforward to employ, due to the logarithmic behaviour of the modified cylinder kernel near $\mathfrak{t} = 0$; on the other hand, the analogue of (6.4) in terms of T has a singularity in the gamma function for $u = 0$. Thus, for the computations in which we are interested, there is no better strategy than using Eq. (6.3) with T and Eq. (6.4) with $\tilde{T}$; this also explains why, in the sequel, we will frequently refer to both kernels.

we will be more synthetic but, in any case, we will always report the expressions for $T(\mathbf{t}; \mathbf{x}, \mathbf{y})$, $\tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y})$ and $\langle 0 | \hat{T}_{\mu\nu} | 0 \rangle_{ren}$; in particular recall the considerations of subsection 2.5 and note that in this case Eq. (2.24) gives

$$\xi_1 = 0 . \quad (6.12)$$

6.4 The total energy. Since no singularity appears, we can use the general prescription (4.11); the latter, along with Eq. (6.8), allow us to derive an explicit expression for the bulk energy, for any one of the several boundary conditions to be considered in the following. More precisely, we have

$$E^{ren} = -\frac{1}{2} \text{Res} \left(\mathbf{t}^{-2} T(\mathbf{t}) ; 0 \right) \quad (6.13)$$

where $T(\mathbf{t})$ is the cylinder trace of Eq. (3.57).

In passing, let us remark that the renormalized boundary energy B^{ren} always vanishes identically in the cases considered, due to the prescribed boundary conditions (indeed, the same statement can be made for the regularized version B^u ; see Eq. (4.9)).

We also mention the following fact: by direct comparison of the results reported in subsections 6.6-6.9 it appears that the results derived using Eq. (6.13) could as well be deduced integrating over $(0, a)$ the conformal part of the renormalized energy density $\langle 0 | \hat{T}_{00} | 0 \rangle_{ren}$. On the contrary, the non-conformal part of the latter appears to diverge in a non-integrable manner near the end-points $x = 0$ and $x = a$.

6.5 The boundary forces. Let us remark that, since the boundary is zero-dimensional, the nominal “pressure” on the boundary points $x^1 = 0$, $x^1 = a$ does in fact coincide with the force on these points; because of this, we adopt the notation

$$F_{ren}(x^1) \equiv p^{ren}(x^1) \quad \text{for } x^1 = 0, a . \quad (6.14)$$

For all the (non periodic) boundary conditions to be analysed in the following subsections, there are in principle two definitions of the renormalized boundary forces; these descend from the two alternatives pointed out in the general discussion on pressure of subsection 4.2.

Let us indicate with $n^1(x^1)$ the unit “outer normal” at the points on the boundary, so that $n^1(0) = -1$ and $n^1(a) = 1$. The first definition reads

$$F_{ren}(x^1) := \langle 0 | \hat{T}_{11}^u(x^1) | 0 \rangle \Big|_{u=0} n^1(x^1) \quad (6.15)$$

(see Eq. (4.22) and notice that the prescription of taking the regular part is superfluous, since no singularity arises; namely, we first compute the regularized stress-energy

tensor at the boundary point x^1 , and then we analytically continue at $u = 0$). The second alternative is to define (see Eq. (4.23))

$$F_{ren}(x^1) := \left(\lim_{x'^1 \in (0, a), x'^1 \rightarrow x^1} \langle 0 | \widehat{T}_{11}(x'^1) | 0 \rangle_{ren} \right) n^1(x^1) \quad (6.16)$$

(i.e., we first renormalize at inner points of the interval $(0, a)$, and then move towards the boundary). As a matter of fact, for all boundary conditions considered in the next subsections, the equivalence between (6.15) and (6.16) will be checked by direct computation.

6.6 Dirichlet boundary conditions. As a first example, let us consider the case where the field fulfills Dirichlet conditions at both the end points of the segment $(0, a)$, that is

$$\widehat{\phi}(t, x^1) = 0 \quad \text{for } t \in \mathbf{R}, \text{ and } x^1 = 0 \text{ or } x^1 = a. \quad (6.17)$$

A complete orthonormal set of eigenfunctions $(F_k)_{k \in \mathcal{K}}$ for $\mathcal{A}$ and the related eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$ are

$$F_k(x^1) := \sqrt{\frac{2}{a}} \sin(k x^1), \quad \omega_k^2 := k^2 \quad \text{for } k \in \mathcal{K} \equiv \left\{ \frac{n\pi}{a} \mid n = 1, 2, 3, \dots \right\}. \quad (6.18)$$

The expansion (6.2) for the cylinder kernel associated to $\mathcal{A}$ reads

$$T(\mathbf{t}; x^1, y^1) = \frac{2}{a} \sum_{n=1}^{+\infty} e^{-\frac{n\pi}{a} \mathbf{t}} \sin\left(\frac{n\pi}{a} x^1\right) \sin\left(\frac{n\pi}{a} y^1\right); \quad (6.19)$$

re-writing the trigonometric functions in terms of complex exponentials, the right-hand side of the above equation reduces to a sum of four geometric series, which can be explicitly evaluated. The final result is

$$T(\mathbf{t}; x^1, y^1) = \frac{1}{2a} \left[\frac{\cos(\frac{\pi}{a}(x^1 - y^1)) - e^{-\frac{\pi}{a} \mathbf{t}}}{\cosh(\frac{\pi}{a} \mathbf{t}) - \cos(\frac{\pi}{a}(x^1 - y^1))} - \frac{\cos(\frac{\pi}{a}(x^1 + y^1)) - e^{-\frac{\pi}{a} \mathbf{t}}}{\cosh(\frac{\pi}{a} \mathbf{t}) - \cos(\frac{\pi}{a}(x^1 + y^1))} \right]; \quad (6.20)$$

the same expression is also reported, e.g., in [47, 50], but therein it is not used to compute the full, renormalized stress-energy VEV.

Expressing the hyperbolic functions in terms of exponentials, we easily obtain the primitive of T which vanishes exponentially for $\mathbf{t} \rightarrow +\infty$, that is $-\tilde{T}$; in conclusion

$$\begin{aligned} \tilde{T}(\mathbf{t}; x^1, y^1) = & -\frac{1}{2\pi} \left[\ln \left(1 - 2e^{-\frac{\pi}{a} \mathbf{t}} \cos \left(\frac{\pi}{a}(x^1 - y^1) \right) + e^{-\frac{2\pi}{a} \mathbf{t}} \right) + \right. \\ & \left. - \ln \left(1 - 2e^{-\frac{\pi}{a} \mathbf{t}} \cos \left(\frac{\pi}{a}(x^1 + y^1) \right) + e^{-\frac{2\pi}{a} \mathbf{t}} \right) \right]. \end{aligned} \quad (6.21)$$

Both T and the space derivatives of $\tilde{T}$ have meromorphic extensions in $\mathfrak{t}$ to a complex neighborhood of $[0, +\infty)$, with poles only at $\mathfrak{t} = 0$; so, we can employ Eq.s (6.5) (6.6) to obtain from them the renormalized Dirichlet kernel and its spatial derivatives. For example, since

$$\left. T(\mathfrak{t}; x^1, y^1) \right|_{y^1=x^1} = \frac{1}{\pi \mathfrak{t}} - \frac{\pi(3 - \sin^2(\frac{\pi}{a} x^1))}{12a^2 \sin^2(\frac{\pi}{a} x^1)} \mathfrak{t} + \frac{\pi^3(15(2 + \cos(\frac{2\pi}{a} x^1)) - \sin^4(\frac{\pi}{a} x^1))}{720a^4 \sin^4(\frac{\pi}{a} x^1)} \mathfrak{t}^3 + O(\mathfrak{t}^5), \quad (6.22)$$

for $\mathfrak{t} \rightarrow 0$, evaluating explicitly the residue in Eq. (6.5), it follows

$$\left. D_{-\frac{1}{2}}(x^1, y^1) \right|_{y^1=x^1} = \frac{\pi}{12a^2} \frac{3 - \sin^2(\frac{\pi}{a} x^1)}{\sin^2(\frac{\pi}{a} x^1)}. \quad (6.23)$$

Proceeding similarly for the derivatives of the Dirichlet kernel, and then using Eq.s (6.9-6.11), one obtains the following expression for the renormalized VEV of the stress-energy tensor:

$$\begin{aligned} \langle 0 | \hat{T}_{\mu\nu}(x^1) | 0 \rangle_{ren} \Big|_{\mu, \nu=0,1} &= A \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} + \xi B(x^1) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \\ A &:= \frac{\pi}{24a^2}, \quad B(x^1) := \frac{\pi}{2a^2} \frac{1}{\sin^2(\frac{\pi}{a} x^1)} \quad \text{for } x^1 \in (0, a). \end{aligned} \quad (6.24)$$

Let us now discuss the renormalized bulk energy. To this purpose, we first note that the expansion (3.58) for the cylinder trace gives

$$T(\mathfrak{t}) = \sum_{n=1}^{+\infty} e^{-\frac{n\pi}{a} \mathfrak{t}} = \frac{1}{e^{\frac{\pi}{a} \mathfrak{t}} - 1}; \quad (6.25)$$

then, using prescription (6.13), we readily infer

$$E^{ren} = -\frac{\pi}{24a}. \quad (6.26)$$

In conclusion, let us consider the boundary forces; it is easily seen that both definitions (6.15) and (6.16) give (with A as in Eq. (6.24))

$$F_{ren}(0) = A, \quad F_{ren}(a) = -A. \quad (6.27)$$

6.7 Dirichlet-Neumann boundary conditions. Let us now consider the case where Dirichlet and Neumann boundary conditions are respectively prescribed at the two end points of the segment $(0, a)$: we assume

$$\hat{\phi}(t, 0) = 0, \quad \partial_{x^1} \hat{\phi}(t, a) = 0 \quad \text{for } t \in \mathbf{R}. \quad (6.28)$$

In this case, a complete orthonormal set of eigenfunctions $(F_k)_{k \in \mathcal{K}}$ for $\mathcal{A}$ and the related eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$ are described by

$$F_k(x^1) := \sqrt{\frac{2}{a}} \sin(k x^1), \quad \omega_k^2 := k^2 \quad \text{for } k \in \mathcal{K} \equiv \left\{ \left(n + \frac{1}{2} \right) \frac{\pi}{a} \mid n = 0, 1, 2, \dots \right\}. \quad (6.29)$$

Using the expansion (6.2), we can determine the cylinder kernel T and then obtain the modified kernel $\tilde{T}$ as minus the primitive of T , vanishing for $\mathfrak{t} \rightarrow +\infty$; the final results are

$$T(\mathfrak{t}; x^1, y^1) = \frac{1}{a} \left[\frac{\sinh(\frac{\pi}{2a} \mathfrak{t}) \cos(\frac{\pi}{2a}(x^1 - y^1))}{\cosh(\frac{\pi}{a} \mathfrak{t}) - \cos(\frac{\pi}{a}(x^1 - y^1))} - \frac{\sinh(\frac{\pi}{2a} \mathfrak{t}) \cos(\frac{\pi}{2a}(x^1 + y^1))}{\cosh(\frac{\pi}{a} \mathfrak{t}) - \cos(\frac{\pi}{a}(x^1 + y^1))} \right], \quad (6.30)$$

$$\begin{aligned} \tilde{T}(\mathfrak{t}; x^1, y^1) = \\ \frac{1}{2\pi} \left[\ln \left(\frac{\cos(\frac{\pi}{2a}(x^1 - y^1)) + \cosh(\frac{\pi}{2a} \mathfrak{t})}{\cos(\frac{\pi}{2a}(x^1 - y^1)) - \cosh(\frac{\pi}{2a} \mathfrak{t})} \right) - \ln \left(\frac{\cos(\frac{\pi}{2a}(x^1 + y^1)) + \cosh(\frac{\pi}{2a} \mathfrak{t})}{\cos(\frac{\pi}{2a}(x^1 + y^1)) - \cosh(\frac{\pi}{2a} \mathfrak{t})} \right) \right]. \end{aligned} \quad (6.31)$$

Using the above expressions along with Eq.s (6.5), (6.6) and (6.9-6.11), one obtains the renormalized VEV of the stress-energy tensor:

$$\begin{aligned} \langle 0 | \hat{T}_{\mu\nu}(x^1) | 0 \rangle_{ren} \Big|_{\mu, \nu=0,1} &= A \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \xi B(x^1) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \\ A &:= \frac{\pi}{48a^2}, \quad B(x^1) := \frac{\pi}{2a^2} \frac{\cos(\frac{\pi}{a} x^1)}{\sin^2(\frac{\pi}{a} x^1)} \quad \text{for } x^1 \in (0, a). \end{aligned} \quad (6.32)$$

Next, we derive the cylinder trace using again the expansion (3.58):

$$T(\mathfrak{t}) = \sum_{n=0}^{+\infty} e^{-(n+\frac{1}{2})\frac{\pi}{a}\mathfrak{t}} = \frac{e^{\frac{\pi}{2a}\mathfrak{t}}}{e^{\frac{\pi}{a}\mathfrak{t}} - 1}. \quad (6.33)$$

Now prescription (6.13) allows us to obtain the renormalized total bulk energy:

$$E^{ren} = \frac{\pi}{48a}. \quad (6.34)$$

Concerning the boundary forces, also in this case definitions (6.15) (6.16) agree and give

$$F_{ren}(0) = -A, \quad F_{ren}(a) = A \quad (6.35)$$

where A is as in Eq. (6.32); notice, in particular, that the above expressions have the opposite sign with respect to the ones of Eq. (6.27), corresponding the case of Dirichlet boundary conditions.

6.8 Neumann boundary conditions. We are now going to study the case where

$$\partial_{x^1} \widehat{\phi}(t, x^1) = 0 \quad \text{for } t \in \mathbf{R}, \text{ and } x^1 = 0 \text{ or } x^1 = \pi_a . \quad (6.36)$$

In this case, according to the considerations of subsection 5.1, the Hilbert space $L^2(0, a)$ has to be replaced with the space $L_0^2(0, a)$ of square integrable functions on $(0, a)$ with mean zero (see Eq. (5.2)); in this space, a complete orthonormal set of eigenfunctions $(F_k)_{k \in \mathcal{K}}$ for $\mathcal{A} = -\partial_{x^1 x^1}$ and the corresponding eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$ are given by

$$F_k(x^1) := \sqrt{\frac{2}{a}} \cos(k x^1), \quad \omega_k^2 := k^2 \quad \text{for } k \in \mathcal{K} \equiv \left\{ \frac{n\pi}{a} \mid n = 1, 2, 3, \dots \right\}. \quad (6.37)$$

The cylinder kernel associated to $\mathcal{A}$ can be evaluated according to Eq. (6.2) to obtain

$$T(\mathbf{t}; x^1, y^1) = \frac{1}{2a} \left[\frac{\cos(\frac{\pi}{a}(x^1 - y^1)) - e^{-\mathbf{t}}}{\cosh \mathbf{t} - \cos(\frac{\pi}{a}(x^1 - y^1))} + \frac{\cos(\frac{\pi}{a}(x^1 + y^1)) - e^{-\mathbf{t}}}{\cosh \mathbf{t} - \cos(\frac{\pi}{a}(x^1 + y^1))} \right]; \quad (6.38)$$

while for the modified cylinder kernel, computed as minus the primitive of T , we obtain

$$\begin{aligned} \tilde{T}(\mathbf{t}; x^1, y^1) = & -\frac{1}{2\pi} \left[\ln \left(1 - 2e^{-\frac{\pi}{a}\mathbf{t}} \cos \left(\frac{\pi}{a}(x^1 - y^1) \right) + e^{-\frac{2\pi}{a}\mathbf{t}} \right) + \right. \\ & \left. + \ln \left(1 - 2e^{-\frac{\pi}{a}\mathbf{t}} \cos \left(\frac{\pi}{a}(x^1 + y^1) \right) + e^{-\frac{2\pi}{a}\mathbf{t}} \right) \right]. \end{aligned} \quad (6.39)$$

Resorting once more to Eq.s (6.5), (6.6) and (6.9-6.11), the renormalized VEV of the stress-energy tensor is found to be

$$\langle 0 | \widehat{T}_{\mu\nu}(x^1) | 0 \rangle_{ren} \Big|_{\mu, \nu=0,1} = A \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} - \xi B(x^1) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad (6.40)$$

where A and $B(x^1)$ are defined as in Eq. (6.24).

In the case under analysis the spectrum of $\mathcal{A}$ coincides with the one obtained for Dirichlet boundary conditions (compare Eq.s (6.18) (6.37)); therefore, the cylinder trace is again given by Eq. (6.25), and we derive the same renormalized bulk energy as in Eq. (6.26):

$$E^{ren} = -\frac{\pi}{24a}.$$

Finally, both definitions (6.15) and (6.16) give for the boundary forces the same results as in the case of Dirichlet boundary conditions (see Eq. (6.27)):

$$F_{ren}(0) = A, \quad F_{ren}(a) = -A \quad (6.41)$$

(again, A is as in Eq. (6.24)).

6.9 Periodic boundary conditions. The last case we consider for the segment configuration is the one where the field satisfies periodic boundary conditions:

$$\widehat{\phi}(t, 0) = \widehat{\phi}(t, a) , \quad \partial_{x^1} \widehat{\phi}(t, 0) = \partial_{x^1} \widehat{\phi}(t, a) \quad \text{for } t \in \mathbf{R} . \quad (6.42)$$

As explained in subsection 5.3, this case would be more properly formulated in terms of a free scalar field on the 1-dimensional torus $\mathbf{T}_a^1 := \mathbf{R}/(a\mathbf{Z})$. Besides, similarly to the case of Neumann boundary conditions, recall that the basic Hilbert space is $L_0^2(\mathbf{T}_a^1) = \{f \in L^2(\mathbf{T}_a^1) \mid \int_0^a dx^1 f(x^1) = 0\}$ (see subsection 5.1 and the footnote 26 of page 52). In this space a complete orthonormal set of eigenfunctions $(F_k)_{k \in \mathcal{K}}$ for $\mathcal{A}$, with the corresponding eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$, is

$$F_k(x^1) := \sqrt{\frac{1}{a}} e^{ikx^1}, \quad \omega_k^2 := k^2 \quad \text{for } k \in \mathcal{K} \equiv \left\{ \pm \frac{2n\pi}{a} \mid n = 1, 2, 3, \dots \right\} . \quad (6.43)$$

Let us pass to determine the cylinder and modified cylinder kernel associated to $\mathcal{A}$; using the same methods of the previous subsections, we obtain

$$T(\mathbf{t}; x^1, y^1) = \frac{\cos(\frac{2\pi}{a}(x^1 - y^1)) - e^{-\frac{2\pi}{a}\mathbf{t}}}{a [\cosh(\frac{2\pi}{a}\mathbf{t}) - \cos(\frac{2\pi}{a}(x^1 - y^1))]} , \quad (6.44)$$

$$\tilde{T}(\mathbf{t}; x^1, y^1) = -\frac{1}{2\pi} \ln \left(1 - 2 e^{-\frac{2\pi}{a}\mathbf{t}} \cos \left(\frac{2\pi}{a}(x^1 - y^1) \right) + e^{-\frac{4\pi}{a}\mathbf{t}} \right) \quad (6.45)$$

(the same expression for T is also reported, e.g., in [47], again for other purposes). Eq.s (6.5), (6.6) and (6.9-6.11), yield the following expression for the renormalized VEV of the stress-energy tensor:

$$\langle 0 | \widehat{T}_{\mu\nu}(x^1) | 0 \rangle_{ren} \Big|_{\mu, \nu=0,1} = \frac{\pi}{6a^2} \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} . \quad (6.46)$$

Let us stress that the above results respects the invariance under translations $x^1 \mapsto x^1 + \alpha$ (for any $\alpha \in \mathbf{R}$) of the given configuration, since it does not depend explicitly on the spatial coordinate x^1 .

To conclude, we discuss the renormalized bulk energy. We first note that expansion (3.58) for the cylinder trace yields, in the present case,

$$T(\mathbf{t}) = \left(\sum_{n=-\infty}^{-1} + \sum_{n=1}^{+\infty} \right) e^{-\frac{2|n|\pi}{a}\mathbf{t}} = 2 \sum_{n=1}^{+\infty} e^{-\frac{2n\pi}{a}\mathbf{t}} = \frac{2}{e^{\frac{2\pi}{a}\mathbf{t}} - 1} ; \quad (6.47)$$

then, using once more prescription (6.13), we obtain for the bulk energy

$$E^{ren} = -\frac{\pi}{6a} . \quad (6.48)$$

Acknowledgments. This work was partly supported by INdAM, INFN and by MIUR, PRIN 2010 Research Project “Geometric and analytic theory of Hamiltonian systems in finite and infinite dimensions”.

A Appendix. On the form (2.11) for the stress-energy tensor

Eq. (2.11) is the quantized version of a classical formula for the stress-energy tensor [12, 17, 19, 79], which we review here for completeness. Following Section 2, we refer to Minkowski spacetime; after an inertial frame has been chosen, the latter is identified with $\mathbf{R}^{d+1} = \mathbf{R} \times \mathbf{R}^d \ni x \equiv (x^\mu) \equiv (t, \mathbf{x})$. We confine the attention to a subset of the form $\mathbf{R} \times \Omega$, where $\Omega \subset \mathbf{R}^d$ is a spatial domain; let us consider a classical scalar field ϕ on $\mathbf{R} \times \Omega$ described by an arbitrary Lagrangian density $\mathcal{L} = \mathcal{L}(\phi, \partial\phi, x)$. The associated canonical stress-energy tensor is

$$T_{\mu\nu}^{can} := -\frac{\partial\mathcal{L}}{\partial(\partial^\mu\phi)} \partial_\nu\phi + \eta_{\mu\nu}\mathcal{L} , \quad (\text{A.1})$$

and fulfills

$$\partial^\mu T_{\mu\nu}^{can} = -\partial_\nu\mathcal{L} \quad (\text{A.2})$$

along the solutions of the field equations. If Σ is any spacelike hypersurface with normal unit vector N^μ and volume element dv , we define the canonical momentum

$$P_\nu^{can}(\Sigma) := \int_\Sigma dv N^\mu T_{\mu\nu}^{can} . \quad (\text{A.3})$$

For simplicity we rescript the attention to the case

$$\Sigma = \{t\} \times \Omega , \quad (\text{A.4})$$

for a fixed $t \in \mathbf{R}$, writing $P_\nu^{can}(t)$ for the corresponding canonical momentum. In this case we can take $(N^\mu) = (1, 0, \dots, 0)$ and $\int_\Sigma dv$ corresponds to integration on Ω with respect to the usual volume element $d\mathbf{x}$, so

$$P_\nu^{can}(t) := \int_\Omega d\mathbf{x} T_{0\nu}^{can}(t, \mathbf{x}) . \quad (\text{A.5})$$

Writing x for $(t, \mathbf{x})$, we have $dP_\nu^{can}/dt(t) = \int_\Omega d\mathbf{x} \partial_0 T_{0\nu}^{can}(x) = \int_\Omega d\mathbf{x} (-\partial^0 T_{0\nu}^{can})(x) = \int_\Omega d\mathbf{x} (\partial^i T_{i\nu}^{can} - \partial^\mu T_{\mu\nu}^{can})(x) = \int_\Omega d\mathbf{x} [\partial^i T_{i\nu}^{can}(x) + (\partial_\nu\mathcal{L})(\phi(x), \partial\phi(x), x)]$, i.e., by the d -dimensional divergence theorem,

$$\frac{dP_\nu^{can}}{dt}(t) = \int_\Omega d\mathbf{x} (\partial_\nu\mathcal{L})(\phi(x), \partial\phi(x), x) + \int_{\partial\Omega} da(\mathbf{x}) n^i(\mathbf{x}) T_{i\nu}^{can}(x) . \quad (\text{A.6})$$

Here (and in the sequel) $\mathbf{n}(\mathbf{x}) = (n^i(\mathbf{x}))$ is the outer unit vector in $\mathbf{R}^d$ normal to the boundary $\partial\Omega$ at $\mathbf{x}$, and da is the $(d-1)$ -dimensional area element ⁽³⁰⁾. For a

³⁰Obviously enough, if Ω is unbounded we intend $\int_{\partial\Omega} da(\mathbf{x}) n^i(\mathbf{x}) := \lim_{\ell \rightarrow +\infty} \int_{\partial\Omega_\ell} da_\ell(\mathbf{x}) n_\ell^i(\mathbf{x})$ where $\Omega_1 \subset \Omega_2 \subset \Omega_3 \subset \dots$ are bounded domains such that $\cup_{\ell=1}^{+\infty} \Omega_\ell = \Omega$.

number of reasons briefly reviewed in the sequel, it is customary (see, e.g., [19, 94]) to consider an “improved stress-energy tensor” of the form

$$T_{\mu\nu} := T_{\mu\nu}^{can} + \partial^\lambda F_{\lambda\mu\nu} , \quad (\text{A.7})$$

where $F_{\lambda\mu\nu}$ is a covariant tensor of rank 3 such that

$$F_{\lambda\mu\nu} = -F_{\mu\lambda\nu} . \quad (\text{A.8})$$

Condition (A.8) implies $\partial^\mu(\partial^\lambda F_{\lambda\mu\nu}) = 0$, thus ensuring

$$\partial^\mu T_{\mu\nu} = \partial^\mu T_{\mu\nu}^{can} . \quad (\text{A.9})$$

We can give definitions similar to (A.3) and (A.5) using the improved stress-energy tensor; in particular, for each $t \in \mathbf{R}$, we define the “improved momentum”

$$P_\nu(t) := \int_{\Omega} d\mathbf{x} T_{0\nu}(t, \mathbf{x}) . \quad (\text{A.10})$$

We claim that

$$P_\nu(t) = P_\nu^{can}(t) + \int_{\partial\Omega} da(\mathbf{x}) n^i(\mathbf{x}) F_{i0\nu}(t, \mathbf{x}) , \quad (\text{A.11})$$

where da and $\mathbf{n}(\mathbf{x})$ have the same meaning as before. To prove this, note that

$$T_{0\nu} - T_{0\nu}^{can} = \partial^\lambda F_{\lambda 0\nu} = \partial^0 F_{00\nu} + \partial^i F_{i0\nu} = \partial^i F_{i0\nu} \quad (\text{A.12})$$

($F_{00\nu} = 0$ due to Eq. (A.8)); thus $P_\nu(t) = P_\nu^{can}(t) + \int_{\Omega} d\mathbf{x} \partial^i F_{i0\nu}(t, \mathbf{x})$, and the d -dimensional divergence theorem yields Eq. (A.11).

In many cases of interest, the boundary term in Eq. (A.11) is zero. In particular, this happens if $\Omega = \mathbf{R}^d$ and $F_{i0\nu}(t, \mathbf{x})$ vanishes rapidly for $\mathbf{x} \rightarrow \infty$. In the case of a bounded domain, the boundary term can be zero if $F_{i0\nu}$ depends suitably on the field ϕ and the latter fulfills appropriate conditions on $\partial\Omega$.

Whether or not the boundary term in Eq. (A.11) vanishes, using Eq. (A.9) we prove that the improved momentum evolves according to the analogue of Eq. (A.6), i.e.,

$$\frac{dP_\nu}{dt}(t) = \int_{\Omega} d\mathbf{x} (\partial_\nu \mathcal{L})(\phi(x), \partial\phi(x), x) + \int_{\partial\Omega} da(\mathbf{x}) n^i(\mathbf{x}) T_{i\nu}(x) . \quad (\text{A.13})$$

The improved stress-energy tensor is symmetric if and only if

$$\partial^\lambda (F_{\lambda\mu\nu} - F_{\lambda\nu\mu}) = - (T_{\mu\nu}^{can} - T_{\nu\mu}^{can}) . \quad (\text{A.14})$$

When $T_{\mu\nu}^{can}$ is not symmetric and it can be found a rank 3 tensor $F_{\lambda\mu\nu}$ fulfilling conditions (A.8) and (A.14), the symmetry of the improved stress-energy tensor (A.7) is itself a good reason to consider this object.

There are reasons to consider the improved stress-energy tensor even in the case when $T_{\mu\nu}^{can}$ is itself symmetric (of course, in this case Eq. (A.14) requires that $\partial^\lambda F_{\lambda\mu\nu}$ be symmetric in μ and ν). One of these reasons has been pointed out by Callan et al. [19]; in few words, after quantization the divergences of the improved tensor can happen to be softer than the divergences of the canonical one, especially in perturbative renormalization.

In this paper we are interested in a field theory governed by the equation $0 = (-\partial_{tt} + \Delta - V)\phi = (\partial_\mu \partial^\mu - V)\phi$ which arises from the Lagrangian

$$\mathcal{L} := -\frac{1}{2} \partial^\mu \phi \partial_\mu \phi - \frac{1}{2} V \phi^2 . \quad (\text{A.15})$$

The corresponding canonical stress-energy tensor is

$$T_{\mu\nu}^{can} = \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} \eta_{\mu\nu} (\partial^\lambda \phi \partial_\lambda \phi + V \phi^2) ; \quad (\text{A.16})$$

this is symmetric and fulfills (along solutions of the field equations)

$$\partial^\mu T_{\mu\nu}^{can} = -\frac{1}{2} (\partial_\nu V) \phi^2 . \quad (\text{A.17})$$

Condition (A.8) is satisfied by the tensor

$$F_{\lambda\mu\nu} := -\xi (\eta_{\lambda\nu} \partial_\mu - \eta_{\mu\nu} \partial_\lambda) \phi^2 , \quad (\text{A.18})$$

where ξ is a real parameter. In this case

$$\partial^\lambda F_{\lambda\mu\nu} = -\xi (\partial_{\mu\nu} - \eta_{\mu\nu} \partial^\lambda \partial_\lambda) \phi^2 ; \quad (\text{A.19})$$

this tensor is symmetric in μ and ν , so $T_{\mu\nu}$ is symmetric as well. The derivatives of ϕ^2 in Eq. (A.19) can be re-expressed using the field equation $\partial_\mu \partial^\mu \phi = V \phi$, yielding

$$\partial^\lambda F_{\lambda\mu\nu} = -2\xi \partial_\mu \phi \partial_\nu \phi + 2\xi \eta_{\mu\nu} (\partial^\lambda \phi \partial_\lambda \phi + V \phi^2) - 2\xi \phi \partial_{\mu\nu} \phi . \quad (\text{A.20})$$

Thus the improved stress-energy tensor (A.7) takes the form

$$T_{\mu\nu} = (1-2\xi) \partial_\mu \phi \partial_\nu \phi - \left(\frac{1}{2} - 2\xi \right) \eta_{\mu\nu} (\partial^\lambda \phi \partial_\lambda \phi + V \phi^2) - 2\xi \phi \partial_{\mu\nu} \phi , \quad (\text{A.21})$$

of which (2.11) is a natural quantization.

Let us recall that the momentum P_ν corresponding to the improved tensor $T_{\mu\nu}$ is related to the canonical one via Eq. (A.11); in the present framework where $F_{\lambda\mu\nu}$ is given by Eq. (A.18), the boundary term in Eq. (A.11) vanishes under Dirichlet boundary conditions ($\phi(t, \mathbf{x}) = 0$ for all $\mathbf{x} \in \partial\Omega$); this term vanishes as well if $\Omega = \mathbf{R}^d$ and $\phi(t, \mathbf{x}), \partial_\lambda \phi(t, \mathbf{x})$ vanish rapidly for $\mathbf{x} \rightarrow \infty$.

In the case $V = 0$, the above improved tensor (A.21) allows another interpretation: this is the functional derivative with respect to the metric of the action describing a scalar field that interacts with a gravitational field via the scalar curvature, in the limit where ϕ is small and the metric is the Minkowski metric $\eta_{\mu\nu}$ plus a perturbation of the second order in ϕ . Concerning this, see [19, 79, 81] and [36, 37].

Again for $V = 0$, the action of the field coupled to gravity is conformally invariant for $\xi = (d - 1)/(4d)$ [101].

B Appendix. Smoothness properties of some integral kernels

We refer to an operator $\mathcal{A}$ in $L^2(\Omega)$ with features described by Eq.s (3.7), (3.8) or (3.9) (recall that the situation (3.7) is more general than (3.8), and (3.8) is more general than (3.9)). In the present appendix we present a number of results which are useful in relation to several integral kernels associated to $\mathcal{A}$. Proving these results would require a heavy use of functional analysis, which is not among the purposes of the present work; therefore, hereafter we just sketch the basic ideas that will be presented with more details elsewhere [41].

General results for case (3.7). As in the cited equation we assume $\mathcal{A}$ to be any strictly positive, selfadjoint operator in $L^2(\Omega)$. A basic step towards our goal consists in associating a scale of Hilbert spaces $\mathcal{H}^r$ to the powers $\mathcal{A}^r$ for $r \in \mathbf{R}$ (see [6] for a similar construction); speaking somehow loosely, we can describe $\mathcal{H}^r$ as the Hilbert space of generalized functions $f : \Omega \rightarrow \mathbf{C}$ such that $\mathcal{A}^r f \in L^2(\Omega)$, equipped with the inner product

$$\langle f | g \rangle_r := \langle \mathcal{A}^{r/2} f | \mathcal{A}^{r/2} g \rangle \quad (\text{B.1})$$

(as usual, $\langle | \rangle$ denotes the inner product of $L^2(\Omega)$) which induces the norm

$$\|f\|_r := \sqrt{\langle f | f \rangle_r} = \|\mathcal{A}^{r/2} f\| \quad (\text{B.2})$$

⁽³¹⁾. Let $(F_k)_{k \in \mathcal{K}}$ be any complete orthonormal set of (generalized) eigenfunctions

³¹Let us sketch the precise definition of $\mathcal{H}^r$, given in [41]. For any real r we consider in $L^2(\Omega)$ the dense linear subspace D^r on which $\mathcal{A}^{r/2}$ is well defined according to the general framework for functional calculus of selfadjoint operators in Hilbert spaces (see, e.g., [84]); if $(F_k)_{k \in \mathcal{K}}$ is a (generalized) complete orthonormal set of eigenfunctions of $\mathcal{A}$ with eigenvalues ω_k^2 ($\omega_k > \varepsilon > 0$), then D^r is formed by the functions $f \in L^2(\Omega)$ such that $\int_{\mathcal{K}} dk \omega_k^{2r} |\langle F_k | f \rangle|^2 < +\infty$. We introduce on D^r an inner product $\langle | \rangle_r$ following Eq. (B.1), and define $\mathcal{H}^r$ to be the completion of D^r with respect to $\langle | \rangle_r$; for $r_2 \geq r_1$ one finds $D^{r_2} \subset D^{r_1}$, a fact implying $\mathcal{H}^{r_2} \subset \mathcal{H}^{r_1}$. Of course $\mathcal{A}^0$ coincides with the identity operator $\mathbf{1}$ of $L^2(\Omega)$; thus $D^0 = \mathcal{H}^0 = L^2(\Omega)$ and $\langle | \rangle_0$ is the usual L^2 inner product $\langle | \rangle$. For any $r \geq 0$, D^r is itself complete, so $\mathcal{H}^r = D^r \subset L^2(\Omega)$; for $r < 0$ the completion $\mathcal{H}^r$ is larger than D^r , and even of $L^2(\Omega) = \mathcal{H}^0$. If r is sufficiently large, $\mathcal{H}^{-r}$ contains elements which cannot even be interpreted as ordinary functions on Ω ; for example, as shown later in this appendix, the Dirac delta $\delta_{\mathbf{x}}$ at any point $\mathbf{x} \in \Omega$ can be interpreted as an element of $\mathcal{H}^{-r}$ for all $r > d/2$.

of $\mathcal{A}$ with eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$ ($\omega_k \geq \varepsilon > 0$ for all $k \in \mathcal{K}$); then $\mathcal{H}^r$ is made of the generalized function f on Ω such that

$$\int_K dk \, \omega_k^{2r} |\langle F_k | f \rangle|^2 < +\infty ; \quad (\text{B.3})$$

moreover, for any pair of functions $f, g \in \mathcal{H}^r$, there holds

$$\langle f | g \rangle_r = \int_K dk \, \omega_k^{2r} \overline{\langle F_k | f \rangle} \langle F_k | g \rangle . \quad (\text{B.4})$$

Of course, $\mathcal{H}^0$ is the usual space $L^2(\Omega)$, and it can be proved that $\mathcal{H}^{r_2} \hookrightarrow \mathcal{H}^{r_1}$ for $r_2 \geq r_1$ ⁽³²⁾. Moreover, from the characterization (B.3) of $\mathcal{H}^r$ and from the identity $\mathcal{A}^s F_k = \omega_k^{2s} F_k$ it follows that

$$\mathcal{A}^{-s} \text{ maps continuously } \mathcal{H}^r \text{ into } \mathcal{H}^{r'} \text{ for } r, r' \in \mathbf{R} \text{ such that } r' - r < 2\Re s ; \quad (\text{B.5})$$

similarly, one proves that

$$e^{-t\mathcal{A}}, e^{-t\sqrt{\mathcal{A}}} \text{ map continuously } \mathcal{H}^r \text{ into } \mathcal{H}^{r'} \text{ for all } r, r' \in \mathbf{R} . \quad (\text{B.6})$$

Next, consider the inner product of $L^2(\Omega)$; for all $r \in \mathbf{R}$ this can be extended to a continuous, sesquilinear map

$$\langle | \rangle : \mathcal{H}^{-r} \times \mathcal{H}^r \rightarrow \mathbf{C} , \quad (\text{B.7})$$

which allows, in particular, to infer the isomorphic identification $(\mathcal{H}^r)' = \mathcal{H}^{-r}$.

Results for case (3.8). In accordance with the cited equation, we now assume $\mathcal{A} = -\Delta + V$ on an open subset Ω of $\mathbf{R}^d$ with given boundary conditions, $V : \Omega \rightarrow \mathbf{R}$ a C^∞ potential; all these ingredients are chosen so as to ensure the strict positivity of $\mathcal{A}$. In this case it can be proved that

$$\mathcal{H}^r \hookrightarrow C^j(\Omega) \quad \text{for } j \in \mathbf{N}, r \in \mathbf{R} \text{ with } r > \frac{d}{2} + j , \quad (\text{B.8})$$

where $C^j(\Omega)$ carries the topology of uniform convergence of the derivatives up to order j on compact subsets of Ω ; this is induced by the family of seminorms

$$p_{jK}(f) := \max_{|\alpha| \leq j, \mathbf{x} \in K} |\partial^\alpha f(\mathbf{x})| \quad (K \subset \Omega \text{ compact}) \quad (\text{B.9})$$

³²Given any pair of topological vector spaces $\mathcal{X}, \mathcal{Y}$, we say that $\mathcal{X}$ is continuously embedded in $\mathcal{Y}$, and we write $\mathcal{X} \hookrightarrow \mathcal{Y}$, if $\mathcal{X}$ is a linear subspace of $\mathcal{Y}$ and the identity map from $\mathcal{X}$ into $\mathcal{Y}$ is continuous. So, for example, $\mathcal{H}^{r_4} \hookrightarrow \mathcal{H}^{r_3} \hookrightarrow L^2(\Omega) \hookrightarrow \mathcal{H}^{r_2} \hookrightarrow \mathcal{H}^{r_1}$ if $r_4 \geq r_3 \geq 0 \geq r_2 \geq r_1$.

⁽³³⁾. By duality, Eq. (B.8) implies (under the same assumptions on j and r)

$$(C^j(\Omega))' \hookrightarrow (\mathcal{H}^r)' = \mathcal{H}^{-r} . \quad (\text{B.10})$$

Now, let $j \in \mathbf{N}$ and $\mathbf{x} \in \Omega$; the prescription

$$\langle \delta_{\mathbf{x}}, f \rangle := f(\mathbf{x}) \quad (\text{B.11})$$

makes sense for all $f \in C^j(\Omega)$ and allows to interpret the Dirac delta function $\delta_{\mathbf{x}}$ as an element of the dual space $(C^j(\Omega))'$. Moreover, it can be shown that the mapping $\delta : \mathbf{x} \in \Omega \mapsto \delta_{\mathbf{x}} \in (C^j(\Omega))'$ is itself of class C^j ⁽³⁴⁾; of course, this fact and Eq.s (B.10) imply that

$$\delta : \Omega \rightarrow \mathcal{H}^{-r} , \quad \mathbf{x} \mapsto \delta_{\mathbf{x}} \quad \text{is } C^j \text{ if } r \in \mathbf{R}, j \in \mathbf{N} \text{ and } r > \frac{d}{2} + j . \quad (\text{B.12})$$

Finally, let $j \in \mathbf{N}$ and suppose that

$$\begin{aligned} \mathcal{B} : \mathcal{H}^{-(d/2+j_2+\eta)} &\rightarrow \mathcal{H}^{d/2+j_1+\eta} \text{ is linear and continuous} \\ &\text{for some } \eta > 0 \text{ and all } j_1, j_2 \in \mathbf{N} \text{ with } j_1 + j_2 \leq j ; \end{aligned} \quad (\text{B.13})$$

we claim that, in this case,

$$\begin{aligned} &\text{the kernel } \Omega \times \Omega \rightarrow \mathbf{C}, (\mathbf{x}, \mathbf{y}) \mapsto \mathcal{B}(\mathbf{x}, \mathbf{y}) := \langle \delta_{\mathbf{x}} | \mathcal{B} \delta_{\mathbf{y}} \rangle \\ &\text{is (well defined and) of class } C^j(\Omega \times \Omega) . \end{aligned} \quad (\text{B.14})$$

In fact, each derivative $\partial \mathcal{B}(\ , \)$ of order $\leq j$ involves (in an arbitrary order) α_i operations of derivation with respect to x^i and β_i operations of derivation with respect to y^i ($i = 1, \dots, d$) where $j_1 := \alpha_1 + \dots + \alpha_d$ and $j_2 := \beta_1 + \dots + \beta_d$ are such

³³Let us sketch the derivation of Eq. (B.8). Due to well-known results on elliptic operators [23], one has

$$\mathcal{H}^r \hookrightarrow H_{loc}^r(\Omega) \quad \text{for all } r \geq 0 ,$$

where $H_{loc}^r(\Omega)$ is the standard local Sobolev space of order r (that is, the space of functions $f : \Omega \rightarrow \mathbf{C}$ such that $(1 - \Delta)^{r/2}(\varphi f) \in L^2(\Omega)$ for all smooth, compactly supported functions $\varphi : \Omega \rightarrow \mathbf{C}$). On the other hand, the usual Sobolev imbedding theorems [5] give

$$H_{loc}^r(\Omega) \hookrightarrow C^j(\Omega) \quad \text{for all } r \in \mathbf{R}, j \in \mathbf{N} \text{ with } r > \frac{d}{2} + j .$$

Summing up, the embeddings discussed in this footnote yield the thesis (B.8).

³⁴For example, setting $j = 1$, one finds that the map $\mathbf{x} \mapsto \delta_{\mathbf{x}} \in (C^1(\Omega))'$ is C^1 with derivatives $(\partial_i \delta)_{\mathbf{x}}$ such that

$$\langle (\partial_i \delta)_{\mathbf{x}}, f \rangle = \partial_i f(\mathbf{x}) \quad (i \in \{1, \dots, d\})$$

for all $f \in C^1(\Omega)$.

that $j_1 + j_2 \leq j$; any such derivative exists and is continuous on $\Omega \times \Omega$, with the explicit expression $\partial \mathcal{B}(\mathbf{x}, \mathbf{y}) = \langle (\partial^\alpha \delta)_\mathbf{x} | \mathcal{B}(\partial^\beta \delta)_\mathbf{y} \rangle$ ⁽³⁵⁾.

Applications to the Dirichlet, heat and cylinder kernels: regularity results. Let $s \in \mathbf{C}$, $j \in \mathbf{N}$; if $\Re s > d/2 + j/2$, using Eq. (B.5) one easily infers that the operator $\mathcal{B} := \mathcal{A}^{-s}$ fulfills the condition (B.13) (with $\eta = \Re s - d/2 - j/2$); thus (B.14) holds. In conclusion, we have the following result for the Dirichlet kernel $D_s(\mathbf{x}, \mathbf{y}) := \langle \delta_\mathbf{x} | \mathcal{A}^{-s} \delta_\mathbf{y} \rangle$:

$$D_s \in C^j(\Omega \times \Omega) \quad \text{for } s \in \mathbf{C}, j \in \mathbf{N} \text{ and } \Re s > \frac{d}{2} + \frac{j}{2}. \quad (\text{B.15})$$

Similarly, for any $\mathfrak{t} > 0$ and any $j \in \mathbf{N}$, due to (B.6) the operators $\mathcal{B} = e^{-\mathfrak{t}\mathcal{A}}$ or $\mathcal{B} = e^{-\mathfrak{t}\sqrt{\mathcal{A}}}$ fulfill the condition (B.13), implying Eq (B.14). So, the heat and cylinder kernels $K(\mathfrak{t}; \mathbf{x}, \mathbf{y}) := \langle \delta_\mathbf{x} | e^{-\mathfrak{t}\mathcal{A}} \delta_\mathbf{y} \rangle$, $T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) := \langle \delta_\mathbf{x} | e^{-\mathfrak{t}\sqrt{\mathcal{A}}} \delta_\mathbf{y} \rangle$ are of class C^j in $\mathbf{x}, \mathbf{y}$, for all $j \in \mathbf{N}$; in conclusion, these kernels are C^∞ in $\mathbf{x}, \mathbf{y}$:

$$K(\mathfrak{t}; \cdot, \cdot), T(\mathfrak{t}; \cdot, \cdot) \in C^\infty(\Omega \times \Omega) \quad \text{for each } \mathfrak{t} > 0. \quad (\text{B.16})$$

Results for case (3.9). We now consider the case described by the cited equation. Thus $\mathcal{A} = -\Delta + V$ on bounded domain Ω with C^∞ boundary $\partial\Omega$, on which Dirichlet conditions are imposed; the potential V is in $C^\infty(\overline{\Omega})$ and $V(\mathbf{x}) \geq 0$ for all $\mathbf{x} \in \overline{\Omega}$ (recall that, due to these assumptions, $\mathcal{A}$ is selfadjoint and strictly positive).

Smoothness up to the boundary. In the setting described above, one can strengthen the results discussed in the previous paragraph of this appendix replacing systematically the space $C^j(\Omega)$ with $C^j(\overline{\Omega})$; this space consists of the functions which are continuous with their derivatives up to order j on the closure $\overline{\Omega} = \Omega \cup \partial\Omega$, and it is equipped with the norm

$$\|f\|_{C^j} := \max_{|\alpha| \leq j, \mathbf{x} \in \overline{\Omega}} |\partial^\alpha f(\mathbf{x})|. \quad (\text{B.17})$$

Similarly, the space $C^j(\Omega \times \Omega)$ can be replaced with $C^j(\overline{\Omega} \times \overline{\Omega})$, endowed with the norm

$$\|f\|_{C^j} := \max_{|\alpha|+|\beta| \leq j, \mathbf{x}, \mathbf{y} \in \overline{\Omega}} |\partial_\mathbf{x}^\alpha \partial_\mathbf{y}^\beta f(\mathbf{x}, \mathbf{y})|. \quad (\text{B.18})$$

In the present case Eq. (B.8) has the stronger version

$$\mathcal{H}^r \hookrightarrow C^j(\overline{\Omega}) \quad \text{for } j \in \mathbf{N}, r \in \mathbf{R} \text{ with } r > \frac{d}{2} + j; \quad (\text{B.19})$$

³⁵note that, due to (B.12), the map $\mathbf{y} \mapsto (\partial^\beta \delta)_\mathbf{y}$ is continuous from Ω to $\mathcal{H}^{-(d/2+j_2+\eta)}$; due to (B.13), $\mathbf{y} \mapsto \mathcal{B}(\partial^\beta \delta)_\mathbf{y}$ is continuous from Ω to $\mathcal{H}^{d/2+j_1+\eta}$; finally, due to (B.12) the map $\mathbf{x} \mapsto (\partial^\alpha \delta)_\mathbf{x}$ is continuous from Ω to $\mathcal{H}^{-(d/2+j_2+\eta)}$, and the sesquilinear form $\langle \cdot | \cdot \rangle$ is continuous on $\mathcal{H}^{-(d/2+j_1+\eta)} \times \mathcal{H}^{d/2+j_1+\eta}$.

similarly, Eq. (B.14) (with the assumptions (B.13) on $\mathcal{B}$) holds with $C^j(\Omega \times \Omega)$ replaced by $C^j(\overline{\Omega} \times \overline{\Omega})$ ⁽³⁶⁾; due to these facts, Eq.s (B.15) and (B.16) for the Dirichlet, heat and cylinder kernels hold in the stronger versions

$$D_s \in C^j(\overline{\Omega} \times \overline{\Omega}) \quad \text{for } s \in \mathbf{C}, j \in \mathbf{N} \text{ with } \Re s > \frac{d}{2} + \frac{j}{2}; \quad (\text{B.20})$$

$$K(\mathbf{t}; \cdot, \cdot), T(\mathbf{t}; \cdot, \cdot) \in C^\infty(\overline{\Omega} \times \overline{\Omega}) \quad \text{for each } \mathbf{t} > 0. \quad (\text{B.21})$$

Estimates for the eigenfunctions and eigenvalues of $\mathcal{A}$. Due to the boundedness of Ω , $\mathcal{A}$ has purely point spectrum; in fact, one can build for this operator a complete orthonormal set of proper eigenfunctions $(F_k)_{k \in \mathcal{K}}$ with eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$, where $\mathcal{K} = \{1, 2, 3, \dots\}$ and the labels are chosen so that $0 < \omega_1 \leq \omega_2 \leq \omega_3 \leq \dots$ (with the possibility that some of these inequalities are equalities, to deal with the case of degenerate eigenvalues).

Let us discuss the smoothness properties of the eigenfunctions and derive some norm bounds for them. Clearly, for each $k \in \{1, 2, 3, \dots\}$ and $r \in \mathbf{R}$, we have $F_k \in \mathcal{H}^r$ and $\|F_k\|_r = \|\mathcal{A}^{r/2} F_k\| = \|\omega_k^r F_k\| = \omega_k^r$. If $r > \frac{d}{2} + j$ these facts and the imbedding (B.19) imply $F_k \in C^j(\overline{\Omega})$; since this holds for each $j \in \mathbf{N}$, we have

$$F_k \in C^\infty(\overline{\Omega}). \quad (\text{B.22})$$

To go on, let us note that (B.19) means the following: for each $j \in \mathbf{N}$ and $r > j + \frac{d}{2}$, there exists a constant $\Lambda_{j,r} \in (0, +\infty)$ such that

$$\|f\|_{C^j} \leq \Lambda_{j,r} \|f\|_r \quad \text{for all } f \in \mathcal{H}^r. \quad (\text{B.23})$$

This statement with $f = F_k$ gives $\|F_k\|_{C^j} \leq \Lambda_{j,r} \omega_k^r$ for $j \in \mathbf{N}$, $r > j + \frac{d}{2}$ or, equivalently,

$$\|F_k\|_{C^j} \leq \Lambda_{j,\eta} \omega_k^{j+d/2+\eta} \quad \text{for all } j \in \mathbf{N}, \eta > 0 \quad (\text{B.24})$$

(where $\Lambda_{j,\eta}$ stands for $\Lambda_{j,j+d/2+\eta}$). These results should be kept in mind in the sequel, together with the already mentioned Weyl asymptotics (3.10)

$$\omega_k \sim C k^{1/d} \quad \text{for } k \rightarrow +\infty,$$

where $C := 2\sqrt{\pi} \Gamma(d/2+1)^{1/d} \text{Vol}(\Omega)^{-1/d}$ ⁽³⁷⁾.

³⁶In order to derive Eq. (B.19) one proceeds similarly to footnote 33 using the embeddings $\mathcal{H}^r \hookrightarrow H^r(\Omega)$ and $H^r(\Omega) \hookrightarrow C^j(\overline{\Omega})$; these follow from Theorem 3 on p. 155 of [71] and some standard interpolation theory (see, e.g., [10, 67]).

³⁷Concerning (3.10), we have already given references [32] [71]. Eq. (B.24) is known as well in the literature, see [61, 97, 104]; here we have proposed an alternative derivation of this result just because it arose naturally from the general framework of the present appendix.

On the eigenfunction expansion for the Dirichlet kernel. In the case we are considering, the eigenfunction expansion (3.18) for the Dirichlet kernel takes the form

$$D_s(\mathbf{x}, \mathbf{y}) = \sum_{k=1}^{+\infty} \frac{1}{\omega_k^{2s}} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) ; \quad (\text{B.25})$$

hereafter we discuss the absolute convergence of this expansion with respect to the norm (B.18) of $C^j(\overline{\Omega} \times \overline{\Omega})$, showing that

$$\sum_{k=1}^{+\infty} \frac{1}{|\omega_k^{2s}|} \|F_k(\cdot) \overline{F_k}(\cdot)\|_{C^j} < +\infty \quad \text{if } \Re s > d + \frac{j}{2} . \quad (\text{B.26})$$

To this purpose, we first note that

$$\begin{aligned} \sum_{k=1}^{+\infty} \frac{1}{|\omega_k^{2s}|} \|F_k(\cdot) \overline{F_k}(\cdot)\|_{C^j} &= \sum_{k=1}^{+\infty} \frac{1}{\omega_k^{2\Re s}} \max_{|\alpha|+|\beta| \leq j} \max_{\mathbf{x} \in \overline{\Omega}, \mathbf{y} \in \overline{\Omega}} |\partial^\alpha F_k(\mathbf{x}) \partial^\beta \overline{F_k}(\mathbf{y})| \leq \\ &\leq \sum_{k=1}^{+\infty} \frac{1}{\omega_k^{2\Re s}} \max_{j_1+j_2 \leq j} \|F_k\|_{j_1} \|F_k\|_{j_2} . \end{aligned} \quad (\text{B.27})$$

Let $k \in \{1, 2, 3, \dots\}$, $j_1 + j_2 \leq j$ and $\eta > 0$. Due to the estimate (B.24) we have

$$\|F_k\|_{j_1} \|F_k\|_{j_2} \leq \Lambda_{j_1 \eta} \Lambda_{j_2 \eta} \omega_k^{d+j_1+j_2+2\eta} ;$$

but (recalling that $\omega_k \geq \omega_1$)

$$\omega_k^{j_1+j_2} = \left(\frac{\omega_k}{\omega_1}\right)^{j_1+j_2} \omega_1^{j_1+j_2} \leq \left(\frac{\omega_k}{\omega_1}\right)^j \omega_1^{j_1+j_2} = \frac{\omega_k^j}{\omega_1^{j-j_1-j_2}} ,$$

whence

$$\|F_k\|_{j_1} \|F_k\|_{j_2} \leq \frac{\Lambda_{j_1 \eta} \Lambda_{j_2 \eta}}{\omega_1^{j-j_1-j_2}} \omega_k^{d+j+2\eta} . \quad (\text{B.28})$$

Inserting Eq. (B.28) into Eq. (B.27) we obtain the following, for all $\eta > 0$:

$$\sum_{k=1}^{+\infty} \frac{1}{|\omega_k^{2s}|} \|F_k(\cdot) \overline{F_k}(\cdot)\|_{C^j} \leq \left(\max_{j_1+j_2 \leq j} \frac{\Lambda_{j_1 \eta} \Lambda_{j_2 \eta}}{\omega_1^{j-j_1-j_2}} \right) \sum_{k=1}^{+\infty} \frac{1}{\omega_k^{2\Re s - d - j - 2\eta}} . \quad (\text{B.29})$$

From here to the end of the paragraph we assume

$$\Re s > d + \frac{j}{2} ; \quad (\text{B.30})$$

hereafter we show that the series on the left-hand side of Eq. (B.29) converges for a suitable $\eta > 0$, a fact yielding the thesis (B.26). Indeed, due to Eq. (B.30) there is $\eta > 0$ such that

$$\Re s = d + \frac{j}{2} + \left(\frac{d}{2} + 1 \right) \eta ; \quad (\text{B.31})$$

expressing $\Re s$ in this way, and using the Weyl asymptotics (3.10) we get

$$\frac{1}{\omega_k^{2\Re s - d - j - 2\eta}} = \frac{1}{\omega_k^{(1+\eta)d}} \sim \frac{1}{C^{(1+\eta)d} k^{1+\eta}} \quad \text{for } k \rightarrow +\infty , \quad (\text{B.32})$$

which implies convergence for the series on the left-hand side of Eq. (B.29).

On the eigenfunction expansions for the heat and cylinder kernels. In the type of configuration under analysis, the expansions (3.47) and (3.48) can be re-expressed, respectively, as

$$K(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \sum_{k=1}^{+\infty} e^{-\mathbf{t}\omega_k^2} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) , \quad (\text{B.33})$$

$$T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \sum_{k=1}^{+\infty} e^{-\mathbf{t}\omega_k} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y}) . \quad (\text{B.34})$$

Using considerations similar to the ones of paragraph c), one shows the absolute convergence of these expansions in the norm (B.18) of $C^j(\overline{\Omega} \times \overline{\Omega})$, for all $\mathbf{t} > 0$ and for each $j \in \mathbf{N}$:

$$\sum_{k=1}^{+\infty} e^{-\mathbf{t}\omega_k^2} \|F_k(\cdot) \overline{F_k}(\cdot)\|_{C^j} < +\infty , \quad \sum_{k=1}^{+\infty} e^{-\mathbf{t}\omega_k} \|F_k(\cdot) \overline{F_k}(\cdot)\|_{C^j} < +\infty . \quad (\text{B.35})$$

Another result. In subsection 3.12 it is stated that, under the assumptions (3.9) for $\mathcal{A}$ and considering a complete orthonormal set $(F_k)_{k=1,2,3,\dots}$ with the usual features for this case, the function

$$\hat{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) := \sum_{k=1}^{+\infty} e^{-\omega_k \mathbf{t}} |F_k(\mathbf{x})| |F_k(\mathbf{y})| \quad (\text{B.36})$$

admits a uniform bound

$$\hat{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) \leq \check{T}(\mathbf{t}) < +\infty \quad \text{for all } \mathbf{x}, \mathbf{y} \in \overline{\Omega} \text{ and } \mathbf{t} > 0 . \quad (\text{B.37})$$

The proof of this statements starts from the inequality (B.24) with $j = 0$ and any $\eta > 0$, this implies $|F_k(\mathbf{x})|, |F_k(\mathbf{y})| \leq \Lambda_{0\eta} \omega_k^{d/2+\eta}$, whence

$$\hat{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}) \leq \check{T}(\mathbf{t}) \text{ for all } \mathbf{x}, \mathbf{y} \in \overline{\Omega} \text{ and } \mathbf{t} > 0 , \quad \check{T}(\mathbf{t}) := \Lambda_{0\eta}^2 \sum_{k=1}^{+\infty} e^{-\omega_k \mathbf{t}} \omega_k^{d+2\eta} . \quad (\text{B.38})$$

Finally, the Weyl estimates (3.10) ensure $\check{T}(\mathbf{t}) < +\infty$ for each $\mathbf{t} > 0$.

C Appendix. A $(d+1)$ -dimensional Green function and its relation with the cylinder kernel

Let $\Omega \subset \mathbf{R}^d$ be an open set, and let $\mathcal{A}$ be a strictly positive selfadjoint operator in $L^2(\Omega)$ (keeping into account suitable boundary conditions on $\partial\Omega$). We consider the cylinder kernel $T(\mathbf{t}; \mathbf{x}, \mathbf{y}) := (e^{-\mathbf{t}\sqrt{\mathcal{A}}})(\mathbf{x}, \mathbf{y})$ ($\mathbf{x}, \mathbf{y} \in \Omega$, $\mathbf{t} \in (0, +\infty)$).

The aim of the present appendix is to illustrate the fact mentioned in subsection 3.11, namely, the possibility to relate T to a $(d+1)$ dimensional Green function. To this purpose we consider in $\mathbf{R}^{d+1}$ the domain $\mathcal{O} := (0, +\infty) \times \Omega \ni (\mathbf{t}, \mathbf{x})$ and the Hilbert space $L^2(\mathcal{O})$; we introduce therein the operator

$$\mathcal{P} := -\partial_{\mathbf{t}} + \mathcal{A} , \quad (\text{C.1})$$

with suitable boundary conditions on $\partial\mathcal{O} := (\{0\} \times \Omega) \cup ((0, +\infty) \times \partial\Omega)$. More precisely, we assume Dirichlet boundary conditions on $\{0\} \times \Omega$ and the previously given boundary conditions for $\mathcal{A}$ on $(0, +\infty) \times \partial\Omega$. The operator $\mathcal{P}$ is selfadjoint; in the sequel we will prove that it is strictly positive.

Let us introduce the Green function

$$G(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y}) := \mathcal{P}^{-1}((\mathbf{t}, \mathbf{x}), (\mathbf{t}', \mathbf{y})) \equiv \langle \delta_{\mathbf{t}} \delta_{\mathbf{x}} | \mathcal{P}^{-1} \delta_{\mathbf{t}'} \delta_{\mathbf{y}} \rangle ; \quad (\text{C.2})$$

this is characterized by the equation

$$(-\partial_{\mathbf{t}} + \mathcal{A}_{\mathbf{x}})G(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y}) = \delta(\mathbf{t} - \mathbf{t}')\delta(\mathbf{x} - \mathbf{y}) , \quad (\text{C.3})$$

and by the boundary conditions prescribed on $\partial\mathcal{O}$. We claim that the cylinder kernel T is related to G by

$$\partial_{\mathbf{t}'} G(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y})|_{\mathbf{t}'=0} = T(\mathbf{t}; \mathbf{x}, \mathbf{y}) . \quad (\text{C.4})$$

To prove this (and the previous statement on the strict positivity of $\mathcal{P}$), let $(F_k)_{k \in \mathcal{K}}$ be a complete orthonormal system of eigenfunctions of $\mathcal{A}$ with corresponding eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$; clearly, the functions $\mathbf{t} \in (0, +\infty) \mapsto \sqrt{\frac{2}{\pi}} \sin(\lambda \mathbf{t})$ ($\lambda \in (0, +\infty)$) are a complete orthonormal system in $L^2((0, +\infty))$ and are eigenfunctions of $-\partial_{\mathbf{t}}$ vanishing for $\mathbf{t} = 0$. These facts ensure that the family of functions

$$Y_{(\lambda, k)}(\mathbf{t}, \mathbf{x}) := \sqrt{\frac{2}{\pi}} \sin(\lambda \mathbf{t}) F_k(\mathbf{x}) \quad \text{for } (\lambda, k) \in (0, +\infty) \times \mathcal{K} \quad (\text{C.5})$$

is a complete orthonormal system of eigenfunctions of $\mathcal{P}$, with

$$\mathcal{P} Y_{(\lambda, k)} = (\lambda^2 + \omega_k^2) Y_{(\lambda, k)} . \quad (\text{C.6})$$

We recall that we are assuming $\omega_k^2 \geq \varepsilon^2$ for some $\varepsilon > 0$; the eigenvalues $(\lambda^2 + \omega_k^2)$ also have ε^2 as a lower bound, so $\mathcal{P}$ is strictly positive.

The Green function of Eq. (C.2) can be expressed via the equation

$$G(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y}) = \int_{(0, +\infty) \times \mathcal{K}} \frac{d\lambda \, dk}{\lambda^2 + \omega_k^2} \left(\sqrt{\frac{2}{\pi}} \sin(\lambda \mathbf{t}) F_k(\mathbf{x}) \right) \left(\sqrt{\frac{2}{\pi}} \sin(\lambda \mathbf{t}') \overline{F}_k(\mathbf{y}) \right), \quad (\text{C.7})$$

which, evaluating explicitly the integral in λ ⁽³⁸⁾, reduces to

$$G(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y}) = \int_{\mathcal{K}} dk \frac{e^{-\omega_k |\mathbf{t} - \mathbf{t}'|} - e^{-\omega_k (\mathbf{t} + \mathbf{t}')}}{2\omega_k} F_k(\mathbf{x}) \overline{F}_k(\mathbf{y}). \quad (\text{C.8})$$

To go on, let us differentiate both sides of Eq. (C.8) with respect to $\mathbf{t}'$; this gives

$$\partial_{\mathbf{t}'} G(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y}) = \int_{\mathcal{K}} dk \left[\frac{\text{sgn}(\mathbf{t} - \mathbf{t}')}{2} e^{-\omega_k |\mathbf{t} - \mathbf{t}'|} + \frac{1}{2} e^{-\omega_k (\mathbf{t} + \mathbf{t}')} \right] F_k(\mathbf{x}) \overline{F}_k(\mathbf{y}). \quad (\text{C.9})$$

Setting $\mathbf{t}' = 0$ (and recalling that $\mathbf{t} > 0$), the last equation yields

$$\partial_{\mathbf{t}'} G(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y})|_{\mathbf{t}'=0} = \int_{\mathcal{K}} dk e^{-\omega_k \mathbf{t}} F_k(\mathbf{x}) \overline{F}_k(\mathbf{y}). \quad (\text{C.10})$$

The right-hand side of the above equality is just the representation of the cylinder kernel T given by Eq. (3.48); thus we have proved Eq. (C.4).

An example. Hereafter, the approach based on (C.4) is used to derive the expression (3.60) for the cylinder kernel T in the case where

$$\Omega := \mathbf{R}^d, \quad \mathcal{A} := -\Delta. \quad (\text{C.11})$$

The associated $(d+1)$ dimensional domain and the operator $\mathcal{P}$ are

$$\mathcal{O} := (0, +\infty) \times \mathbf{R}^d, \quad \mathcal{P} := -\partial_{\mathbf{t}\mathbf{t}} - \Delta = -\Delta_{d+1}, \quad (\text{C.12})$$

with Dirichlet boundary conditions on $\partial\mathcal{O}$, which coincides with the hyperplane $\{0\} \times \mathbf{R}^d$; of course, in the above Δ_{d+1} indicates the $(d+1)$ -dimensional Laplacian. The Green function $\mathcal{G}$ of $-\Delta_{d+1}$ on the half-space $\mathcal{O}$ can be obtained by the familiar method of images from the Green function $\mathcal{G}_0$ of $-\Delta_{d+1}$ on the full space $\mathbf{R}^{d+1}$; thus

$$\mathcal{G}(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y}) = \mathcal{G}_0(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y}) - \mathcal{G}_0(\mathbf{t}, \mathbf{x}; -\mathbf{t}', \mathbf{y}), \quad (\text{C.13})$$

³⁸Just observe that, by symmetry arguments and the residue theorem, we have

$$\int_0^{+\infty} d\lambda \frac{\sin(\lambda \mathbf{t}) \sin(\lambda \mathbf{t}')}{\lambda^2 + \omega_k^2} = \frac{1}{2} \int_{-\infty}^{+\infty} d\lambda \frac{\sin(\lambda \mathbf{t}) \sin(\lambda \mathbf{t}')}{\lambda^2 + \omega_k^2} = \frac{\pi}{4\omega_k} \left(e^{-\omega_k |\mathbf{t} - \mathbf{t}'|} - e^{-\omega_k (\mathbf{t} + \mathbf{t}')} \right).$$

$$\mathcal{G}_0(\mathbf{t}, \mathbf{x}; \mathbf{t}', \mathbf{y}) := \begin{cases} \frac{1}{2\pi} \ln((\mathbf{t} - \mathbf{t}')^2 + (\mathbf{x} - \mathbf{y})^2) & \text{if } d = 1, \\ \frac{\Gamma(\frac{d+1}{2})}{(d-1)2\pi^{\frac{d+1}{2}}((\mathbf{t} - \mathbf{t}')^2 + |\mathbf{x} - \mathbf{y}|^2)^{\frac{d-1}{2}}} & \text{if } d \geq 2. \end{cases} \quad (\text{C.14})$$

Inserting Eq.s (C.13) (C.14) into Eq. (C.4) we obtain for the cylinder kernel T the expression (3.60)

$$T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \frac{\Gamma(\frac{d+1}{2}) \mathbf{t}}{\pi^{\frac{d+1}{2}} (\mathbf{t}^2 + |\mathbf{x} - \mathbf{y}|^2)^{\frac{d+1}{2}}}$$

(in all dimensions, including $d = 1$).

D Appendix. Derivation of Eq. (3.90)

Let $\mathbf{t} \mapsto h(\mathbf{t})$ be a complex-valued function, analytic in a neighborhood of $[0, +\infty)$ and exponentially vanishing for $\Re \mathbf{t} \rightarrow +\infty$. For any given $s \in \mathbf{C}$ with $\Re s > 0$, consider the integral

$$I(s, h) := \int_{\mathfrak{H}} d\mathbf{t} \mathbf{t}^{s-1} h(\mathbf{t}) , \quad (\text{D.1})$$

where $\mathfrak{H}$ is a Hankel contour (see below Eq. (3.89) and Fig. 1 on page 32 for the description of this path); the complex power $\mathbf{t}^{s-1}$ in the above equation is defined following Eq.s (3.15) (3.16). For any $\delta > 0$, $\mathfrak{H}$ is homotopic to the path $\mathfrak{H}_\delta$ described as follows:

$$\begin{aligned} \mathfrak{H}_\delta &= \mathfrak{H}_\delta^+ \cup \mathfrak{H}_\delta^0 \cup \mathfrak{H}_\delta^- , \quad \text{with} \\ \mathfrak{H}_\delta^\pm &:= \{\mathbf{t} \in \mathbf{C} \mid \mathbf{t} = v \pm i\delta, \ v \in [0, +\infty)\} , \\ \mathfrak{H}_\delta^0 &:= \{\mathbf{t} \in \mathbf{C} \mid \mathbf{t} = \delta e^{i\theta}, \ \theta \in (\pi/2, 3\pi/2)\} . \end{aligned} \quad (\text{D.2})$$

Due to this remark and to the analyticity of h we can replace $\mathfrak{H}$ with $\mathfrak{H}_\delta$ in Eq. (D.1); so, for any $\delta > 0$ we have

$$I(s, h) = I_\delta^+(s, h) + I_\delta^0(s, h) + I_\delta^-(s, h) , \quad (\text{D.3})$$

$$I_\delta^\pm(s, h) := \mp \int_0^{+\infty} dv (v \pm i\delta)^{s-1} h(v \pm i\delta) , \quad I_\delta^0(s, h) := i \int_{\pi/2}^{3\pi/2} d\theta (\delta e^{i\theta})^s h(\delta e^{i\theta}) .$$

We are now going to consider the limit $\delta \rightarrow 0^+$. Notice that in this limit $(v+i\delta)^{s-1} \rightarrow v^{s-1}$ while $(v-i\delta)^{s-1} \rightarrow e^{2i\pi(s-1)} v^{s-1} = e^{2i\pi s} v^{s-1}$; moreover, $h(v \pm i\delta) \rightarrow h(v)$. Due to these results, we easily infer

$$\lim_{\delta \rightarrow 0^+} I_\delta^+(s, h) = - \int_0^{+\infty} dv v^{s-1} h(v) , \quad (\text{D.4})$$

$$\lim_{\delta \rightarrow 0^+} I_\delta^-(s, h) = e^{2i\pi s} \int_0^{+\infty} dv v^{s-1} h(v) . \quad (\text{D.5})$$

Passing to the integral $I_\delta^0(s, h)$, noting that $|h(\delta e^{i\theta})| \leq C$ for small δ and recalling that $\Re s > 0$ by hypothesis, we obtain

$$|I_\delta^0(s, h)| \leq C \delta^{\Re s} \int_{\pi/2}^{3\pi/2} d\theta e^{-(\Im s)\theta} \rightarrow 0 \quad \text{for } \delta \rightarrow 0^+ . \quad (\text{D.6})$$

Summing up, in the limit $\delta \rightarrow 0^+$ one obtains from Eq. (D.3) that

$$I(s, h) = (e^{2i\pi s} - 1) \int_0^{+\infty} dv v^{s-1} h(v) ; \quad (\text{D.7})$$

noting that $e^{2i\pi s} - 1 = 2ie^{i\pi s} \sin(\pi s)$, the above relation yields

$$I(s, h) = 2ie^{i\pi s} \sin(\pi s) \int_0^{+\infty} dv v^{s-1} h(v) , \quad (\text{D.8})$$

which is equivalent to Eq. (3.90) for $\Re s > 0$, $s \notin \{1, 2, 3, \dots\}$. For more information concerning the Mellin transform and its contour integral representations see, e.g., [42, 98].

E Appendix. Derivation of Eq.s (3.112-3.114)

We refer to the framework of subsection 3.18 about the slab $\Omega = \Omega_1 \times \mathbf{R}^{d_2}$; we retain all the assumptions and notations of the cited subsection. In particular, $(\mathfrak{F}_{k_1})_{k_1 \in \mathcal{K}_1}$ is a complete orthonormal system of eigenfunctions of $\mathcal{A}_1$ with related eigenvalues $\varpi_{k_1}^2$. A complete orthonormal set of eigenfunctions $(F_k)_{k \in \mathcal{K}}$ with eigenvalues $(\omega_k^2)_{k \in \mathcal{K}}$ for the operator $\mathcal{A}$ in $L^2(\Omega)$ is given by

$$F_k(\mathbf{x}) = \mathfrak{F}_{k_1}(\mathbf{x}_1) \frac{e^{i\mathbf{k}_2 \cdot \mathbf{x}_2}}{(2\pi)^{d_2/2}} , \quad \omega_k^2 = \varpi_{k_1}^2 + |\mathbf{k}_2|^2 \quad (\text{E.1})$$

for $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2)$ and $k = (k_1, \mathbf{k}_2) \in \mathcal{K}_1 \times \mathbf{R}^{d_2}$

In the present case, the eigenfunction expansion (3.18) of the Dirichlet kernel at any two points $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2)$ and $\mathbf{y} = (\mathbf{y}_1, \mathbf{y}_2)$ can be re-expressed as follows:

$$\begin{aligned} D_s(\mathbf{x}_1, \mathbf{x}_2; \mathbf{y}_1, \mathbf{y}_2) &= \int_{\mathcal{K}_1 \times \mathbf{R}^{d_2}} \frac{dk_1 d\mathbf{k}_2}{(\varpi_{k_1}^2 + |\mathbf{k}_2|^2)^s} \mathfrak{F}_{k_1}(\mathbf{x}_1) \overline{\mathfrak{F}_{k_1}(\mathbf{y}_1)} \frac{e^{i\mathbf{k}_2 \cdot (\mathbf{x}_2 - \mathbf{y}_2)}}{(2\pi)^{d_2}} = \\ &= \int_{\mathcal{K}_1 \times \mathbf{R}^{d_2}} \frac{dk_1 d\mathbf{h}}{\varpi_{k_1}^{2s-d_2} (|\mathbf{h}|^2 + 1)^s} \mathfrak{F}_{k_1}(\mathbf{x}_1) \overline{\mathfrak{F}_{k_1}(\mathbf{y}_1)} \frac{e^{i\varpi_{k_1} \mathbf{h} \cdot (\mathbf{x}_2 - \mathbf{y}_2)}}{(2\pi)^{d_2}} \end{aligned} \quad (\text{E.2})$$

where, in the last passage, we performed the change of variables $\mathbf{k}_2 = \varpi_{k_1} \mathbf{h}$. On the other hand, it is known that, for any $\mathbf{z} \in \mathbf{R}^{d_2}$,

$$\int_{\mathbf{R}^{d_2}} \frac{d\mathbf{h}}{(2\pi)^{d_2}} \frac{e^{i\mathbf{h} \cdot \mathbf{z}}}{(|\mathbf{h}|^2 + 1)^s} = \frac{|\mathbf{z}|^{s-\frac{d_2}{2}}}{(2\pi)^{d_2/2} 2^{s-1} \Gamma(s)} K_{s-\frac{d_2}{2}}(|\mathbf{z}|) \quad \text{if } s \in \mathbf{C}, \Re s > \frac{d_2}{2} , \quad (\text{E.3})$$

with K_ν denoting the modified Bessel function of the second kind of order $\nu \in \mathbf{C}$ (see, e.g., [7, 80, 103]). Thus

$$D_s(\mathbf{x}_1, \mathbf{x}_2; \mathbf{y}_1, \mathbf{y}_2) = \int_{\mathcal{K}_1} \frac{dk_1}{\varpi_{k_1}^{2s-d_1}} \mathfrak{F}_{k_1}(\mathbf{x}_1) \overline{\mathfrak{F}_{k_1}}(\mathbf{y}_1) \frac{(\varpi_{k_1}|\mathbf{x}_2 - \mathbf{y}_2|)^{s-\frac{d_2}{2}}}{(2\pi)^{d_2/2} 2^{s-1}\Gamma(s)} K_{s-\frac{d_2}{2}}(\varpi_{k_1}|\mathbf{x}_2 - \mathbf{y}_2|) . \quad (\text{E.4})$$

For the sake of brevity, for any $\nu \in \mathbf{C}$, we put

$$\mathfrak{G}_\nu : (0, +\infty) \rightarrow \mathbf{C} , \quad z \mapsto \mathfrak{G}_\nu(z) := z^{\nu/2} K_\nu(\sqrt{z}) ; \quad (\text{E.5})$$

due to the asymptotic behaviour of the Bessel function K_ν near zero (see [80], p.252, Eq.10.30.2), for $\Re \nu > 0$ this function has continuous extension to $z = 0$, given by

$$\mathfrak{G}_\nu(0) = 2^{\nu-1}\Gamma(\nu) . \quad (\text{E.6})$$

To proceed, note that Eq. (E.4) can then be rephrased in terms of the function $\mathfrak{G}_\nu$ as follows:

$$\begin{aligned} D_s(\mathbf{x}_1, \mathbf{x}_2; \mathbf{y}_1, \mathbf{y}_2) &= \hat{D}_s(\mathbf{x}_1, \mathbf{y}_1; |\mathbf{x}_2 - \mathbf{y}_2|^2) , \\ \hat{D}_s(\mathbf{x}_1, \mathbf{y}_1; q) &= \frac{2^{1-s}}{(2\pi)^{d_2/2}\Gamma(s)} \int_{\mathcal{K}_1} \frac{dk_1}{\varpi_{k_1}^{2s-d_2}} \mathfrak{F}_{k_1}(\mathbf{x}_1) \overline{\mathfrak{F}_{k_1}}(\mathbf{y}_1) \mathfrak{G}_{s-\frac{d_1}{2}}(\varpi_{k_1}^2 q) . \end{aligned} \quad (\text{E.7})$$

This proves Eq. (3.112), also giving an explicit expression for the function $\hat{D}_s$. To proceed note that, due to the well-known facts on the derivatives of the Bessel function K_ν for any $\nu \in \mathbf{C}$ (see [80], p.252, Eq.10.29.4), we have

$$\begin{aligned} \frac{d\mathfrak{G}_\nu}{dz}(z) &= \frac{d}{dv} \left(v^\nu K_\nu(v) \right) \Big|_{v=\sqrt{z}} \frac{1}{2\sqrt{z}} = \left(-v^\nu K_{\nu-1}(v) \right)_{v=\sqrt{z}} \frac{1}{2\sqrt{z}} = \\ &= -\frac{1}{2} z^{\frac{\nu-1}{2}} K_{\nu-1}(\sqrt{z}) = -\frac{1}{2} \mathfrak{G}_{\nu-1}(z) . \end{aligned} \quad (\text{E.8})$$

Using the above identity it can be proved by induction that

$$\frac{d^n \mathfrak{G}_\nu}{dz^n}(z) = \left(-\frac{1}{2} \right)^n \mathfrak{G}_{\nu-n}(z) \quad \text{for } n \in \{1, 2, 3, \dots\} ; \quad (\text{E.9})$$

this fact, along with Eq. (E.7), implies

$$\frac{\partial^n \hat{D}_s}{\partial q^n}(\mathbf{x}_1, \mathbf{y}_1; q) = \frac{(-1)^n 2^{1-s-n}}{(2\pi)^{d_2/2}\Gamma(s)} \int_{\mathcal{K}_1} \frac{dk_1}{\varpi_{k_1}^{2s-d_2-2n}} \mathfrak{F}_{k_1}(\mathbf{x}_1) \overline{\mathfrak{F}_{k_1}}(\mathbf{y}_1) \mathfrak{G}_{s-\frac{d_2}{2}-n}(\varpi_{k_1}^2 q) . \quad (\text{E.10})$$

Now, recalling Eq. (E.6) we conclude that, for any $n \in \{1, 2, 3, \dots\}$ and any $s \in \mathbb{C}$ with $\Re s > \frac{d}{2} + n$,

$$\frac{\partial^n \hat{D}_s}{\partial q^n}(\mathbf{x}_1, \mathbf{y}_1; 0) = \frac{(-1)^n \Gamma(s - \frac{d_2}{2} - n)}{(4\pi)^{d_2/2} 4^n \Gamma(s)} \int_{\mathcal{K}_1} \frac{dk_1}{\varpi_{k_1}^{2s-d_2-2n}} \mathfrak{F}_{k_1}(\mathbf{x}_1) \overline{\mathfrak{F}_{k_1}(\mathbf{y}_1)}. \quad (\text{E.11})$$

Due to a representation analogous to (3.18) holding for the reduced Dirichlet kernel $D_s^{(1)}$, Eq. (E.11) implies Eq. (3.114). Finally, Eq. (3.113) is just the case $n = 0$ of Eq. (3.114).

F Appendix. Some results on boundary forces

As in the final part of subsections 4.3 and 4.4, we work on a domain Ω with Dirichlet boundary conditions.

F.1 Derivation of Eq. (4.25) for the pressure. We start from Eq. (4.24), holding for general boundary conditions. In the Dirichlet case that we are considering, only the terms involving mixed derivatives (with respect to both $\mathbf{x}$ and $\mathbf{y}$) of the Dirichlet kernel yield non-vanishing contributions on the boundary $\partial\Omega$ ⁽³⁹⁾; thus, for any $\mathbf{x} \in \partial\Omega$, Eq. (4.24) reduces to

$$p_i^u(\mathbf{x}) = \kappa^u \left[\left(-\left(\frac{1}{4} - \xi\right) \delta_{ij} \partial^{x^\ell} \partial_{y^\ell} + \left(\frac{1}{2} - \xi\right) \partial_{x^i y^j} \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} n^j(\mathbf{x}). \quad (\text{F.1})$$

We now claim that the terms proportional to ξ in Eq. (F.1) vanish, i.e., that

$$\left[\left(\delta_{ij} \partial^{x^\ell} \partial_{y^\ell} - \partial_{x^i y^j} \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}} n^j(\mathbf{x}) = 0 \quad \text{for all } \mathbf{x} \in \partial\Omega; \quad (\text{F.2})$$

this will yield Eq. (4.25). Using the eigenfunction expansion (3.18) for the Dirichlet kernel, we see that Eq. (F.2) holds if we are able to prove that

$$\left(\delta_{ij} \partial^\ell F_k \partial_\ell \overline{F_k} - \frac{1}{2} \partial_i F_k \partial_j \overline{F_k} - \frac{1}{2} \partial_j F_k \partial_i \overline{F_k} \right)(\mathbf{x}) n^j(\mathbf{x}) = 0 \quad \text{for } k \in \mathcal{K}, \mathbf{x} \in \partial\Omega. \quad (\text{F.3})$$

The simplest way to prove Eq. (F.3) is to derive the following, equivalent statement: for all $k \in \mathcal{K}$ and all (sufficiently smooth) vector field $\mathfrak{S} \equiv (\mathfrak{S}^i) : \partial\Omega \rightarrow \mathbf{R}^d$,

$$\int_{\partial\Omega} da \mathfrak{S}^i \left(\delta_{ij} \partial^\ell F_k \partial_\ell \overline{F_k} - \frac{1}{2} \partial_i F_k \partial_j \overline{F_k} - \frac{1}{2} \partial_j F_k \partial_i \overline{F_k} \right) n^j = 0. \quad (\text{F.4})$$

³⁹One can easily infer this statement using the eigenfunction expansion (3.18).

Let us sketch a derivation of Eq. (F.4), for given $k \in \mathcal{K}$ and $\mathfrak{S} : \partial\Omega \rightarrow \mathbf{R}^d$. To this purpose we consider a smooth extension of $\mathfrak{S}$ to a vector field $\mathfrak{S} : \partial\Omega \cup \Omega \rightarrow \mathbf{R}^d$ and fix the attention on the integral

$$\frac{1}{2} \int_{\Omega} d\mathbf{x} (\partial^j \partial_j)(\partial_i \mathfrak{S}^i) |F_k|^2 = \frac{1}{2} \int_{\Omega} d\mathbf{x} \partial_i (\partial^j \partial_j \mathfrak{S}^i) |F_k|^2 \quad (\text{F.5})$$

(note that $\partial^j \partial_j = \Delta$). We re-express both sides in the above identity integrating by parts with respect to all the derivatives appearing therein ⁽⁴⁰⁾, considering them in the two orders proposed in the two sides; some of the boundary terms arising in this way vanish since F_k is zero on $\partial\Omega$. The difference between the two expressions thus obtained, which is obviously zero, is found to coincide with the left-hand side of Eq. (F.4).

F.2 Derivation of Eq. (4.29). Let us stick to the framework of subsection 4.4 in which the domain Ω is bounded, and Dirichlet boundary conditions are prescribed; the operator $\mathcal{A} = -\Delta + V$, acting in $L^2(\Omega)$, has a complete orthonormal system of eigenfunctions F_k with eigenvalues ω_k^2 , labelled by a countable set $\mathcal{K}$. We consider, for small $\epsilon > 0$, a deformation of the domain Ω of the form (4.26-4.27), controlled by a vector field $\mathfrak{S}$ on $\mathbf{R}^d$. The operator $\mathcal{A}_\epsilon := -\Delta + V$ acting in $L^2(\Omega_\epsilon)$ has a complete orthonormal system of eigenfunctions $F_{\epsilon,k}$ with eigenvalues $\omega_{\epsilon,k}^2$. In subsection 4.4 we have already considered the regularized bulk energy corresponding to Ω_ϵ ; this is (see Eq. (4.28))

$$E_\epsilon^u = \frac{\kappa^u}{2} \sum_{k \in \mathcal{K}} (\omega_{\epsilon,k}^2)^{\frac{1-u}{2}}.$$

We now consider the limit $\epsilon \rightarrow 0$, and expand everything to the first order in ϵ . Eq. (4.29) that we want to derive concerns the expansion in ϵ of the bulk energy E_ϵ^u ; as already mentioned, Eq. (4.28) can be used to make contact with the expansion of the eigenvalues $\omega_{\epsilon,k}^2$, on which we now fix our attention.

The variation of the eigenvalues under a deformation of the spatial domain for the Dirichlet Laplacian (or similar operators) has been the subject of classical investigations. Here we refer to the book of Rellich [85] (see Chapter II, at the end of §6), whose results can be expressed in this way with our notations:

$$\omega_{\epsilon,k}^2 = \omega_k^2 + \epsilon \varpi_k^2 + O(\epsilon^2) \quad \text{with} \quad \varpi_k^2 := \langle F_k | \mathcal{B} F_k \rangle, \quad (\text{F.6})$$

where $\mathcal{B}$ is the selfadjoint operator in $L^2(\Omega)$ defined by

$$\mathcal{B}f := \partial_i \left((\partial^i \mathfrak{S}^j + \partial^j \mathfrak{S}^i) \partial_i f \right) + \left(\frac{1}{2} \Delta \partial_\ell \mathfrak{S}^\ell + \mathfrak{S}^\ell \partial_\ell V \right) f \quad (\text{F.7})$$

⁴⁰See the footnote on page 39.

(as matter of fact, [85] gives the expression of $\mathcal{B}$ for $V = 0$, but the extension to a nonzero V is straightforward). Keeping in mind these facts, we return to Eq. (4.28) for the regularized bulk energy; this implies

$$E_\epsilon^u = E^u + \epsilon \mathfrak{E}^u + O(\epsilon^2), \quad \mathfrak{E}^u := \left(\frac{1-u}{2} \right) \frac{\kappa^u}{2} \sum_{k \in \mathcal{K}} \omega_k^{-1-u} \varpi_k^2. \quad (\text{F.8})$$

To go on, we note that the definition of ϖ_k^2 in Eq. (F.6), with $\mathcal{B}$ as in Eq. (F.7), yields

$$\varpi_k^2 = \int_{\Omega} d\mathbf{x} \, \overline{F_k} \left[\partial_i \left((\partial^i \mathfrak{S}^j + \partial^j \mathfrak{S}^i) \partial_i F_k \right) + \left(\frac{1}{2} \Delta \partial_\ell \mathfrak{S}^\ell + \mathfrak{S}^\ell \partial_\ell V \right) F_k \right]. \quad (\text{F.9})$$

The above result can be re-expressed in terms of surface integrals on $\partial\Omega$ via suitable integrations by parts⁽⁴¹⁾; while making these computations, one must use the identity $\Delta F_k = (V - \omega_k^2) F_k$ (and its complex conjugate), and recall that F_k vanishes on $\partial\Omega$. In this way we obtain

$$\varpi_k^2 = \int_{\partial\Omega} da(\mathbf{x}) \, n^j \mathfrak{S}^i \left[\delta_{ij} \partial^\ell \overline{F_k} \partial_\ell F_k - (\partial_i \overline{F_k} \partial_j F_k + \partial_j \overline{F_k} \partial_i F_k) \right]. \quad (\text{F.10})$$

We plug this relation into Eq. (F.8), exchange the summation with the integration and use the expansion (3.18), which in this case reads $D_s(\mathbf{x}, \mathbf{y}) = \sum_{k \in \mathcal{K}} \frac{1}{\omega_k^{2s}} F_k(\mathbf{x}) \overline{F_k}(\mathbf{y})$; in this way we infer

$$\begin{aligned} \mathfrak{E}^u = & -(1-u) \kappa^u \int_{\partial\Omega} da(\mathbf{x}) \, n^j(\mathbf{x}) \mathfrak{S}^i(\mathbf{x}) \cdot \\ & \cdot \left[\left(-\frac{1}{4} \delta_{ij} \partial^{x^\ell} \partial_{y^\ell} + \frac{1}{2} \partial_{x^i y^j} \right) D_{\frac{u+1}{2}}(\mathbf{x}, \mathbf{y}) \right]_{\mathbf{y}=\mathbf{x}}. \end{aligned} \quad (\text{F.11})$$

Now, comparing the above result with Eq. (4.25) for the regularized pressure we see that

$$\mathfrak{E}^u = -(1-u) \int_{\partial\Omega} da(\mathbf{x}) \, \mathfrak{S}^i(\mathbf{x}) p_i^u(\mathbf{x}); \quad (\text{F.12})$$

the first equality in (F.8) and Eq. (F.12) give the thesis (4.29).

G Appendix. Derivation of Eq. (5.21)

Consider the framework developed in subsection 5.2 for a fundamental operator $\mathcal{A}$, such that $\sigma(\mathcal{A}) \subset [0, +\infty)$ and 0 is in the continuous spectrum of $\mathcal{A}$. Hereafter we are using the deformed fundamental operator

$$\mathcal{A}_\varepsilon := (\sqrt{\mathcal{A}} + \varepsilon)^2. \quad (\text{G.1})$$

⁴¹See again the footnote 21 on page 39.

We already observed in subsection 5.2 (see Eq. (5.17)) that the cylinder kernels T and T^ε , respectively associated to $\mathcal{A}$ and $\mathcal{A}_\varepsilon$, are related by

$$T^\varepsilon(\mathbf{t}; \mathbf{x}, \mathbf{y}) = e^{-\varepsilon \mathbf{t}} T(\mathbf{t}; \mathbf{x}, \mathbf{y}) ; \quad (\text{G.2})$$

we also showed (see Eq. (5.18)) that, assuming the map $\mathbf{t} \mapsto T(\mathbf{t}; \mathbf{x}, \mathbf{y})$ (for fixed $\mathbf{x}, \mathbf{y} \in \Omega$) to admit an extension in a neighborhood of the real half-axis $[0, +\infty)$ to a meromorphic function of $\mathbf{t}$, we have

$$D_s^\varepsilon(\mathbf{x}, \mathbf{y}) = \frac{e^{-2i\pi s} \Gamma(1-2s)}{2\pi i} \int_{\mathfrak{J}} d\mathbf{t} \, \mathbf{t}^{2s-1} T^\varepsilon(\mathbf{t}; \mathbf{x}, \mathbf{y}) . \quad (\text{G.3})$$

The integral in the right-hand side of the above equation is an analytic function of s on the whole complex plane, while the gamma function is meromorphic with simple poles at positive half-integer values of s . Taking into account these facts, hereafter we show how to evaluate the renormalized kernels

$$D_{s_0}^{(\kappa)}(\mathbf{x}, \mathbf{y}) := \lim_{\varepsilon \rightarrow 0^+} RP \Big|_{s=s_0} \left(k^{2(s-s_0)} D_s^\varepsilon(\mathbf{x}, \mathbf{y}) \right) \quad (\text{G.4})$$

considering, separately, the cases $s_0 = -n/2$ ($n \in \{0, 1, 2, \dots\}$) and $s_0 = n/2$ ($n \in \{1, 2, 3, \dots\}$). Putting together the results obtained for $s_0 = -n/2$ ($n \in \{0, 1, 2, \dots\}$) and for $s_0 = 1/2$, we will finally obtain the proof of Eq. (5.21).

We remark that, for $s_0 = \pm 1/2$, the above renormalized kernels coincide with the functions introduced in Eq. (5.9).

Case 1: $s_0 = -\frac{n}{2}$, $n \in \{0, 1, 2, \dots\}$. The right-hand side of Eq. (G.3) is clearly an analytic function of s for $\Re s < \frac{1}{2}$; thus, for $n \in \{0, 1, 2, \dots\}$ $D_s^\varepsilon(\mathbf{x}, \mathbf{y})$ has an analytic continuation at $s = -n/2$, hereafter indicated with $D_{-n/2}^\varepsilon(\mathbf{x}, \mathbf{y})$, that is simply obtained substituting this value of s in the integral representation (G.3). The resulting integrand is meromorphic so that, by the residue theorem,

$$D_{-\frac{n}{2}}^\varepsilon(\mathbf{x}, \mathbf{y}) = (-1)^n \Gamma(n+1) \text{Res} \left(\mathbf{t}^{-(n+1)} e^{-\varepsilon \mathbf{t}} T(\mathbf{t}; \mathbf{x}, \mathbf{y}) ; 0 \right) . \quad (\text{G.5})$$

Of course, in the present case the prescription of taking the regular part in Eq. (5.9) is pleonastic, and the cited equation is reduced to

$$D_{-\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) = \lim_{\varepsilon \rightarrow 0^+} D_{-\frac{n}{2}}^\varepsilon(\mathbf{x}, \mathbf{y}) . \quad (\text{G.6})$$

On the other hand, computation of the previous limit gives ⁽⁴²⁾

$$D_{-\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) = (-1)^n \Gamma(n+1) \operatorname{Res}\left(\mathbf{t}^{-(n+1)} T(\mathbf{t}; \mathbf{x}, \mathbf{y}); 0\right). \quad (\text{G.7})$$

The above result can be reformulated in terms of the modified cylinder kernel $\tilde{T}$ associated to $\mathcal{A}$. Indeed, recall that $T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = -\partial_{\mathbf{t}} \tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y})$ (see Eq. (3.54)) and note that, for any pair of functions f, g meromorphic near a point $\mathbf{t}_0$, we have $\operatorname{Res}(fg'; \mathbf{t}_0) = -\operatorname{Res}(f'g; \mathbf{t}_0)$; these facts (and the standard identity $z \Gamma(z) = \Gamma(1+z)$) give

$$D_{-\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) = (-1)^{n+1} \Gamma(n+2) \operatorname{Res}\left(\mathbf{t}^{-(n+2)} \tilde{T}(\mathbf{t}; \mathbf{x}, \mathbf{y}); 0\right). \quad (\text{G.8})$$

Case 2: $s_0 = +\frac{n}{2}$, $n \in \{1, 2, 3, \dots\}$ (with a special attention for the subcase $n = 1$). A substantial difference occurs with respect to Case 1. In fact, due to the gamma function appearing in Eq. (G.3), the function $D_s^\varepsilon(\mathbf{x}, \mathbf{y})$ described by this equation has a genuine singularity at $s = n/2$; in order to remove this singularity, it is essential to retain only the regular part in Eq. (5.9). To this purpose, for s in a neighborhood of $n/2$, we introduce the variable $u := 2s - n$ and note that, for $u \rightarrow 0$,

$$\begin{aligned} & \kappa^{2s-n} e^{-2i\pi s} \Gamma(1-2s) \mathbf{t}^{2s-1} = \\ & = \frac{\mathbf{t}^{n-1}}{(n-1)!} \left[\frac{1}{u} + \left(\ln(\kappa \mathbf{t}) + \gamma_{EM} - i\pi - H_{n-1} \right) + O(u) \right], \end{aligned} \quad (\text{G.9})$$

where $\gamma_{EM} \simeq 0.577216$ is the Euler-Mascheroni constant and, for $m \in \{0, 1, 2, \dots\}$, $H_m := \sum_{k=1}^m \frac{1}{k}$ ($H_0 := 0$) denotes the m -th harmonic number ⁽⁴³⁾.

⁴²To prove this, one can proceed as follows. In the present case the cylinder kernel is assumed to be a meromorphic function of $\mathbf{t}$, so there is an expansion $T(\mathbf{t}; \mathbf{x}, \mathbf{y}) = \frac{1}{\mathbf{t}^q} \sum_{k=0}^{+\infty} e_k(\mathbf{x}, \mathbf{y}) \mathbf{t}^k$ for some $q \in \mathbf{Z}$ (converging, at least, for small $\mathbf{t}$; note that under the assumptions for the validity of (3.64) we have $q = d$); of course $e^{-\varepsilon \mathbf{t}} = \sum_{k=0}^{+\infty} \frac{(-\varepsilon)^k}{k!} \mathbf{t}^k$, so the Cauchy formula for the product of two series yields

$$\operatorname{Res}\left(\mathbf{t}^{-(n+1)} e^{-\varepsilon \mathbf{t}} T(\mathbf{t}; \mathbf{x}, \mathbf{y}); 0\right) = \sum_{k=0}^{q+n} \frac{(-\varepsilon)^k}{k!} e_{q+n-k}(\mathbf{x}, \mathbf{y}) \xrightarrow{\varepsilon \rightarrow 0} e_{d+n}(\mathbf{x}, \mathbf{y}) = \operatorname{Res}\left(\mathbf{t}^{-(n+1)} T(\mathbf{t}; \mathbf{x}, \mathbf{y}); 0\right).$$

This proves Eq. (G.7).

⁴³ To obtain Eq. (G.9), one uses the following well known facts [80]: for $m \in \{0, 1, 2, \dots\}$,

$$\Gamma(-u-m) = \frac{(-1)^m \Gamma(-u)}{(u+1)\dots(u+m)}; \quad (u+1)\dots(u+m) = m! \left[1 + H_m u + O(u^2) \right],$$

$$\Gamma(-u) = -\frac{1}{u} - \gamma + O(u), \quad e^{-i\pi u} (\kappa \mathbf{t})^u = e^{u(\ln(\kappa \mathbf{t}) - i\pi)} = 1 + (\ln(\kappa \mathbf{t}) - i\pi)u + O(u^2) \quad \text{for } u \rightarrow 0.$$

It follows that

$$\begin{aligned} RP \Big|_{s=\frac{n}{2}} \left(\kappa^{2s-n} D_s^\varepsilon(\mathbf{x}, \mathbf{y}) \right) = \\ = \frac{1}{2\pi i} \int_{\mathfrak{H}} d\mathfrak{t} \frac{\mathfrak{t}^{n-1} e^{-\varepsilon \mathfrak{t}}}{(n-1)!} \left(\ln(\kappa \mathfrak{t}) + \gamma - i\pi - H_{n-1} \right) T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) . \end{aligned} \quad (\text{G.10})$$

According to Eq. (5.9), the renormalized function $D_{\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y})$ is the limit $\varepsilon \rightarrow 0^+$ of the above expression. Under suitable hypotheses on the behaviour of T for $\Re \mathfrak{t} \rightarrow +\infty$ (namely, $|T(\mathfrak{t}; \mathbf{x}, \mathbf{y})| \leq C |\mathfrak{t}|^{-a-n}$ for some $C, a > 0$), we can exchange the limit and the integral to obtain

$$D_{\frac{n}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) = \frac{1}{2\pi i} \int_{\mathfrak{H}} d\mathfrak{t} \frac{\mathfrak{t}^{n-1}}{(n-1)!} \left(\ln(\kappa \mathfrak{t}) + \gamma - i\pi - H_{n-1} \right) T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; \quad (\text{G.11})$$

the term $\ln(\kappa \mathfrak{t})$ prevents us from using the residue theorem, so we must find alternative ways to evaluate explicitly the above integral. In the special case $n = 1$, we can proceed as follows. First we recall that $T(\mathfrak{t}; \mathbf{x}, \mathbf{y}) = -\partial_{\mathfrak{t}} \tilde{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y})$ and integrate by parts Eq. (G.11) to obtain

$$D_{\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) = \frac{1}{2\pi i} \int_{\mathfrak{H}} d\mathfrak{t} \mathfrak{t}^{-1} \tilde{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; \quad (\text{G.12})$$

the resulting integrand is meromorphic in $\mathfrak{t}$ so that we can resort to the residue theorem to obtain

$$D_{\frac{1}{2}}^{(\kappa)}(\mathbf{x}, \mathbf{y}) = \text{Res} \left(\mathfrak{t}^{-1} \tilde{T}(\mathfrak{t}; \mathbf{x}, \mathbf{y}) ; 0 \right) . \quad (\text{G.13})$$

Conclusion. Eq. (G.8) for $D_{-\frac{n}{2}}^{(\kappa)}$ with $n \in \{0, 1, 2, \dots\}$ and Eq. (G.13) for $D_{\frac{1}{2}}^{(\kappa)}$ prove Eq. (5.21) for $D_{-\frac{n}{2}}^{(\kappa)}$ with $n \in \{-1, 0, 1, 2, \dots\}$.

References

- [1] A.A. Actor, *Multiple harmonic oscillator zeta functions*, J.Phys.A:Math.Gen. **20**, 927–936 (1987).
- [2] A.A. Actor, *Local analysis of a quantum field confined within a rectangular cavity*, Ann.Phys. **230**, 303–320 (1994).
- [3] A.A. Actor, *Scalar quantum fields confined by rectangular boundaries*, Fortsch.Phys. **43**, 141–205 (1995).
- [4] A.A. Actor, I. Bender, *Boundaries Immersed in a Scalar Quantum Field*, Fortsch.Phys. **44**(4), 281–322 (1996).
- [5] R.A. Adams, J.J.F. Fournier, “Sobolev Spaces”, Academic Press (2003).
- [6] S. Albeverio, G. Cognola, M. Spreafico, S. Zerbini, *Singular perturbations with boundary conditions and the Casimir effect in the half space*, J.Math.Phys. **51**(06), 063502 (2010).
- [7] N. Aronszajn, K.T. Smith, *Theory of Bessel potentials. I*, Ann.Inst.Fourier (Grenoble) **11**, 385–475 (1961).
- [8] M. Benini, C. Dappiaggi, T.P. Hack, *Quantum Field Theory on Curved Backgrounds – A Primer*, Int.Jour.Mod.Phys.A **28**, 1330023 (2013).
- [9] I.M. Berezanskii, “Selfadjoint Operators in Spaces of Functions of Infinitely Many Variables”, Am.Math.Soc. (1986).
- [10] J. Bergh and J. Löfström. *Interpolation Spaces*, Springer (1976).
- [11] N. Berline, E. Getzler, M. Vergne, “Heat Kernels and Dirac Operators”, Springer Science & Business Media (1992).
- [12] N.D. Birrell, P.C.W. Davies, “Quantum Fields in Curved Space”, Cambridge University Press (1984).
- [13] S.K. Blau, M. Visser, A. Wipf, *Zeta functions and the Casimir energy*, Nucl.Phys.B **310**(1), 163–180 (1988).
- [14] M. Bordag, U. Mohideen, V. M. Mostepanenko, *New developments in the Casimir effect*, Phys.Rep.-Rev.Sect.Phys.Lett. **353**(1), 1–205 (2001).
- [15] M. Bordag, G.L. Klimchitskaya, U. Mohideen, V.M. Mostepanenko, “Advances in the Casimir Effect”, Oxford University Press (2009).

- [16] S.T. Bunch, P.C.W. Davies, *Quantum field theory in de Sitter space: renormalization by point-splitting*, Proc.R.Soc.Lond.A Math.Phys.Sc. **360**(1700), 117–134 (1978).
- [17] A. A. Bytsenko, G. Cognola, E. Elizalde, V. Moretti, S. Zerbini, “Analytic aspects of quantum fields”, World Scientific (2003).
- [18] O. Calin, D.-C. Chang, K. Furutani, C. Iwasaki, “Heat Kernels for Elliptic and Sub-elliptic Operators: Methods and Techniques”, Springer Science & Business Media (2010).
- [19] C.G. Callan, S.R. Coleman, R. Jackiw, *A new improved energy-momentum tensor*, Ann.Phys. **59**, 42–73, (1970).
- [20] H.B.G. Casimir, *On the attraction between two perfectly conducting plates*, Proc.R.Neth.Acad.Arts Sci. **51**, 793–795 (1948).
- [21] I. Cavero-Peláez, K.A. Milton, K. Kirsten, *Local and global Casimir energies for a semitransparent cylindrical shell*, J.Phys.A-Math.Theor. **40**(13), 133607 (2007).
- [22] I. Chavel, “Eigenvalues in Riemannian Geometry”, Academic Press (1984).
- [23] J. Chazarain, A. Piriou, “Introduction to the Theory of Linear Partial Differential Equations”, Elsevier (2011).
- [24] G. Cognola, L. Vanzo, and S. Zerbini, *A New Algorithm for Asymptotic Heat Kernel Expansion for Manifolds with Boundary*, Phys.Lett.B **241**(3), 381–386 (1990).
- [25] G. Cognola, E. Elizalde, S. Zerbini, *Fluctuations of quantum fields via zeta function regularization*, Phys.Rev.D **65**, 085031 (2002).
- [26] G. Cognola, S. Zerbini, *Variances of relativistic quantum field fluctuations*, Int.J.Mod.Phys.A **18**(12), 2067–2072 (2003).
- [27] C. Dappiaggi, G. Nosari, N. Pinamonti, *The Casimir effect from the point of view of algebraic quantum field theory*, arXiv:1412.1409 (2014).
- [28] E.B. Davies, “Heat Kernels and Spectral Theory”, Cambridge University Press (1990).
- [29] D. Deutsch, P. Candelas, *Boundary effects in quantum field theory*, Phys.Rev.D **20**(12), 3063–3080 (1979).

- [30] J.S. Dowker and R. Critchley, *Effective Lagrangian and energy-momentum tensor in de Sitter space*, Phys.Rev.D **13**, 3224–3232 (1976).
- [31] D.G. Duffy, “Green’s Functions with Applications”, CRC Press (2001).
- [32] Yu. V. Egorov, M.A. Shubin, *Linear Partial Differential Equations* in “Elements of the Modern Theory of Partial Differential Equations”, Springer-Verlag (1999).
- [33] E. Elizalde, S.D. Odintsov, A. Romeo, A.A. Bytsenko, S. Zerbini, “Zeta regularization techniques with applications”, World Scientific (1994).
- [34] E. Elizalde, “Ten Physical Applications of Spectral Zeta Functions”, Springer Science & Business Media (1995).
- [35] R. Estrada, S.A. Fulling, *Distributional asymptotic expansions of spectral functions and of the associated Green kernels*, Elect.J.Diff.Eqs. **1999**, 1–37 (1999).
- [36] D. Fermi, L. Pizzocchero, *Local zeta regularization and the Casimir effect*, Prog.Theor.Phys. **126**(3), 419–434 (2011); see also arXiv:1104.4330.
- [37] D. Fermi, *L’effetto Casimir e la regolarizzazione zeta*, Master Thesis, University of Milan (2012).
- [38] D. Fermi, L. Pizzocchero, *Local zeta regularization and the scalar Casimir effect II. Some explicitly solvable cases*, arXiv:1505.01044 (2015).
- [39] D. Fermi, L. Pizzocchero, *Local zeta regularization and the scalar Casimir effect III. The case with a background harmonic potential*, arXiv:1505.01651 (2015).
- [40] D. Fermi, L. Pizzocchero, *Local zeta regularization and the scalar Casimir effect IV. The case of a rectangular box*, arXiv:1505.03276 (2015).
- [41] D. Fermi, L. Pizzocchero, In preparation.
- [42] P. Flajolet, X. Gourdon, P. Dumas, *Mellin transforms and asymptotics: Harmonic sums*, Theor.Comp.Sc. **144**, 3–58 (1995).
- [43] L.H. Ford, N.F. Svaiter, *Vacuum energy density near fluctuating boundaries*, Phys.Rev.D **58**, 065007 (1998).
- [44] G. Fucci, K. Kirsten, *Conical Casimir pistons with hybrid boundary conditions*, J.Phys.A-Math.Theor. **44**(29), 295403 (2011).

- [45] G. Fucci, K. Kirsten, *The Spectral Zeta Function for Laplace Operators on Warped Product Manifolds of the type $I \times f_N$* , Comm.Math.Phys **317**(3), 635-665 (2013).
- [46] S.A. Fulling, “Aspects of quantum field theory in curved space time”, Cambridge University Press (1989).
- [47] S.A. Fulling, R.A. Gustafson, *Some properties of Riesz means and spectral expansions*, Elect.J.Diff.Eqs., **1999**(6), 1–39 (1999).
- [48] S.A. Fulling, *Systematics of the relationship between vacuum energy calculations and heat kernel coefficients*, J.Phys.A: Math.Gen. **36**, 6857–6873 (2003).
- [49] S.A. Fulling, *Mass dependence of vacuum energy*, Phys.Lett.B **624**, 281–286 (2005).
- [50] S.A. Fulling, *Vacuum Energy as Spectral Geometry*, SIGMA **3**, 094 (2007).
- [51] S.A. Fulling, C.S. Trendafilova, P.N. Truong, J. Wagner, *Wedges, cones, cosmic strings and their vacuum energy*, J.Phys.A: Math.Theor. **45**, 374018 (2012).
- [52] I.M. Gel’fand, G.E. Shilov, “Generalized Functions. Volume III: Theory of Differential Equations”, Academic Press (1967).
- [53] P.B. Gilkey, “Invariance Theory: The Heat Equation and the Atiyah-Singer Index Theorem”, CRC Press (1994).
- [54] A. Grigor’yan, “Heat Kernel and Analysis on Manifolds”, AMS/IP Studies in Advanced Mathematics **47** (2009).
- [55] T.P. Hack, V. Moretti, *On the stress–energy tensor of quantum fields in curved spacetimes – comparison of different regularization schemes and symmetry of the Hadamard/SeeleyDeWitt coefficients*, J.Phys.A:Math.Theor. **45**, 374019 (2012).
- [56] G.H. Hardy, M. Riesz, “The general theory of Dirichlet’s series”, Cambridge Tracts in Mathematics **18**, Cambridge University Press (1915).
- [57] J.M. Harrison, K. Kirsten, C. Texier, *Spectral determinants and zeta functions of Schrödinger operators on metric graphs*, J.Phys.A-Math.Theor. **45**(12), 125206 (2011).
- [58] S.W. Hawking, *Zeta function regularization of path integrals in curved space-time*, Commun. Math. Phys. **55**(2), 133–148 (1977).

- [59] P. Henrici, “Applied and computational complex analysis. Volume I”, Wiley and Sons (1974).
- [60] D. Iellici, V. Moretti, *ζ -function regularization and one-loop renormalization of field fluctuations in curved space-time*, Phys.Lett.B **425**, 33–40 (1998).
- [61] V.A. Il’in, I.A. Shishmarev, *Uniform estimates in a closed domain for the eigenfunctions of an elliptic operator and their derivatives*, Izv.Akad.Nauk SSSR.Set.Mat. **24**(6), 883–896 (1960).
- [62] K. Kirsten, “Spectral Functions in Mathematics and Physics”, CRC Press (2001).
- [63] K. Kirsten, *Basic zeta functions and some applications in physics* in “A Window into Zeta and Modular Physics”, Mat.Sc.Res.Inst.Pub. **57**, Cambridge University Press (2010).
- [64] N.V. Krylov, “Lectures on Elliptic and Parabolic Equations in Sobolev Spaces”, American Mathematical Soc. (2008).
- [65] P.K. Kythe, “Green’s Functions and Linear Differential Equations: Theory, Applications, and Computation”, CRC Press (2011).
- [66] T. Lu, T. Jeffres, K. Kirsten, *Zeta function of self-adjoint operators on surfaces of revolution*, J.Phys.A-Math.Theor. **48**(14), 145204 (2015).
- [67] A. Lunardi, *Interpolation Theory. Appunti*, Scuola Normale Superiore, Pisa (1999).
- [68] S.G. Mamaev, N.N. Trunov, *Vacuum means of energy-momentum tensor of quantized fields on manifolds of different topology and geometry. I*, Sov.Phys.J. **22**(7), 766–770 (1979).
- [69] F. Mandl, G. Shaw, “Quantum Field Theory”, John Wiley & Sons (2013).
- [70] F. D. Mera, S. A. Fulling, *Vacuum energy density and pressure of a massive scalar field*, J. Phys. A: Math. Theor. **48**, 245402 (18pp) (2015).
- [71] V.P. Mikhailov, “Partial differential equations”, Mir Publishers (1978).
- [72] K.A. Milton, “The Casimir effect - Physical manifestations of zero-point energy”, World Scientific Publishing Co.Pte.Ltd. (2001).
- [73] S. Minakshisundaram, *A generalization of Epstein zeta functions*, Canad.J.Math. **1**, 320–327 (1949).

- [74] S. Minakshisundaram, A. Pleijel, *Some properties of the eigenfunctions of the Laplace-Operator on Riemannian manifolds*, Canad.J.Math. **1**, 242–256 (1949).
- [75] V. Moretti, *Direct ζ -function approach and renormalization of one-loop stress tensors in curved spacetimes*, Phys.Rev.D **56**(12), 7797–7819 (1997).
- [76] V. Moretti, *Local ζ -function techniques vs point-splitting procedure: a few rigorous results*, Commun.Math.Phys. **201**, 327–363 (1999).
- [77] V. Moretti, *A review on recent results of the ζ -function regularization procedure in curved spacetime*, in D. Fortunato, M. Francaviglia, A. Masiello, “Recent developments in General Relativity”, Springer-Verlag (1999); also see arXiv:gr-qc/9902056.
- [78] V. Moretti, *Local ζ -functions, stress-energy tensor, field fluctuations, and all that, in curved static spacetime*, Springer Proc.Phys. **137**, 323–332 (2011).
- [79] H.C. Ohanian, *Gravitation and the new improved energy-momentum tensor*, J.Math.Phys. **14**(12), 1892–1897 (1973).
- [80] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, C.W. Clark, “NIST Handbook of mathematical functions”, Cambridge University Press (2010).
- [81] L.E. Parker, D.J. Toms, “Quantum field theory in curved spacetime: Quantized Fields and Gravity”, Cambridge University Press (2009).
- [82] G. Plunien, B. Müller, W. Greiner, *The Casimir Effect*, Phys.Rep. **134**(2,3), 87–193 (1986).
- [83] M.J. Radzikowski, *Micro-local approach to the Hadamard condition in quantum field theory on curved space-time*, Comm.Math.Phys. **179**(3), 529–553 (1996).
- [84] M. Reed, B. Simon, “Methods of Modern Mathematical Physics I: Functional Analysis”, Academic Press (1980).
- [85] F. Rellich, “Perturbation Theory of Eigenvalue Problems”, CRC Press (1969).
- [86] R.B. Rodrigues, N.F. Svaiter, *Vacuum Polarization of a Scalar Field in a Rectangular Waveguide*, arXiv:hep-th/0111131 (2001).
- [87] R.B. Rodrigues, N.F. Svaiter, R.D.M. De Paola, *Vacuum Stress Tensor of a Scalar Field in a Rectangular Waveguide*, arXiv:hep-th/0110290 (2001).

- [88] J.R. Ruggiero, A. Villani, A.H. Zimerman, *Application of Analytic Regularization to the Casimir Forces*, Rev.Bras.Fis. **7**(3), 663–687 (1977).
- [89] J.R. Ruggiero, A. Villani, A.H. Zimerman, *Some comments on the application of analytic regularisation to the Casimir forces*, J.Phys.A:Math.Gen. **13**, 761–766 (1980).
- [90] F. Sauvigny, “Partial Differential Equations 2: Functional Analytic Methods”, Springer Science & Business Media (2012).
- [91] J. Schlemmer, J. Zahn, *The current density in quantum electrodynamics in external potentials*, arXiv:1501.05912 (2015).
- [92] L. Schwartz, “Théorie des noyaux” , Proc. Internat. Congress Mathematicians (Cambridge, 1950) **1**, 220–230, Amer. Math. Soc. (1952), .
- [93] S.S. Schweber, “An Introduction to Relativistic Quantum Field Theory”, Courier Corporation (2005).
- [94] J. Schwinger, “Particles, sources, and fields”, Perseus Books (1998).
- [95] R.T. Seeley, *Complex Powers of an Elliptic Operator*, AMS Proc.Symp.Pure Math. **X**, 288–307 (1967).
- [96] N. Shimakura, “Partial Differential Operators of Elliptic Type”, American Mathematical Soc. (1992).
- [97] I.A. Shishmarev, *Uniform estimates for the derivatives of solutions to the Dirichlet problem and the eigenvalue problem for the operator $Lu - \operatorname{div}(p(x) \operatorname{grad} u) + q(x)u$ with discontinuous coefficients*, Dokl.Akad.Nauk SSSR **137**(1), 45–47 (1961); see also, Soviet Math.Dokl. **2**(2), 244–246 (1961).
- [98] I.N. Sneddon, “The use of integral transforms”, McGraw-Hill (1972).
- [99] A. Sommerfeld, “Partial differential equations in physics”, Academic Press (1949).
- [100] I. Stakgold, M.J. Holst, “Green’s Functions and Boundary Value Problems”, John Wiley & Sons (2011).
- [101] R.M. Wald, *On the Euclidean approach to quantum field theory in curved spacetime*, Commun. Math. Phys. **70**(3), 221–242 (1979).
- [102] R.M. Wald, “General relativity”, University of Chicago Press (2010).

- [103] G.N. Watson, “A Treatise on the Theory of Bessel Functions”, Cambridge University Press (1995).
- [104] V.Ya. Yakubov, *Estimates for Eigenfunctions of Elliptic Operators with Respect to the Spectral Parameter*, Func.Anal.App. **33**(2), 128–136 (1999).